

Kinetic Products Under Thermal Conditions: Rapid Entry into α/β -D-Galactofuranosides Using Microwave Irradiation and Selective Lewis Acids

Soumava Santra , Emily Jonas , Jean-Paul Bourgault , Tarick El-Baba & Peter R. Andreana

To cite this article: Soumava Santra , Emily Jonas , Jean-Paul Bourgault , Tarick El-Baba & Peter R. Andreana (2011) Kinetic Products Under Thermal Conditions: Rapid Entry into α/β -D-Galactofuranosides Using Microwave Irradiation and Selective Lewis Acids, Journal of Carbohydrate Chemistry, 30:1, 27-40, DOI: [10.1080/07328303.2011.585259](https://doi.org/10.1080/07328303.2011.585259)

To link to this article: <http://dx.doi.org/10.1080/07328303.2011.585259>



Published online: 27 Jun 2011.



Submit your article to this journal [↗](#)



Article views: 212



View related articles [↗](#)



Citing articles: 2 View citing articles [↗](#)

Kinetic Products Under Thermal Conditions: Rapid Entry into α/β -D-Galactofuranosides Using Microwave Irradiation and Selective Lewis Acids

Soumava Santra, Emily Jonas, Jean-Paul Bourgault,
Tarick El-Baba, and Peter R. Andreana

Department of Chemistry, Wayne State University, Detroit, Michigan, USA 48202

A modified Fischer-Lubineau reaction, employing microwave irradiation and selected Lewis acids such as $\text{Mn}(\text{ClO}_4)_2$, $\text{Mn}(\text{C}_2\text{H}_3\text{O}_2)_3$, FeCl_3 , CoCl_2 , and AgF as independent additives, induces rapid entry into α/β -D-galactofuranosides in very good yield and purity.

Keywords D-Galactofuranosides; Lewis acid coordination; Zwitterionic polysaccharide PS A1; Microwave irradiation; M,P arrangement; Fisher-Lubineau reaction

INTRODUCTION

D-Galactofuranosides (D-Galf) have been identified and well characterized in oligosaccharides and glycoconjugates existing within many infectious organisms including *Mycobacterium tuberculosis*, *Mycobacterium leprae*, *Escherichia coli*, *Trypanosoma cruzi*, *Leishmania*, *Paracoccidioides brasiliensis*, and *Bacteroides fragilis*, to name a few.^[1] Considerable attention to the constituents of these pathogens has revealed that their survival and virulence is directly reliant on D-galactofuranoside moieties. These sugars have mostly been found as terminal, nonreducing units such as in the CD4^+ T-cell immunostimulatory capsular zwitterionic polysaccharide PS A1 (**1**) (Fig. 1).^[2]

Ironically, D-galactofuranoside is nonexistent in mammalian tissue and has been shown to be an antigenic determinant representing a valuable

Received October 18, 2010; accepted April 28, 2011.

Address correspondence to Peter R. Andreana, Department of Chemistry, Wayne State University, Detroit, Michigan, USA 48202. E-mail: pra@chem.wayne.edu





Figure 1: Immunostimulatory zwitterionic polysaccharide PS A1 (**1**) from *Bacteroides fragilis* NCTC 9343 containing a terminal D-Galf unit.

target for new drug agents and therapies.^[3–6] Although of immense importance, our current biological understanding of many aspects of D-Galf, such as its metabolic processes, have proven difficult to elucidate due to complications in obtaining derivatives.^[7]

One drawback is that unlike their thermodynamically favorable pyranoside counterparts, hexofuranosides are not commercially available. Equilibration and anomeric control of both hexo-pyranoside and hexo-furanoside syntheses must be considered in addition to the competing donor/acceptor reactions for its construction. To this extent, pioneering work by H.E. Fischer from 1893 to 1895^[8–11] demonstrated that furanosides could be obtained from corresponding unprotected pyranoses using simple alcohols and Brønsted-Lowry acids.

Further studies by J.C. Fischer and A. Lubineau^[12] established that when unprotected monosaccharides were subjected to thermal heating in the presence of anhydrous Lewis acid FeCl_3 and methanol, the reaction yielded kinetically favored methyl furanoside (**3**) as the major product in a mixture of α - and β -anomers (Sch. 1). The reaction took approximately 28 h and in addition the purification proved to be challenging and cumbersome.



Scheme 1: Fischer-Lubineau D-Galf synthesis.


Ferrières and Plusquellec have also utilized FeCl_3 as a promoter but with CaCl_2 as an additive in THF to obtain D-Galf.^[13–16] Although other methods have been reported, including base-mediated furanoside formation,^[17,18] reduction of lactones,^[19] acid-catalyzed ring opening of 1,4-anhydrosugars,^[20] high-temperature benzylation,^[21] cyclization with mercuric salts,^[22,23] and the utilization of dithioacetal derivatives of galactose,^[24–26] the chemical synthesis of D-Galf remains to be an important yet challenging endeavor.

RESULTS AND DISCUSSION

We elected to revisit the Fischer-Lubineau D-Galf synthesis and rationalize the use of microwave irradiation as a thermal source for kinetic product formation. As our current research program focuses on chemical modifications of D-Galf in ZPS PS A1 (**1**) for immunogen development,^[27] having a reliable means to obtain D-galactofuranoside in very good yield and high purity is of great importance.

Since the spectroscopic data of methyl α/β -D-galacto-furanoside and -pyranoside (D-Galp) was known,^[12,16,28,29] methanol was chosen as the initial acceptor for direct comparison. Therefore, when unprotected D-galactose (**2**) was dissolved in anhydrous methanol and subjected to microwave irradiation in the presence of 1 mole equivalent anhydrous FeCl_3 for 5 min at stage 1 (microwave was equilibrated to 300 W, 60°C, 4 bar) and then for 5 min at stage 2 (microwave was equilibrated to 300 W, 70°C, 5 bar), the overall conversion was 35% to 45% (Table 1, entries 1 and 2). In realizing that an extended incubation

Table 1: Optimization of methyl D-galactofuranoside synthesis using microwave irradiation and anhydrous FeCl_3 ^a



Entry	Stage 1		Stage 2		% Yield ^b / α : β ^c		
	Temp (°C)	Time (min)	Temp (°C)	Time (min)	D-Galf (3)	D-Galp (4)	Overall % yield
1	60	5	70	5	25 / 1:2	5 / 2:1	35
2	60	10	70	5	35 / 1:3	10 / 2:1	45
3	60	20	70	15	60 / 1:5	15 / 3:1	75
4	60	30	70	25	70 / 1:7	17 / 5:1	87
5	60	30	70	35	75 / 1:8	20 / 6:1	95

^aCEM Discover Microwave; microwave power was set to 300 W.

^bIsolated.


^cDetermined from ^1H NMR sample of unpurified reaction mixture; chemical shifts of anomers compared to literature precedent.

period would have a bearing on the overall conversion (Table 1, entries 3 and 4), further optimization of the dual-stage time course provided 75% yield of the desired D-Galf with an α/β selectivity of 1:8 (Table 1, entry 5). It is important to note that allowing the reaction to run in excess of 70 min or increasing the temperature in stage 2 led to decreased yields of methyl D-Galf and in some cases decomposition. The microwave-assisted efforts yielded an improvement in reaction rate, from the reported 240 min to 65 min, and we noted an improvement in yield and α/β selectivity of methyl D-galactofuranoside.

With the described protocol in hand, we next turned our attention to examining the Lewis acid/additive scope. A literature search revealed that carbohydrates and derivatives thereof are known to form complexes with metal cations.^[30,31] Furthermore, experimental evidence has shown that the outcomes in acid-mediated glycosylation may influence the α/β and D-Galf/D-Galp selectivities.^[31,32]

From our data we observed that Lewis acids, in which the central atom is in +I, +III, or +IV oxidation state (Table 2, entries 1, 2, 6, 8, 11, and 13–16), provided excellent overall yield of the glycoside. However, when Zn(II), Cu(I), and Cu(II) (Table 2, entries 3–5) were used, there was no observed product. Use of Amberlyst 15 as an additive gave the methyl α -D-galactopyranoside (**4**) in very good yield and high α -selectivity while TMSOTf did not lead to product formation (or at best a trace amount) (Table 2, entries 17 and 18). Remarkably, only FeCl₃ and AgF (Table 2, entries 8 and 12) gave the best overall yields and α/β selectivities of methyl D-Galf. Interestingly, the α/β ratio of D-Galf was compromised when MnCl₂, and Mn(ClO₄)₂ was used, whereas Mn(C₂H₃O₂)₃ provided only methyl α -D-Galf in a mixture with α/β -D-Galp (Table 2, entries 9–11). On the other hand, CoCl₂ (Table 2, entry 7) was selective for α -D-Galf. It is worth noting that all the manganese and cobalt salts used (Table 2, entries 7, and 9–11) provided D-Galf as the major product, albeit with an overall decrease in yield as compared to entries 8 and 12. Our data suggest that the oxidation state of the central atom in the Lewis acid does not control α/β selectivities but rather other properties have a direct affect on product outcome.

One likely explanation is that in general, metal cations are known to coordinate with ligands via O-atoms.^[33] A noted steric arrangement in which three O-atoms found on three consecutive C-atoms in such a conformation that the first and second are *gauche*⁺ and the second and third *gauche*[−] or (*threo-threo*) has a profound bearing on metal cation coordination due to no unfavorable interactions.^[34] This has been commonly referred to as the “M,P arrangement.”^[35] Findings suggest that the most common occurrence is a sequence of axial (ax), equatorial (eq), and axial OH groups on chair-formed 6-membered rings (**5**, **6a–b**) or envelope-formed 5-membered rings (**6c–d**) (Fig. 2).^[30] The “M,P arrangement,” in subsequent flipped orientations of the pyranoside or furanoside, can also be noted as an equatorial (eq), axial (ax), and equatorial (eq) convention.

Table 2: Effect of Lewis acids/additives on microwave-assisted methyl D-Galf synthesis^a

Entry	Lewis acid/ additive ^b	Central metal atom oxidation state	Ionic radii (Å) ^c	% Yield ^d /α:β ^e		Overall % yield
				3	4	
1	BF ₃ ·OEt ₂	III	0.87	25/1:5	63/5:1	88
2	AlCl ₃	III	1.18	15/1:7	80/5:1	95
3	ZnCl ₂	II	1.42	0/—	0/—	0
4	CuI	I	1.45	0/—	0/—	0
5	CuSO ₄	II	1.45	0/—	0/—	0
6	InCl ₃	III	1.46	23/1:7	74/5:1	97
7	CoCl ₂	II	1.52	48/2:1	30/5:1	78
8	FeCl ₃	III	1.56	75/1:8	20/6:1	95
9	MnCl ₂	II	1.61	49/1:3	28/2:1	77
10	Mn(ClO ₄) ₂	II	1.61	59/1:1	20/10:1	79
11	Mn(C ₂ H ₃ O ₂) ₃	III	1.61	55/100:1	24/3:1	79
12	AgF	I	1.65	75/1:6	21/6:1	96
13	TiCl ₄	IV	1.76	15/1:4	75/5:1	90
14	Ti(O- <i>i</i> -Pr) ₄	IV	1.76	16/1:4	73/5:1	89
15	Sc(OTf) ₃	III	1.84	25/1:7	70/5:1	95
16	Yb(OTf) ₃	III	2.22	20/1:7	72/5:1	92
17	Amberlyst 15	—	—	15/1:5	75/9:1	90
18	TMSOTf	—	—	0/—	0/—	0

^a CEM Discover Microwave; microwave power was set to 300 W.^b Anhydrous.^c See reference 36.^d Isolated.^e Determined from ¹H NMR of unpurified reaction mixture; chemical shift of anomers compared to literature precedent.

However, coordination of metal cations via a C-1, C-2, C-3 eq-eq-ax conformation, such as in D-glucose (**7**), is highly unfavorable. Importantly, documented studies of complex formation with cations have been carried out using acyclic polyols and they can adapt the “M,P arrangement.”^[34]

In addition, it has also been observed that the radii of metal ions affecting the extent of coordination can shift the equilibrium in favor of α-anomers,^[31] although based on our findings, in conjunction with microwave irradiation, one can expect favorable β-D-Galf outcomes except in cases where CoCl₂ and Mn(C₂H₃O₂)₃ are employed. Importantly, we argue that the ionic radii of Ag, Mn, Fe, and Co (1.65 Å, 1.61 Å, 1.56 Å, and 1.52 Å, respectively)^[36] allow for





Figure 2: Coordination of alkaline-earth metal ions with D-allopyranosides (**5**, **6a,b**) and 3-O-methyl α -D-allofuranose (**6c,d**).^[31]

the best eq-ax-eq coordination at the C-3, C-4, and C-5 hydroxyls. We also believe that the coordination number and outer orbital electron configuration of the central atom are responsible for our observed product yields and conformational forms.

A plausible mechanism for the formation of alkyl D-galacto-furanoside and -pyranoside is shown in Scheme 2. It is well known that D-galactose (**2**) exists in equilibrium with the open-chain form, which can readily undergo isomerization through the C-5 or C-4 hydroxyls. However, in the instances where Lewis acids are used, two metal coordinating paths are highly plausible. In pathway a (when M = Mn, Fe, Co, or Ag), and for formation of **9**, the oxygen atom of the pyranose ring can coordinate to the Lewis acid, which leads to the acyclic aldose sugar in the highly stable eq-ax-eq (3,4,5-erythro-threo or “MP arrangement”) configuration.^[37] The C-4 axial hydroxyl group can then attack the carbonyl aldehyde and lead to the cyclization forming a five-membered furanose ring, which can then undergo dehydration to form the oxocarbenium intermediate **10**. Glycosylation can then occur to render the furanoside as the major kinetic product, which, dependent upon the Lewis acid used, can give rise to either α - or β -anomers (Table 2; entries 8 versus 11 for example) in very good yields.^[38,39]

In pathway b (when M \neq Mn, Fe, Co, or Ag), and for the formation of **12**, metal coordination at the anomeric hydroxyl can lead to the oxocarbenium ion intermediate through the loss of M-OH. Once this intermediate is formed, the anomeric effect is attributed in controlling the orientation of the acceptor as predominantly α . In fact, in all cases we observed the α -anomer dominated, with only a minor amount of β -anomer being characterized.

A notable feature of this microwave-assisted method for furanose formation is that solvent polarity apparently does not influence anomeric ratios as other solvents were tested and yielded similar stereochemical outcomes, albeit at significant loss in overall product yields. In our experiments we believe that




Scheme 2: Proposed mechanism for the Lewis acid-mediated β -galacto-furanoside and -pyranoside formation.

initially methanol will coordinate to the Lewis acid in a monodentate fashion but then be displaced by a more entropically favored tridentate coordination complex such as in **9** due to the chelate effect. Lewis acid coordination to the final D-galactofuranoside is also possible, but the 3,4,5-erythro-threo ($eq\text{-}ax\text{-}eq$) coordination prevails such as in **9**. It is important to consider that in a recent report, it was shown that stereoselective glycosylations of 1-thioimidoyl hexofuranosides could be achieved by adjusting the concentration of promoter $\text{Cu}(\text{OTf})_2$.^[37b] In that report, ^{13}C NMR spectrum was used to illustrate that copper(II) ions could assist in five-membered ring preservation through the complexation of the *O*-5 of the furanosidic derivative. Although another report describing how ultrasound affects bond rotation in the Diels-Alder reaction

exists,^[40] we do not believe, at this point, that microwave irradiation affects bond rotation for selective β -formation in our examples.

In order to probe the reaction scope, we elected to use allyl alcohol and 2-phenylethanol as acceptors in the furanoside synthesis. Both allyl D-galactofuranosides (**15a,b**) and phenylethyl D-galactofuranosides (**16a,b**) were obtained in good yield and anomeric selectivity (Sch. 3).^[41] The minor allyl α -D-galactofuranoside (**15a**) was unequivocally confirmed by x-ray crystal structure analysis.



Scheme 3: Synthesis of allyl- and phenylethyl-D-galactosides.

CONCLUSION

In conclusion, we have successfully employed microwaves in a modified Fischer-Lubineau reaction for alkyl D-galactofuranoside synthesis, which led to an increase in reaction rate and product/anomeric ratios. The reaction scope was examined with acceptors such as allyl- and phenylethyl-alcohols, the aforementioned acceptor yielding an X-ray crystal. Our results also reveal a direct correlation between the properties of Lewis acids and furanoside synthesis. A *C*-3, *C*-4, *C*-5-*erythro-threo* eq-ax-eq Lewis acid complex, such as **9**, led to galactofuranoses as kinetic products under thermal conditions. One can expect further applicable work in this area as we utilize the constructed furanosides as acceptors/donors for oligomer synthesis. Importantly, the allyl β -D-galactofuranoside building block has been utilized to synthesize lipid-bearing β -1,6-linked D-Galf disaccharide as a biological probe.^[42] We also plan to probe oxidative protocols for vicinal diol interconversions leading to novel glycoconjugation methods.

EXPERIMENTAL

Supplementary information (SI) available: NMR spectra for all compounds and X-ray data for **15a**. Contact the corresponding author for this information.

Allyl β -D-galactofuranoside (**15b**) was also synthesized via a Schmidt glycosylation strategy as a reference sample for comparison purposes and for determining α/β ratios.

Materials and Methods

All reagents were purchased from Aldrich unless otherwise noted. Anhydrous ferric chloride was purchased from Alfa Aesar. Anhydrous methanol was purchased from EMD Chemicals (DriSolv). Palladium chloride was purchased from STREM Chemicals. Indium chloride, copper iodide, and copper sulfate were purchased from Alfa Aesar. TMSOTf was purchased from Acros Organics. Allyl alcohol was distilled prior to use. Celite was purchased from Aldrich (catalog no. 221791). Except as otherwise indicated, reactions were carried out under argon. Microwave reactions were conducted using a capped vial on a CEM Discover Microwave System. All reactions were monitored by thin layer chromatography using 0.25-mm Dynamic Adsorbents' precoated silica gel (particle size 0.03–0.07 mm, catalog no. 84111). Column chromatography was performed using Whatman Purasil 60 Å (230–400 mesh ASTM) silica gel. Yields refer to chromatographically and spectroscopically pure compounds, except as otherwise noted. Diastereomeric ratios were determined from ^1H NMR spectra of nonpurified reaction mixtures. Proton and carbon-13 NMR spectra were recorded on Varian Mercury 400, Varian Unity 500, and Varian 500 Direct Drive System spectrometers. The residual CDCl_3 singlet at δ 7.26 ppm and δ 77 ppm was used as the standard for ^1H NMR and ^{13}C NMR spectra, respectively. Mass spectra were recorded on Micromass GCT at 70 eV.

General Method for Microwave-Assisted D-Galactofuranoside Synthesis

A 10-mL CEM Discover microwave vial was oven dried, capped, and cooled under argon. To the vial D-galactose (0.05 g, 0.28 mmol) was added followed by FeCl_3 (Lewis acid) (0.045 g, 0.28 mmol). To the mixture, anhydrous methanol (alcohol) (2.5 mL) was added slowly. The vial was capped according to the manufacturer's instructions and placed in the single-mode microwave cavity. Then the microwave was run at two stages: (a) 30 min at 60°C, 10 bar, and 300 W, and (b) 35 min, 70°C, 10 bar, and 300 W. After allowing the vial to cool to rt, saturated NaHCO_3 (3 mL) was added and stirred for 5 min. The reaction mixture was then filtered through celite and the filtrate was dried on a rotor evaporator to afford a yellow slurry. The slurry was then extracted with a mixture of hot ethyl acetate/ethanol mixture (2:1). The extract was then filtered through celite and dried over a rotor evaporator. The extraction process was repeated again. The filtrate was concentrated under vacuum and purified by gradient

column chromatography using silica gel as the stationary phase and mixture of methanol and dichloromethane (1:15 to 1:5) as the mobile phase.

NMR Data

1. Methyl α -D-Galactofuranoside (**3 α**)^[12,16a,28,29] (Main Data)

¹H NMR (CD₃OD, 500 MHz): δ 4.73 (1 H, J = 1.5 Hz, H-1).

¹³C NMR (CD₃OD, 125 MHz): δ 103.8 (C-1).

2. Methyl β -D-Galactofuranoside (**3 β**)^[12,16a,28,29] (Main Data)

¹H NMR (CD₃OD, 500 MHz): δ 4.71 (1 H, J = 4.6 Hz, H-1).

¹³C NMR (CD₃OD, 125 MHz): δ 110.8 (C-1).

3. Methyl α -D-Galactopyranoside (**4 α**)

¹H NMR (CD₃OD, 500 MHz): δ 4.62 (d, 1 H, J = 3.5 Hz, H-1), 3.79 (d, 1 H, J = 2.5 Hz), 3.57–3.72 (m, 5 H), 3.31 (s, 3 H, OMe).

¹³C NMR (CD₃OD, 125 MHz): δ 101.5 (C-1), 72.3, 71.5, 71.0, 70.2, 62.8 (C-6), 55.6 (OMe).

HRMS (EIMS, M⁺): calcd for C₇H₁₄O₆ 194.0790, found 194.0798.

4. Methyl β -D-Galactopyranoside (**4 β**)

¹H NMR (CD₃OD, 500 MHz): δ 4.02 (d, 1 H, J = Hz), 3.73 (d, 1 H, J = Hz), 3.64 (ddd, 2 H, J = Hz), 3.43 (s, 3 H, OMe), 3.33–3.42 (m, 3 H).

¹³C NMR (CD₃OD, 125 MHz): δ 106.0 (C-1), 76.7, 74.9, 72.5, 70.3, 62.5 (C-6), 57.2 (OMe).

HRMS (EIMS, M⁺): calcd for C₇H₁₄O₆ 194.0790, found 194.0788.

5. Allyl α -D-galactofuranoside (**15 α**)^[17]

¹H NMR (400 MHz, CD₃OD): δ 5.86 (dddd, 1 H, J = 17.4, 10.4, 6.2, 5.2 Hz, H^b), 5.19–5.28 (m, 1 H, H^a), 5.04–5.12 (m, 1 H, H^a), 4.81 (d, 1 H, J = 4.9 Hz, H-1), 4.15–4.25 (m, 1 H, H-3), 3.95–4.05 (m, 2 H, H-4 and H^c overlap), 3.84–3.90 (m, 1 H, H-2), 3.60–3.66 (m, 1 H, H^c), 3.50–3.58 (m, 2 H, H-6 and H-5 overlap), 3.41–3.50 (m, 1 H, H-6).

¹³C NMR (100 MHz, CD₃OD): δ 135.7, 117.5, 101.8, 83.5, 78.8, 76.2, 74.5, 69.7, 64.0.

HRMS (EIMS, M⁺ + Na): calcd for C₉H₁₆NaO₆ 243.0845, observed 243.0847.

6. *Allyl β-D-galactofuranoside (15β)*^[17]

¹H NMR (400 MHz, CD₃OD): δ 5.88 (dddd, 1 H, *J* = 17.4, 10.4, 6.2, 5.2 Hz, H^b), 5.28 (qd, 1 H, *J* = 17.0, 1.6 Hz, H^a), 5.11–5.14 (qd, 1 H, *J* = 10.5, 1.6 Hz, H^a), 4.89 (d, 1 H, *J* = 2.0 Hz, H-1), 4.16–4.23 (m, 1 H, H-3), 3.98–4.03 (m, 1 H, H-6), 3.95–3.98 (m, 3 H, H-2 and H^c overlap), 3.93 (dd, 1 H, *J* = 6.5, 3.2 Hz, H-6), 3.68–3.74 (m, 1 H, H-5), 3.62 (br s, -OH), 3.60 (d, 1 H, *J* = 1.3 Hz, H-4).

¹³C NMR (100 MHz, CD₃OD): δ 135.9 (C^b), 117.1 (C^a), 108.7 (H-1), 84.3 (C-6), 83.4 (C-2), 78.8 (C^c), 72.4 (C-5), 69.4 (C-3), 64.5 (C-4).

HRMS (EIMS, M⁺ + Na): calcd for C₉H₁₆NaO₆ 243.0845, observed 243.0848.

Crystallographic Data for 15β

CCDC 795662 <http://www.ccdc.cam.ac.uk/>

Formula: C₉ H₁₆ O₆

Unit cell parameters: a 9.2341(2) b 9.4269(2) c 12.2070(3) space group P212121

7. *Phenylethyl α-D-galactofuranoside (16α)*

¹H NMR (500 MHz, CDCl₃): δ 7.23–7.29 (m, 2 H, Ph), 7.15–7.21 (m, 3 H, Ph), 4.90 (d, 1 H, *J* = 3.4 Hz, H-1), 3.66–4.6 (m, 7 H, H-2, H-3, H-4, H-5, H-6 and -OCH₂CH₂Ph overlap), 3.40–3.46 (m, 1 H, -OCH₂CH₂Ph), 2.88 (t, 2 H, *J* = 6.7 Hz, -OCH₂CH₂Ph).

¹³C NMR (125 MHz, acetone-D₆): δ 140.1, 129.8, 129.0, 126.9, 99.9, 85.6, 83.8, 78.9, 76.1, 71.4, 62.5, 36.7.

¹³C NMR (125 MHz, CD₃OD): δ 140.3, 130.1, 129.3, 127.2, 100.2, 84.2, 83.3, 78.7, 76.6, 72.2, 71.5, 62.8, 36.9.

HRMS (EIMS, M⁺ + Na): calcd for C₁₄H₂₀NaO₆ 307.1158, observed 307.1163.

ACKNOWLEDGMENTS

We would like to thank Dr. Bashar Ksebati (Wayne State University) for NMR assistance, Dr. Lew Hryhorczuk (Wayne State University) for obtaining mass spectral data, and Dr. Mary Jane Heeg (Wayne State University) for x-ray data. P.R.A. also acknowledges WSU for start-up funds, and the NIH/NCI for an R03 CA153936-01.

REFERENCES

1. Peltier, P.; Euzen, R.; Daniellou, R.; Nugier-Chauvin, C.; Ferrieres, V. Recent knowledge and innovations related to hexofuranosides: structure, synthesis and applications. *Carbohydr. Res.* **2008**, *343*, 1897–1923.

2. Cobb, B.A.; Wang, Q.; Tzianabos, A.O.; Kasper, D.L. Polysaccharide processing and presentation by the MHC-II pathway. *Cell* **2004**, *117*, 677–687.
3. Van Bruggen-Van Der Lugt, A.W.; Kamphuis, H.J.; De Ruiter, G.A.; Mischnick, P.; Van Boom, J.H.; Rombouts, F.M. New structural features of the antigenic extracellular polysaccharides of *Penicillium* and *Aspergillus* species revealed with exo-beta-D-galactofuranosidase. *J. Bacteriol.* **1992**, *174*, 6096–6102.
4. Bennett, J.E.; Bhattacharjee, A.K.; Glaudemans, C.P.J. Galactofuranosyl groups are immunodominant in *Aspergillus fumigatus* galactomannan. *Mol. Immunol.* **1985**, *22*, 251–254.
5. Sarvas, M.; Nikaido, H. Biosynthesis of Tl antigen in *Salmonella*: origin of D-galactofuranose and D-ribofuranose residues. *J. Bacteriol.* **1971**, *105*, 1063–1072.
6. Marlow, A.L.; Kiessling, L.L. Improved chemical synthesis of UDP-galactofuranose. *Org. Lett.* **2001**, *3*, 2517–2519.
7. Pedersen, L.L.; Turco, S.J. Galactofuranose metabolism: a potential target for antimicrobial chemotherapy. *Cell. Mol. Life Sci.* **2003**, *60*, 259–266.
8. Fischer, E.H. Alkyl derivatives of benzoin. *Ber. Dtsch. Chem. Ges.* **1893**, *26*, 2400–2412.
9. Fischer, E.H.; Beensch, L. Synthetic glucosides. *Ber. Dtsch. Chem. Ges.* **1894**, *27*, 2478–2486.
10. Fischer, E.H. Sugar derivatives of alcohols and ketones. *Ber. Dtsch. Chem. Ges.* **1895**, *28*, 1145–1167.
11. Review: Capon, B. Mechanism in carbohydrate chemistry. *Chem. Rev.* **1969**, *69*, 407–498.
12. Lubineau, A.; Fischer, J.C. High-yielding one-step conversion of D-glucose and D-galactose to the corresponding methyl α - and β -glucofuranosides and galactofuranosides. *Synth. Commun.* **1991**, *21*, 815–818.
13. Euzen, R.; Ferrieres, V.; Plusquellec, D. Synthesis of galactofuranose-containing disaccharides using thioimidoyl-type donors. *Carbohydr. Res.* **2006**, *341*, 2759–2768.
14. Ferrieres, V.; Blanchard, S.; Fischer, D.; Plusquellec, D. A novel synthesis of D-galactofuranosyl, D-glucofuranosyl and D-mannofuranosyl 1-phosphates based on remote activation of new and free hexofuranosyl donors. *Bioorg. Med. Chem. Lett.* **2002**, *12*, 3515–3518.
15. Ferrieres, V.; Roussel, M.; Gelin, M.; Plusquellec, D. A new approach to a disaccharidic hapten containing a galactofuranosyl entity. *J. Carbohydr. Chem.* **2001**, *20*, 855–865.
16. (a) Ferrieres, V.; Bertho, J.-N.; Plusquellec, D. A new synthesis of O-glycosides from totally O-unprotected glycosyl donors. *Tetrahedron Lett.* **1995**, *36*, 2749–2752; (b) Bertho, J.-N.; Ferrieres, V.; Plusquellec, D. A new synthesis of D-glycoduronates from unprotected D-uronic acids. *J. Chem. Soc., Chem. Commun.* **1995**, *13*, 1391–1393.
17. Klotz, W.; Schmidt, R.R. Anomeric O-alkylation. 11. Anomeric O-alkylation of O-unprotected hexoses and pentoses – convenient synthesis of decyl, benzyl, and allyl glycosides. *Liebigs Ann. Chem.* **1993**, 683–690.
18. Baldoni, L.; Marino, C. Facile synthesis of per-O-tert-butyldimethylsilyl- β -D-galactofuranose and efficient glycosylation *via* the galactofuranosyl iodide. *J. Org. Chem.* **2009**, *74*, 1994–2003.

19. Kohn, P.; Samaritano, R.H.; Lerner, L.M. A new method for the synthesis of furanose derivatives of aldohexoses. *J. Am. Chem. Soc.* **1965**, *87*, 5469–5472.
20. Kovensky, J.; Sinay, P. Synthesis of galactofuranosides by regioselective ring opening of a 1,4-anhydrogalactopyranose derivative: A possible chemical model for an unprecedented enzymatic reaction. *Eur. J. Org. Chem.* **2000**, 3523–2525.
21. D'Accorso, N.B.; Thiel, I.M.E.; Schueller, M. Proton and carbon-13 nuclear magnetic resonance spectra of some benzoylated aldohexoses. *Carbohydr. Res.* **1983**, *124*, 177–184.
22. Green, J.W.; Pacsu, E. Glucofuranosides and thioglucofuranosides. III. New crystalline furanosides of D-galactose and l-arabinose. *J. Am. Chem. Soc.* **1938**, *60*, 2056–2057.
23. McAuliffe, J.C.; Hindsgaul, O. Use of acyclic glycosyl donors for furanoside synthesis. *J. Org. Chem.* **1997**, *62*, 1234–1239.
24. Szarek, W.A.; Zamojski, A.; Tiwari, K.N.; Ison, E.R. A new, facile method for cleavage of acetals and dithioacetals in carbohydrate derivatives. *Tetrahedron Lett.* **1986**, *27*, 3827–3830.
25. Completo, G.C.; Lowary, T.L. Synthesis of galactofuranose-containing acceptor substrates for Mycobacterial galactofuranosyltransferases. *J. Org. Chem.* **2008**, *73*, 4513–4525.
26. Richards, M.R.; Lowary, T.L. Chemistry and biology of galactofuranose-containing polysaccharides. *ChemBioChem.* **2009**, *10*, 1920–1938.
27. De Silva, R.A.; Wang, Q.; Chidley, T.; Appulage, D.K.; Andreana, P.R. Immunological response from an entirely carbohydrate antigen: design of synthetic vaccines based on Tn-PS A1 conjugates. *J. Am. Chem. Soc.* **2009**, *131*, 9622–9623.
28. Vojtech, M.; Petrusova, M.; Pribulova, B.; Petrus, L. Direct conversion of 1-deoxy-1-nitro-alditols to methyl glycofuranosides. *Tetrahedron Lett.* **2008**, *49*, 3112–3116.
29. Bordoni, A.; Lima, C.; Marino, K.; de Lederkremer, R.M.; Marino, C. Facile synthesis of methyl α - and β -D-[6-³H]galactofuranosides from D-galacturonic acid. Substrates for the detection of galactofuranosidases. *Carbohydr. Res.* **2008**, *343*, 10–11.
30. Angyal, S.J. Complexing the polyols with cations. *Tetrahedron*, **1974**, *30*, 1695–1702.
31. Evans, M.E.; Angyal, S.J. Complexes of carbohydrates with metal cations. II. Glycosidations of D-allose in the presence of strontium and calcium ions. *Carbohydr. Res.* **1972**, *25*, 43–48.
32. Arasappan, A.; Fraser-Reid, B. *n*-Pentenyl furanosides: synthesis and glycosidation reactions of some galacto derivatives. *Tetrahedron Lett.* **1995**, *36*, 7967–7970.
33. Angyal, S.J.; Bodkin, C.L.; Parrish, F.W. Complexes of carbohydrates with metal cations. V. Syntheses of methyl glycosides in the presence of metal ions. *Aust. J. Chem.* **1975**, *28*, 1541–1549.
34. Angyal, S.J.; Greeves, D.; Mills, J.A. Conformations of acyclic sugar derivatives. IV. Complexes of carbohydrates with metal cations. III. Conformations of alditols in aqueous solutions. *Aust. J. Chem.* **1974**, *27*, 1447–1456.
35. Cahn, R.S.; Ingold, C.; Prelog, V. Specification of molecular chirality. *Angew. Chem. Int. Ed.* **1966**, *5*, 385–415.
36. Clementi, E.; Raimondi, D.L.; Reinhardt, W.P. Atomic screening constants from SCF functions. II. Atoms with 37 to 86 electrons. *J. Chem. Phys.* **1967**, *47*, 1300–1307.

37. (a) Bordoni, A.; Lima, C.; Mariño, K.; Lederkremer, R.M.D.; Marino, C. Facile synthesis of methyl α - and β -D-[6- ^3H]galactofuranosides from D-galacturonic acid. Substrates for the detection of galactofuranosidases. *Carbohydr. Res.* **2008**, *343*, 1863–1869; (b) Euzen, R.; Guégan, J.-P.; Ferrières, V.; Plusquellec, D. First O-glycosylation from unprotected 1-thioimidoyl hexofuranosides assisted by divalent cations. *J. Org. Chem.* **2007**, *72*, 5743–5747; (c) Raunhardt, O.; Schmidt, H.W.H.; Hans, N. Gas-Chromatographische Untersuchungen An Uronsäuren Und Urons Äurederivaten. *Helv. Chim. Acta.* **1967**, *50*, 1267–1274.
38. Ellervik, U.; Magnusson, G. Anomeric effect in furanosides. Experimental evidence from conformationally restricted compounds. *J. Am. Chem. Soc.* **1994**, *116*, 2340–2347.
39. Kopf, J.; Koell, P. Influence of anomeric effect on the conformation of furanose. X-ray structural analysis of five isomeric methyl 3,6-anhydrohexofuranosides. *Carbohydr. Res.* **1984**, *135*, 29–46.
40. Hickenboth, C.R.; Moore, J.S.; White, S.R.; Sottos, N.R.; Baudry, J.; Wilson, S.R. Biasing reaction pathways with mechanical force. *Nature*, **2007**, *446*, 423–427.
41. Larsen, K.; Worm-Leonhard, K.; Olsen, P.; Hoel, A.; Jensen, K.J. Reconsidering glycosylations at high temperature: precise microwave heating. *Org. Biomol. Chem.* **2005**, *3*, 3966–3970.
42. Splain, R.A.; Kiessling, L.L. Synthesis of galactofuranose-based acceptor substrates for the study of the carbohydrate polymerase GlfT2. *Bioorg. Med. Chem. Lett.* **2010**, *18*, 3753–3759.