



Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/lcyc20>

Xanthan Sulfuric Acid: An Efficient, Biosupported, and Recyclable Solid Acid Catalyst for the Synthesis of 4,4'-(Arylmethylene)bis(1H-pyrazol-5-ols)

B. Suresh Kuarm^a & B. Rajitha^a

^a Department of Chemistry, National Institute of Technology, Warangal, India

Accepted author version posted online: 15 Dec 2011. Version of record first published: 14 May 2012.

To cite this article: B. Suresh Kuarm & B. Rajitha (2012): Xanthan Sulfuric Acid: An Efficient, Biosupported, and Recyclable Solid Acid Catalyst for the Synthesis of 4,4'-(Arylmethylene)bis(1H-pyrazol-5-ols), *Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry*, 42:16, 2382-2387

To link to this article: <http://dx.doi.org/10.1080/00397911.2011.557516>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.


The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

XANTHAN SULFURIC ACID: AN EFFICIENT, BIOSUPPORTED, AND RECYCLABLE SOLID ACID CATALYST FOR THE SYNTHESIS OF 4,4'-(ARYLMETHYLENE)BIS(1H-PYRAZOL-5-OLS)

B. Suresh Kuarm and B. Rajitha

Department of Chemistry, National Institute of Technology, Warangal, India

GRAPHICAL ABSTRACT



Abstract A simple and efficient method has been developed for the synthesis of 4,4'-(arylmethylene)bis(1H-pyrazol-5-ols) by the condensation reaction between substituted aldehydes, and 1-phenyl-3-methylpyrazol-5-one in the presence of xanthan sulfuric acid (XSA) as a solid acid catalyst. This method is simple and cost-effective with short reaction times. Yields are excellent with high purity, and the catalyst could be easily recycled.

Keywords Aldehydes; 4,4'-(arylmethylene)bis(1H-pyrazol-5-ols); biosupported; 1-phenyl-3-methylpyrazol-5-one; xanthan sulfuric acid

INTRODUCTION

Synthesis of nitrogen-containing heterocyclic systems occupies an important role in the realm of natural and synthetic organic chemistry because of their therapeutic and pharmacological properties.^[1] In particular, pyrazoles and their derivatives have attracted considerable attention because of their wide variety of biological activities, including anti-inflammatory,^[2] antipyretic,^[3] gastric secretion stimulatory,^[4] antidepressant,^[5] antibacterial,^[6] antifilarial,^[7] bactericidal, fungicidal,^[8] and promising inhibitory activities against monoamine oxidase for the treatment of diseases such as Parkinson's and Alzheimer's.^[9] The conventional chemical approach to 4,4'-(arylmethylene)bis(3-methyl-1-phenyl-pyrazol-5-ols) involves the successive Knoevenagel synthesis of the corresponding arylidenepyrazolones, its base-promoted Michael reaction, and one-pot tandem Knoevenagel–Michael

Received November 23, 2010.

Address correspondence to B. Rajitha, Department of Chemistry, National Institute of Technology, Warangal, India. E-mail: rajithabhargavi@gmail.com


reaction of arylaldehydes with 2 equiv. of 5-methyl-2-phenyl-2,4-dihydro-3H-pyrazol-3-one performed under a variety of reaction conditions.^[10,11] The first set of procedures utilizes the catalysis of the components with piperidine in ethanolic solution.^[12] The second set of methods involves the uncatalyzed tandem Knoevenagel–Michael reaction under neutral conditions in either ethanol^[13] or benzene^[14] solutions. Although it affords the corresponding 4,4'-(arylmethylene)bis(1H-pyrazol-5-ols) in reliable 70–90% yields, the reaction requires 3–12 h of initial reflux with a further 24 h under ambient temperature to go to completion. Wang et al.^[15] reported its synthesis in water using sodium dodecyl sulfate as the surfactant catalyst over a 1-h period, but the process needs a temperature of 100 °C. Finally, Elinson et al. utilized electrocatalytic procedure for its synthesis.^[16] Further, Perumal and coworkers reported the synthesis and antiviral activity of 4,4'-(arylmethylene)bis(3-methyl-1-phenyl-1H-pyrazol-5-ols) using ceric ammonium nitrate (CAN) as a catalyst.^[17] However, most of the methods suffer from at least one limitation that may include moderate yields, long reaction times, harsh reaction conditions, or tedious workup procedures. Therefore, the search continues for a better catalyst for the synthesis of 4,4'-(arylmethylene)bis(3-methyl-1-phenyl-pyrazol-5-ols) in terms of operational simplicity, reusability of catalyst, low cost, and greater selectivity.

Xanthan and its derivatives^[18–20] have some unique properties that make them attractive alternatives for conventional organic or inorganic supports for catalytic applications. Xanthan is the most abundant bacterial exopolysaccharide in the world, which is produced through fermentation, and it has been widely studied during recent decades, as it is a biodegradable material and a renewable resource. Unlike other gums, it is very stable under a wide range of temperatures and pH values. Xanthan sulfuric acid^[21] can be easily prepared by the reaction of xanthan with chlorosulfonic acid; the number of acidic (H⁺) sites in the xanthan sulfuric acid determined by acid–base titration was 0.6 meq/g.

RESULTS AND DISCUSSION

Our research group has developed various new synthetic methodologies for the synthesis of both building blocks and heterocyclic compounds using new reagents.^[22] Thus, in continuation of our interest in the development of new methodologies, we describe herein the synthesis of a 4,4'-(arylmethylene)bis(3-methyl-1-phenyl-pyrazol-5-ols) in ethanol catalyzed by xanthan sulfuric acid (Scheme 1).

We investigated the efficiency of xanthan sulfuric acid compared to various sulfur analog acidic catalysts. The results are summarized in Table 1. Xanthan sulfuric



Scheme 1. Synthesis of 4,4'-(arylmethylene)bis(1H-pyrazol-5-ols) with xanthan sulfuric acid.

Table 1. Effect of catalysts on yield

Entry	Catalyst	Yield (%) ^a
1	Xanthan sulfuric acid	95
2	Silica sulfuric acid	88
3	Sulfuric acid in acetic acid	52
4	No catalyst	15

^aIsolated yield.**Table 2.** Influence of the catalytic amounts of xanthan sulfuric acid

Entry	Catalyst (g)	Time (min)	Yield (%) ^a
1	None	60	Trace
2	0.01	15	25
3	0.03	15	55
4	0.05	15	79
5	0.08	30	95
6	0.08	15	95

^aIsolated yield.

acid was found to be the most effective catalyst based on the product yield. The reaction did not proceed in the absence of catalyst (yield less than 15%).

We examined the amount of catalyst required in this reaction. The best results were obtained using 0.08 g of catalyst (95%). Using lesser amounts of catalyst resulted in lesser yields, and in the absence of catalyst the yield of the product was found to be very poor Table 2.

EXPERIMENTAL

All the melting points are uncorrected. The progress of the reaction was monitored by thin-layer chromatography (TLC). Infrared (IR) spectra (KBr) were recorded on a Shimadzu FTIR model 8010 spectrometer, and the ¹H NMR spectra was determined on a Varian Gemini 200-MHz spectrometer using tetramethylsilane (TMS) as internal standard. Mass spectra were recorded on a Jeol JMS D-300 spectrometer. CHN analysis was done on a Carlo Erba EA 1108 automatic elemental analyzer. All solvents and reagents were purchased from Aldrich and Fluka.

Typical Procedure

A mixture of aldehyde (1 mmol), 5-methyl-2-phenyl-2,4-dihydro-3H-pyrazol-3-one (2 mmol), and xanthane sulfuric acid (0.08 g) in ethanol (5 mL) was heated under refluxing conditions for an appropriate time according to Table 3. After the completion of the reaction (as monitored by TLC), the reaction mixture was filtered and washed with ethanol (2 × 20 mL) to separate the catalyst. Then the filtrate was evaporated under reduced pressure to afford a pure product. Further purification

Table 3. Synthesis of 4,4'-(arylmethylene)bis(1H-pyrazol-5-ols) with xanthan sulfuric acid

Entry	Aldehydes	Product	Time (min)	Yield (%) ^a	Mp (°C)
1	Benzaldehyde	3a	15	95	166–167 ^[16]
2	3-Methylbenzaldehyde	3b	25	91	242–243 ^[17]
3	4-Methylbenzaldehyde	3c	25	90	203–204 ^[17]
4	3-Hydroxybenzaldehyde	3d	25	92	165–168 ^[23]
5	4-Hydroxybenzaldehyde	3e	30	91	152–153 ^[15]
6	4-Nitrobenzaldehyde	3f	15	90	230–232 ^[15]
7	2-Chlorobenzaldehyde	3g	25	90	236–237 ^[15]
8	4-Chlorobenzaldehyde	3h	20	94	207–209 ^[15]
9	4-Methoxybenzaldehyde	3i	25	92	148 ^[11]
10	2-Furfuraldehyde	3j	15	95	189–190 ^[17]
11	2-Naphthaldehyde	3k	25	85	185–186
12	4-Oxo-4 <i>H</i> -chromene-3-carbaldehyde	3l	30	78	202
13	6-Nitro-4-oxo-4 <i>H</i> -chromene-3-carbaldehyde	3m	30	76	190

^aYields refer to pure solid products; all products were characterized by spectral data.

Table 4. Effect of reusability of catalyst on yield^a

Run	Cycle	Yield (%) ^b
1	0	95
2	1	92
3	2	88
4	3	76

^aReaction conditions: mixture of benzaldehyde (1 mmol), 5-methyl-2-phenyl-2,4-dihydro-3*H*-pyrazol-3-one (2 mmol), and xanthane sulfuric acid (0.08 g) in ethanol (5 mL) was heated under reflux.

^bYields refer to the pure, isolated, recovered catalyst.

was followed by crystallization from ethanol. The recovered catalyst was reused for subsequent runs (Table 4).

Analytical Data

Table 3, Entry 11. IR (KBr): 3421, 2949, 1610 cm⁻¹; ¹H NMR (DMSO-*d*₆): δ = 2.33 (s, 6 H, 2 CH₃), 5.23 (s, 1H, CH), 6.95–7.78 (m, 17 H, Ar); MS (*m/z*): 487 (M⁺). Anal. Calcd. for C₃₁H₂₆N₄O₂: C, 76.52; H, 5.39; N, 11.51; Found: C, 76.25; H, 5.24; N, 12.10.

Table 3, Entry 12. IR (KBr): 3439, 2945, 1604 cm⁻¹; ¹H NMR (DMSO-*d*₆): δ = 2.46 (s, 6 H, 2 CH₃), 5.21 (s, 1H, CH), 6.19 (s, 1H, CH) 7.21–7.86 (m, 14 H, Ar); MS (*m/z*): 504 (M⁺). Anal. calcd. for C₃₀H₂₄N₄O₄: C, 71.42; H, 4.79; N, 11.10; found C, 71.21; H, 4.70; N, 11.18.

Table 3, Entry 13. IR (KBr): 3411, 2951, 1609 cm⁻¹; ¹H NMR (DMSO-*d*₆): δ = 2.34 (s, 6 H, 2 CH₃), 5.24 (s, 1H, CH), 6.21 (s, 1H, CH) 7.18–7.88 (m, 13 H,

Ar); MS (m/z): 549 (M^+). Anal. calcd. for $C_{30}H_{23}N_5O_6$: C, 65.57; H, 4.22; N, 12.74. found: C, 65.46; H, 4.34; N, 12.61.

CONCLUSION

We have developed a simple and efficient method for the synthesis of 4,4'-(arylmethylene)bis(1H-pyrazol-5-ols) using xanthan sulfuric acid. The short reaction times, simple workup, good yields, mild reaction conditions, and recyclability of the catalyst are features of this new procedure.

ACKNOWLEDGMENT

B. S. K. is grateful to the Council of Scientific and Industrial Research, New Delhi, India, for providing financial support in the form of a senior research fellowship.

REFERENCES

1. Elguero, J. Pyrazoles. In *Comprehensive Heterocyclic Chemistry II*; A. R. Katritzky, C. W. Rees, and E. F. V. Scriven, (Eds.); Pergamon: Oxford, UK, 1996; vol. 3, pp. 1–76.
2. Sugiura, S.; Ohno, S.; Ohtani, O.; Izumi, K.; Kitamikado, T.; Asai, H.; Kato, K. Syntheses and anti-inflammatory and hypnotic activity of 5-alkoxy-3-(N-substituted carbamoyl)-1-phenylpyrazoles. *J. Med. Chem.* **1977**, *20*, 80–85.
3. Behr, L. C.; Fusco, R.; Jarboe, C. H. In *The Chemistry of Heterocyclic Compounds, Pyrazoles, Pyrazolines, Pyrazolidines, Indazoles, and Condensed Rings*; A. Weissberger, (Eds.); Interscience: New York, 1967.
4. Rosiere, C. E.; Grossman, M. I. An analog of histamine that stimulates gastric acid secretion without other actions of histamine. *Science* **1951**, *113*, 651.
5. Bailey, D. M.; Hansen, P. E.; Hlavac, A. G.; Baizman, E. R.; Pearl, J.; Defelice, A. F.; Feigenson, M. E. 3,4-Diphenyl-1H-pyrazole-1-propanamine antidepressants. *J. Med. Chem.* **1985**, *28*, 256–260.
6. Mahajan, R. N.; Havaladar, F. H.; Fernandes, P. S. Syntheses and biological activity of heterocycles derived from 3-methoxy-1-phenyl-1H-pyrazole-5-carboxylate. *J. Indian Chem. Soc.* **1991**, *68*, 245–246.
7. Chauhan, P. M. S.; Singh, S.; Chatterjee, R. K. Antimalarial profile of substituted pyrazoles: A new class of antimalarial agents. *Indian J. Chem., Sect. B* **1993**, *32*, 858–861.
8. Djuric, S. W.; Bmaung, N. Y.; Basha, A.; Liu, H.; Luly, J. R.; Madar, D. J.; Sciotti, R. J.; Tu, N. P.; Wagenaar, F. L.; Wiedman, P. E.; Zhou, X.; Ballaron, S.; Bauch, J.; Chen, Y.-W.; Chiou, X. G.; Fey, T.; Gauvin, D.; Gubbins, E.; Hsieh, G. C.; Marsch, K. C.; Mollison, K. W.; Pong, M.; Shaughnessy, T. K.; Sheets, M. P.; Smith, M.; Trevillyan, J. M.; Warrior, U.; Wegner, C. D.; Carter, G. W. 3,5-Bis(trifluoromethyl)pyrazoles: A novel class of NFAT transcription factor regulator. *J. Med. Chem.* **2000**, *43*, 2975.
9. Chimenti, F.; Maccioni, E.; Secci, D.; Bolasco, A.; Chimenti, P.; Granese, A.; Befani, O.; Turini, P.; Alcaro, S.; Ortuso, F.; Cirilli, R.; La Torre, F.; Cardia, M. C.; Simona, D. Synthesis, molecular modeling studies, and selective inhibitory activity against monoamine oxidase of 1-thiocarbamoyl-3,5-diaryl-4,5-dihydro-(1H)-pyrazole derivatives. *J. Med. Chem.* **2005**, *48*, 7113.
10. Hamama, W. S. Pyrazolones as versatile precursors for the synthesis of fused and binary heterocycles. *Synth. Commun.* **2001**, *31*, 1335–1345.

11. Li, X.-L.; Wang, Y.-M.; Tian, B.; Matsuura, T.; Meng, J.-B. J. The solid-state Michael addition of 3-methyl-1-phenyl-5-pyrazolone. *J. Heterocycl. Chem.* **1998**, *35*, 129–134.
12. Singh, D.; Singh, D. Syntheses of 1,3-disubstituted 4-arylidene-pyrazolin-5-ones and the keto and enol forms of 4,40-arylidenebis(1,3-disubstituted pyrazolin-5-ones). *J. Chem. Eng. Data* **1984**, *29*, 355–356.
13. Pavlov, P. T.; Goleneva, A. F.; Lesnov, A. E.; Prokhorova, T. S. Biological activity of some pyrazolone derivatives. *Pharm. Chem. J. (Engl. Trans.)* **1998**, *32*, 370–372.
14. Buzykin, B. I.; Lonshchakova, T. I. Reaction of azines with 5-pyrazolones. *Bull. Acad. Sci. USSR* **1971**, 2224–2226.
15. Wang, W.; Wang, S.-X.; Qin, X.-Y.; Li, J.-T. Reaction of aldehydes and pyrazolones in the presence of sodium dodecyl sulfate in a aqueous media. *Synth. Commun.* **2005**, *35*, 1263–1269.
16. Elinson, M. N.; Dorofeev, A. S.; Nasybullin, R. F.; Nikishin, G. I. Facile and convenient synthesis of 4,4'-(arylmethylene)bis(1H-pyrazol-5-ols) by electrocatalytic tandem Knoevenagel–Michael reaction. *Synthesis* **2008**, *12*, 1933–1937.
17. Sujatha, K.; Shanthi, G.; Selvam, N. P.; Manoharan, S.; Perumal, P. T.; Rajendran, M. Synthesis and antiviral activity of 4,4'-(arylmethylene)bis(1H-pyrazol-5-ols) against peste des petits ruminant virus (PPRV). *Bioorg. Med. Chem. Lett.* **2009**, *19*, 4501–4503.
18. Sargent, E. V.; Adolph, J.; Clemmons, M. K.; Kirk, G. D.; Pena, B. M.; Fedoruk, M. J. Evaluation of flu-like symptoms in workers handling xanthan gum powder, *J. Occup. Med.* **1990**, *32*, 625.
19. Davidson, R. L. (Ed.). *Handbook of Water-Soluble Gums and Resins*; McGraw-Hill: 1980.
20. Whistler, R. L.; BeMiller, J. N. (Eds.). *Industrial Gums: Polysaccharides and Their Derivatives*; Academic Press: 1973.
21. Shaabani, A.; Maleki, A.; Soudi, M. R.; Mofakham, H. Xanthan sulfuric acid: A new and efficient biosupported solid acid catalyst for the synthesis of α -amino nitriles by condensation of carbonyl compounds, amines, and trimethylsilylcyanide. *Catal. Commun.* **2009**, *10*, 945.
22. (a) Venu Madhav, J.; Thirupathi Reddy, Y.; Narsimha Reddy, P.; Nikhil Reddy, M.; Suresh, K.; Crooks, P. A.; Rajitha, B. Cellulose sulfuric acid: An efficient biodegradable and recyclable solid acid catalyst for the one-pot synthesis of aryl-14H-dibenzo[a,j]xanthenes under solvent-free conditions. *J. Mol. Catal. A* **2009**, *304*, 85–87; (b) Venu Madhav, J.; Suresh Kuarm, B.; Rajitha, B. Dipyrindine cobalt chloride: A novel and efficient catalyst for the synthesis of aryl-4H-dibenzo[a,j]xanthenes under solvent-free conditions. *Arkivoc*, **2008**, *12*, 204; (c) Venu Madhav, J.; Suresh Kuarm, B.; Someshwar, P.; Rajitha, B. Thirupathi Reddy, Y.; Crooks, P. A. CuPy₂Cl₂: A novel and efficient catalyst for synthesis of propargylamines under conventional method and microwave irradiation. *Synth. Commun.* **2008**, *38*, 3215; (d) Venu Madhav, J.; Suresh Kuarm, B.; Rajitha, B. The solid-state synthesis of 1,3-selenazoles by employing the CuPy₂Cl₂ as a Lewis acid catalyst. *Synth. Commun.* **2008**, *38*, 3514–3522; (e) Venu Madhav, J.; Suresh Kuarm, B.; Rajitha, B. Dipyrindine copper chloride as a mild and efficient catalyst for the solid-state synthesis of 2-substituted benzimidazoles. *Arkivoc* **2008**, *13*, 145–150 (f) Venu Madhav, J.; Suresh Kuarm, B.; Someshwar, P.; Rajitha, B. Thirupathi Reddy, Y.; Crooks, P. A. Dipyrindine cobalt chloride: A novel catalyst for the synthesis of coumarins via Pechmann condensation. *J. Chem. Res.* **2008**, 232–234; (g) Suresh Kuarm, B.; Venu Madhav, J.; Vijaya Laxmi, S.; Rajitha, B.; Thirupathi Reddy, Y.; Narsimha Reddy, P.; Crooks, P. A. An expeditious Pechmann condensation by using biodegradable cellulose sulfuric acid as a solid acid catalyst. *Synth. Commun.* **2010**, *40*, 3358; (h) Suresh Kuarm, B.; Thirupathi Reddy, Y.; Venu Madhav, J.; Crooks, P. A.; Rajitha, B. 3-[Benzimidazo- and 3-[benzothiadiazoleimidazo-(1,2-c)quinazolin-5-yl]-2H-chromene-2-ones as potent antimicrobial agents. *Bioorg. Med. Chem. Lett.* **2011**, *21*, 524–527.
23. Niknam, K.; Saberi, D.; Sadegheyan, M.; Deris, A. Silica-bonded S-sulfonic acid: An efficient and recyclable solid acid catalyst for the synthesis of 4,4'-(arylmethylene)bis(1H-pyrazol-5-ols). *Tetrahedron Lett.* **2010**, *51*, 692–694.