

## Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/lncn20>

### STRUCTURE-ACTIVITY RELATIONSHIPS OF APIO NUCLEOSIDES AS POTENTIAL ANTIVIRAL AGENTS

Lak Shin Jeong<sup>a</sup>, Hyung Ryong Moon<sup>a</sup>, Jun Hee Hong<sup>a</sup>, Su Jeong Yoo<sup>a</sup>, Won Jun Choi<sup>a</sup>, Hea Ok Kim<sup>b</sup>, Hee Sung Ahn<sup>a</sup>, Hye Woo Baek<sup>a</sup>, Moon Woo Chun<sup>c</sup>, Hee-Doo Kim<sup>d</sup>, Jeongmin Kim<sup>e</sup> & Jong-Ryoo Choi<sup>e</sup>

<sup>a</sup> College of Pharmacy, Ewha Womans University, Seoul, Korea

<sup>b</sup> College of Medicine, Yonsei University, Seoul, Korea

<sup>c</sup> College of Pharmacy, Seoul National University, Seoul, Korea

<sup>d</sup> College of Pharmacy, Sookmyung Women's University, Seoul, Korea

<sup>e</sup> LG Chem/Biotech Research Institute, Taejon, Korea

Published online: 07 Feb 2007.

To cite this article: Lak Shin Jeong, Hyung Ryong Moon, Jun Hee Hong, Su Jeong Yoo, Won Jun Choi, Hea Ok Kim, Hee Sung Ahn, Hye Woo Baek, Moon Woo Chun, Hee-Doo Kim, Jeongmin Kim & Jong-Ryoo Choi (2001) STRUCTURE-ACTIVITY RELATIONSHIPS OF APIO NUCLEOSIDES AS POTENTIAL ANTIVIRAL AGENTS, *Nucleosides, Nucleotides and Nucleic Acids*, 20:4-7, 657-660, DOI: [10.1081/NCN-100002344](https://doi.org/10.1081/NCN-100002344)

To link to this article: <http://dx.doi.org/10.1081/NCN-100002344>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

NUCLEOSIDES, NUCLEOTIDES & NUCLEIC ACIDS, 20(4–7), 657–660 (2001)

## STRUCTURE-ACTIVITY RELATIONSHIPS OF APIO NUCLEOSIDES AS POTENTIAL ANTIVIRAL AGENTS

Lak Shin Jeong,<sup>1,\*</sup> Hyung Ryong Moon,<sup>1</sup> Jun Hee Hong,<sup>1</sup>  
Su Jeong Yoo,<sup>1</sup> Won Jun Choi,<sup>1</sup> Hea Ok Kim,<sup>2</sup> Hee Sung Ahn,<sup>1</sup>  
Hye Won Baek,<sup>1</sup> Moon Woo Chun,<sup>3</sup> Hee-Doo Kim,<sup>4</sup>  
Jeongmin Kim,<sup>5</sup> and Jong-Ryoo Choi<sup>5</sup>

<sup>1</sup>College of Pharmacy, Ewha Womans University, Seoul, Korea

<sup>2</sup>College of Medicine, Yonsei University, Seoul, Korea

<sup>3</sup>College of Pharmacy, Seoul National University, Seoul, Korea

<sup>4</sup>College of Pharmacy, Sookmyung Women's University, Seoul, Korea

<sup>5</sup>LG Chem/Biotech Research Institute, Taejeon, Korea

### ABSTRACT

Several types of novel apio nucleosides were synthesized starting from 1,3-dihydroxyacetone and evaluated for antiviral activity. Among compounds tested, amino substituted apio dideoxynucleosides exhibited anti-HBV activity, while thioapio dideoxynucleosides were found to be active against HIV-1. Apio dideoxydidehydro nucleosides showed moderate to potent anti-HCMV activity, but their bioisosteric thioapio dideoxydidehydro nucleosides did not exhibit any significant antiviral activity.

### INTRODUCTION

One of the nonclassical nucleosides, apio dideoxynucleoside (1–5) in which 4'-hydroxymethyl group of the classical 2',3'-dideoxynucleosides moves to the 3'-position has been shown anti-HIV activity (1) in MT-4 cells. In addition to antiviral activity, these nucleosides also showed metabolic advantages such as

---

\*Corresponding author.




Figure 1.

resistance to adenosine deaminase and glycosyl bond hydrolysis (3). In order to study structure-activity relationships of apio nucleosides, we synthesized several types of apio nucleosides such as azido or amino substituted apio dideoxynucleosides (1), apio dideoxydidehydro nucleosides (2), thioapio dideoxynucleosides (3), and thioapio dideoxydidehydro nucleosides (4) and evaluated them for antiviral activities (Fig. 1).


Here, we report the synthesis and structure-activity relationship study of novel apio nucleosides as antiviral agents.

## RESULTS AND DISCUSSION

The amino substituted apio nucleosides (1) and apio dideoxydidehydro nucleosides (2a–2f) were synthesized using 1,3-dihydroxyacetone as a common starting material via lactone 5 as the common intermediate (Scheme 1). These two classes of nucleosides were synthesized according to the known procedure reported by our laboratory (6,7).


Thioapio dideoxynucleosides which may act as bioisosteres of apio dideoxynucleosides were synthesized starting from the lactone 5 (Scheme 2).

The lactone 5 was protected as *tert*-butyldiphenylsilyl ether to give 7 which was hydrogenated under palladium on carbon to afford saturated lactone (±)-8 in quantitative yield. Treatment of (±)-8 with 1 N NaOH followed by methylation of the resulting acid gave the hydroxyester (±)-9 which was converted to the thioacetate (±)-10 in two steps. Cyclization of (±)-10 was achieved using DIBAL at  $-78^{\circ}\text{C}$  to give lactol, which was acetylated to give the thioapio acetate (±)-11.



Scheme 1.






Scheme 2.

The glycosyl donor (±)-**11** was condensed with silylated uracil, thymine, and 6-chloropurine in the presence of TMSOTf to give the inseparable anomeric mixtures of protected nucleosides to give the inseparable anomeric mixtures of protected nucleosides (±)-**12**, respectively. The protected nucleosides (±)-**12** was treated with tetra-*n*-butylammonium fluoride to give β anomers ((±)-**13a**, **14a**, and **15a**) and α-anomers ((±)-**13b**, **14b**, and **15b**), respectively. The 6-chloropurine derivatives, (±)-**15a** and (±)-**15b** were converted to the adenine ((±)-**16a** and (±)-**16b**), N-methyladenine ((±)-**17a** and (±)-**17b**), and hypoxanthine ((±)-**18a** and (±)-**18b**) derivatives, respectively according to the conventional methods.

Synthesis of the thioapio dideoxydidehydro nucleosides is shown in Scheme 3. Treatment of (±)-**10** with 1 N NaOH gave the mercapto acid which without purification, was cyclized using DCC to the thiolactone (±)-**19**. Treatment of (±)-**19**



Scheme 3.



with LiHMDS at  $-78^{\circ}\text{C}$  followed by the trapping of the enolate with TMSCl gave silylenol ether, which was reacted with phenyl selenenyl bromide at  $-78^{\circ}\text{C}$  to afford ( $\pm$ )-**20** (cis/trans = 10/1 as determined by  $^1\text{H}$  NMR). Reduction of ( $\pm$ )-**20** with DIBAL followed by acetylation gave the key intermediate ( $\pm$ )-**21**. Condensation of ( $\pm$ )-**21** with silylated uracil and thymine gave the inseparable mixture of protected nucleosides ( $\pm$ )-**22** and ( $\pm$ )-**23**, respectively. Syn-elimination using *m*CPBA and catalytic amounts of pyridine followed by desilylation yielded the final nucleosides ( $\pm$ )-**24** and ( $\pm$ )-**25**, respectively.

All synthesized compounds were assayed against several viruses such as against HIV-1, HSV-1, HSV-2, HCMV, and HBV. Among compounds tested, amino substituted apio dideoxynucleoside (**1**) exhibited anti-HBV activity, while their bioisosteric thioapio dideoxynucleosides were found to be only active against HIV-1. Most apio dideoxydidehydro nucleosides showed moderate to potent anti-HCMV activity without cytotoxicity, among which 5-fluorouracil derivative **2c** was found to be the most potent. However, thioapio analogues, ( $\pm$ )-**24** and ( $\pm$ )-**25** did not exhibit any significant antiviral activities.

### ACKNOWLEDGMENTS

This research was supported by the grant of the Good Health R & D Project, Ministry of Health and Welfare, Korea (HMP-98-D-4-0057).

### REFERENCES

1. Terao, Y.; Akamatsu, M.; Achiwa, K. *Chem. Pharm. Bull.* **1991**, *39*, 823–825.
2. Bamford, M. J.; Humber, D. C.; Storer, R. *Tetrahedron Lett.* **1991**, *32*, 271–274.
3. Nair, V.; Nuesca, Z. *J. Am. Chem. Soc.* **1992**, *114*, 7951–7952.
4. Sells, T. B.; Nair, V. *Tetrahedron Lett.* **1992**, *33*, 7639–7642.
5. Sells, T. B.; Nair, V. *Tetrahedron Lett.* **1993**, *34*, 3527–3530.
6. Jeong, L. S.; Lee, Y. A.; Moon, H. R.; Chun, M. W. *Nucleosides Nucleotides* **1998**, *17*, 1473–1487.
7. Jeong, L. S.; Lee, Y. A.; Moon, H. R.; Yoo, S. J.; Kim, S. Y. *Tetrahedron Lett.* **1998**, *39*, 7517–7520.



## **Request Permission or Order Reprints Instantly!**

Interested in copying and sharing this article? In most cases, U.S. Copyright Law requires that you get permission from the article's rightsholder before using copyrighted content.

All information and materials found in this article, including but not limited to text, trademarks, patents, logos, graphics and images (the "Materials"), are the copyrighted works and other forms of intellectual property of Marcel Dekker, Inc., or its licensors. All rights not expressly granted are reserved.

Get permission to lawfully reproduce and distribute the Materials or order reprints quickly and painlessly. Simply click on the "Request Permission/Reprints Here" link below and follow the instructions. Visit the [U.S. Copyright Office](#) for information on Fair Use limitations of U.S. copyright law. Please refer to The Association of American Publishers' (AAP) website for guidelines on [Fair Use in the Classroom](#).

The Materials are for your personal use only and cannot be reformatted, reposted, resold or distributed by electronic means or otherwise without permission from Marcel Dekker, Inc. Marcel Dekker, Inc. grants you the limited right to display the Materials only on your personal computer or personal wireless device, and to copy and download single copies of such Materials provided that any copyright, trademark or other notice appearing on such Materials is also retained by, displayed, copied or downloaded as part of the Materials and is not removed or obscured, and provided you do not edit, modify, alter or enhance the Materials. Please refer to our [Website User Agreement](#) for more details.

**[Order now!](#)**

Reprints of this article can also be ordered at

<http://www.dekker.com/servlet/product/DOI/101081NCN100002344>