## Palladium-Catalyzed Intramolecular Nucleophilic Substitution at the Alkoxycarbonyl Group\*\*

Daniel Solé\* and Olga Serrano

Transition-Metal Catalysis

The widespread application of palladium catalysts in organic synthesis owes mainly to their tolerance of most important functional groups, largely as a result of the low nucleophilicity of organopalladium compounds.<sup>[1]</sup> Thus, aryl and vinyl palladium complexes are commonly used as electrophiles in C-C coupling reactions, and there are only a few examples of their use as nucleophiles in reactions with polar electrophilic multiple bonds  $^{[2-5]}$  Alkoxycarbonyl groups have long been considered inert toward organopalladium reagents. In many cases, substrates that bear an ester moiety can undergo palladium-mediated processes in which the alkoxycarbonyl group is not modified in any way.<sup>[6]</sup> For example, the palladium-catalyzed  $\alpha$  arylation of esters,<sup>[7]</sup> which involves the nucleophilic attack of an enolate at a  $\sigma$ -aryl palladium complex, has emerged recently as a straightforward method for the preparation of  $\alpha$ -aryl esters.<sup>[8,9]</sup>

We described recently the Pd<sup>0</sup>-catalyzed intramolecular nucleophilic addition of aryl halides to amino-tethered ketones [Eq. (1); Bn = benzyl].<sup>[3c]</sup> This process appears to

$$(1)$$

involve the formation of a four-membered azapalladacyclic intermediate, which strongly modifies the interaction of the metal center with the carbonyl group and forces the otherwise unexpected nucleophilic addition. Herein we report that we can take advantage of the formation of such transient intermediates to force nucleophilic attack at functional groups less electrophilic than the ketone carbonyl group, such as an alkoxycarbonyl group. In particular, we describe the Pd<sup>0</sup>-catalyzed intramolecular coupling of  $\beta$ -(2-iodoanilino)esters to give dihydroquinolin-4-ones [Eq. (2)].<sup>[10]</sup>

[*]	Dr. D. Solé, O. Serrano
	Laboratori de Química Orgànica
	Facultat de Farmàcia
	Universitat de Barcelona
	Av. Joan XXIII s/n, 08028-Barcelona (Spain)
	Fax: (+34) 934-024-539
	E-mail: dsole@ub.edu

- [\*\*] This research was supported by the MEC (Spain) and FEDER (CTQ2004-04701/BQU) and by the DURSI (Catalonia) (grant 2005SGR-00442). D.S. thanks the DURSI for "Distinció per a la Promoció de la Recerca Universitària".
- Supporting information for this article is available on the WWW under http://www.angewandte.org or from the author.



Our initial efforts were focused on the Pd<sup>0</sup>-catalyzed intramolecular coupling reaction of ester **1a** (Table 1). The use of  $[Pd(PPh_3)_4]$  as the catalyst and  $Cs_2CO_3$  as the base, a

**Table 1:** Optimization of the Pd<sup>0</sup>-catalyzed intramolecular nucleophilic substitution of **1a**.

Me	X N Me 1a, X = I 1b, X = Br	OMe Me		Me		O OMe Me 3
Entry	Catalyst <sup>[a]</sup>	Additives ([equiv])	Solvent	Т [°С] <sup>[b]</sup>	t [h]	Products (yield [%]) <sup>[c]</sup>
1	[Pd(PPh <sub>3</sub> ) <sub>4</sub> ]	Cs <sub>2</sub> CO <sub>3</sub> (2)	THF	110	24	1a (40), 3 (40)

1	[Pd(PPh <sub>3</sub> ) <sub>4</sub> ]	$Cs_2CO_3$ (2)	THF	110	24	1 a (40), 3 (40)
2	[Pd(PPh₃)₄]	K <sub>3</sub> PO <sub>4</sub> (3)	THF	110	31	<b>2</b> (30), <b>3</b> (25) <sup>[d]</sup>
3	[Pd(PPh <sub>3</sub> ) <sub>4</sub> ]	K <sub>3</sub> PO <sub>4</sub> (3)	toluene	110	72	<b>2</b> (65), <b>3</b> (30)
		Et₃N (10)				
4	[Pd(PPh <sub>3</sub> ) <sub>4</sub> ]	K <sub>3</sub> PO <sub>4</sub> (3)	toluene	130	48	<b>2</b> (45), <b>3</b> (45)
		Et <sub>3</sub> N (10)				
5	[Pd(PPh <sub>3</sub> ) <sub>4</sub> ]	K <sub>3</sub> PO <sub>4</sub> (3)	toluene	90	100	<b>1</b> a (79), <b>2</b> (10)
		Et <sub>3</sub> N (10)				<b>3</b> (5)
6	[Pd(PPh <sub>3</sub> ) <sub>4</sub> ]	K <sub>3</sub> PO <sub>4</sub> (2)	toluene	110	72	<b>3</b> (36) <sup>[e]</sup>
		Ag <sub>3</sub> PO <sub>4</sub> (1)				
		Et <sub>3</sub> N (10)				
7	[Pd₂(dba)₃]	K <sub>3</sub> PO <sub>4</sub> (3)	toluene	110	72	<b>1</b> a (40), <b>3</b> (43)
	P(c-C <sub>6</sub> H <sub>11</sub> ) <sub>3</sub> <sup>[f]</sup>	Et₃N (10)				
8	Pd(OAc) <sub>2</sub>	K <sub>3</sub> PO <sub>4</sub> (3)	toluene	110	72	<b>1</b> a (72), <b>3</b> (15)
	P(o-tolyl) <sub>3</sub> <sup>[f]</sup>	Et <sub>3</sub> N (10)				
9 <sup>[g]</sup>	[Pd(PPh <sub>3</sub> ) <sub>4</sub> ]	K <sub>3</sub> PO <sub>4</sub> (3)	toluene	110	72	<b>2</b> (26), <b>3</b> (64)
		Et <sub>3</sub> N (10)				

[a] Catalyst loading: 0.2 equivalents of Pd with respect to **1a**. [b] All reactions were carried out in a sealed tube. [c] Yields of the isolated products. [d] Compound **1a** was also recovered (10%). [e] Methyl 3-(N-(4-methylphenyl)amino)propionate was also isolated in 5% yield. [f] Ligand/Pd 2:1. [g] Bromide **1b** was used instead of **1a**. dba = dibenz-ylideneacetone.

combination that was effective in promoting nucleophilic addition in  $\beta$ -(2-iodoanilino)ketones,<sup>[3c]</sup> resulted exclusively in the reduction of the aryl halide (Table 1, entry 1).<sup>[11]</sup> In contrast, when K<sub>3</sub>PO<sub>4</sub> was used instead of Cs<sub>2</sub>CO<sub>3</sub>, a mixture of the reduction product **3** and ketone **2** was obtained. The latter compound results from nucleophilic substitution at the alkoxycarbonyl group (Table 1, entry 2). During our studies



© 2007 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

toward the optimization of the cyclization process, we found that the use of toluene in place of THF and in combination with Et<sub>3</sub>N led to an increase in the yield of ketone **2** to 65 % (Table 1, entry 3). The use of higher reaction temperatures resulted in an increase in the amount of the reduction product obtained (Table 1, entry 4), whereas low reaction rates were observed at lower reaction temperatures (Table 1, entry 5). The addition of Ag<sub>3</sub>PO<sub>4</sub> to remove the iodide ligand and facilitate the coordination of the carbonyl group (Table 1, entry 6) resulted in the formation of reduction products only, as did the use of other ligands on the palladium catalyst (Table 1, entries 7 and 8). Finally, the substitution reaction was less efficient with bromide **1b** as the substrate: Under the same reaction conditions, **3** was obtained as the main product (Table 1, entry 9).

A range of differently substituted dihydroquinolin-4-ones were synthesized under the optimized reaction conditions (Table 2). In general, substituents on the aromatic ring were

Table 2: Synthesis of substituted dihydroquinolin-4-ones.[a]



[a] Reaction conditions:  $[Pd(PPh_3)_4]$  (0.2 equiv),  $K_3PO_4$  (3 equiv),  $Et_3N$  (10 equiv), toluene, 72 h, 110 °C (sealed tube). [b] Yield of the isolated product. [c] Product of hydrodehalogenation (for details, see the Supporting Information). [d] Methyl 4-(methylamino)benzoate was also isolated. [e]  $[Pd(PPh_3)_4]$ : 0.1 equivalents. [f] Reaction time: 40 h.

found to have little effect on the success of the carbopalladation reaction. Thus, the nucleophilicity of the aryl palladium species does not appear to be affected by the electronic properties of the substituent. The low yield of the acylation product derived from **4b** is mainly a consequence of an increase in the rate of the competitive retro-Michael fragmentation of the  $\beta$ -aminoester (Table 2, entry 2). We found that benzyl esters can also be used as substrates in the reaction, although the desired product was formed in slightly lower yield from the benzyl ester counterpart of **1a** than from the methyl ester **1a** itself (compare Table 2, entry 3 with Table 1, entry 3).

The reaction of amino esters without hydrogen atoms  $\alpha$  to the carbonyl group proceeded smoothly to give the corresponding ketones in high yields (Table 2, entries 5-11), even when the amount of catalyst used was decreased to 0.1 equivalents (Table 2, entry 7). On the other hand, no competition between nucleophilic attack at the carbonyl group and  $\alpha$  arylation was observed in the reactions of amino esters 1a, 4a-c, and 7, which contain hydrogen atoms  $\alpha$  to the carbonyl group. The Pd-catalyzed intramolecular  $\alpha$  arylation of a-amino esters was described recently for substrates with (2-halobenzyl)amino or (2-(2-halophenyl)ethyl)amino moieties:<sup>[9]</sup> No interference from nucleophilic attack at the ester group was observed.<sup>[12]</sup> As the main difference between these amino esters and those examined in the present study is that there are no carbon atoms between the nitrogen atom and the aromatic ring, our initial assumption that the intermediacy of four-membered azapalladacycles could force otherwise unfavorable processes may be accurate. In fact, carbamate 14a and sulphonamide 14b, in which the coordination ability of the N atom has been suppressed, failed to undergo nucleophilic attack at the ester group, and the reduction products 15a (87%) and 15b (63%; Ts = p-toluenesulfonyl), respectively, were formed exclusively. Moreover, the treatment of 1a with [Pd<sub>2</sub>(dba)<sub>3</sub>] and PPh<sub>3</sub> in benzene at room



temperature for 24 h afforded the four-membered azapalladacycle 16 (see Scheme 1), which was transformed into ketone 2 together with ester 3 when heated at 110 °C in toluene in the presence of  $K_3PO_4$ .

A plausible mechanism for the observed  $Pd^{0}$ -catalyzed cyclization<sup>[13]</sup> is shown in Scheme 1. The oxidative addition of the aryl iodide to a  $Pd^{0}$  species would give the four-membered azapalladacycle **16**. A carbopalladation reaction between the



Scheme 1. Proposed mechanism for the acylation reaction.

## Communications

σ-aryl palladium moiety and the alkoxycarbonyl group would then lead to the Pd<sup>II</sup> alkoxide **A**. The coordination of the N atom to the palladium center in **16** not only brings the carbonyl group nearer to the metal to facilitate the formation of a transient chelated intermediate in which the carbonyl group is coordinated to the palladium center,<sup>[14]</sup> but also increases the electron density on the palladium center to enable the otherwise unfavorable carbopalladation reaction to occur. β-Alkoxide elimination from **A** would afford ketone **2** and a Pd<sup>II</sup> alkoxide, which would finally undergo β-hydride elimination to regenerate the Pd<sup>0</sup> catalyst. The isolation of significant amounts of benzaldehyde in the reaction of benzyl ester **4c** supports this reduction sequence.<sup>[15]</sup>

In summary, we have described the first examples of a palladium-catalyzed acylation of aryl iodides by alkoxycarbonyl groups. Further studies to expand the intramolecular reaction and provide deeper insight into the mechanism are under way.

## **Experimental Section**

Typical procedure: A mixture of the amino ester (0.04 M),  $K_3\text{PO}_4$  (3 equiv), Et<sub>3</sub>N (10 equiv), and [Pd(PPh\_3)\_4] (20 mmol%) in toluene was stirred at 110 °C in a sealed tube for 72 h. The reaction mixture was then poured into water, and the resulting mixture was extracted with Et<sub>2</sub>O. The organic extracts were washed with brine, dried, and concentrated. The residue was purified by flash chromatography to give the corresponding dihydroquinolin-4-one.

Received: May 17, 2007 Published online: August 14, 2007

**Keywords:** acylation · cyclization · nitrogen heterocycles · nucleophilic substitution · palladium

- Handbook of Organopalladium Chemistry for Organic Synthesis, Vols. I and II (Ed.: E. Negishi), Wiley-VCH, New York, 2002.
- [2] For the reaction of aryl and vinyl palladium complexes with aldehydes, see: a) R. C. Larock, M. J. Doty, S. Cacchi, *J. Org. Chem.* **1993**, *58*, 4579–4583; b) V. Gevorgyan, L. G. Quan, Y. Yamamoto, *Tetrahedron Lett.* **1999**, *40*, 4089–4092; c) L. Zhao, X. Lu, *Angew. Chem.* **2002**, *114*, 4519–4521; *Angew. Chem. Int. Ed.* **2002**, *41*, 4343–4345; d) J. Vicente, J.-A. Abad, B. López-Peláez, E. Martínez-Viviente, *Organometallics* **2002**, *21*, 58–67.
- [3] For the reaction of aryl and vinyl palladium complexes with ketones, see: a) L. G. Quan, V. Gevorgyan, Y. Yamamoto, J. Am. Chem. Soc. 1999, 121, 3545–3546; b) L. G. Quan, M. Lamrani, Y. Yamamoto, J. Am. Chem. Soc. 2000, 122, 4827–4828; c) D. Solé, L. Vallverdú, X. Solans, M. Font-Bardia, J. Bonjoch, J. Am. Chem. Soc. 2003, 125, 1587–1594; d) G. Liu, X. Lu, J. Am. Chem. Soc. 2006, 128, 16504–16505; see also ref. [2c–d].

- [4] For the reaction of aryl palladium complexes with acetic anhydride, see: S. Cacchi, G. Fabrizi, F. Gavazza, A. Goggiamani, Org. Lett. 2003, 5, 289–291.
- [5] For the reaction of aryl and vinyl palladium complexes with nitriles, see: a) C.-C. Yang, P.-J. Sun, J.-M. Fang, J. Chem. Soc. Chem. Commun. 1994, 2629–2630; b) R. C. Larock, Q. Tian, A. A. Pletnev, J. Am. Chem. Soc. 1999, 121, 3238–3239; c) A. A. Pletnev, Q. Tian, R. C. Larock, J. Org. Chem. 2002, 67, 9276–9287; d) A. A. Pletnev, R. C. Larock, J. Org. Chem. 2002, 67, 9428–9438; e) Q. Tian, A. A. Pletnev, R. C. Larock, J. Org. Chem. 2002, 67, 9428–9438; e) Q. Tian, A. A. Pletnev, R. C. Larock, J. Org. Chem. 2003, 68, 339–347; f) C. Zhou, R. C. Larock, J. Am. Chem. Soc. 2004, 126, 2302–2303; g) C. Zhou, R. C. Larock, J. Org. Chem. 2006, 71, 3551–3558; h) B. Zhao, X. Lu, Tetrahedron Lett. 2006, 47, 6765–6768; see also ref. [2c].
- [6] The few Pd-mediated reactions in which an ester functionality is modified do not involve the carbopalladation of the carbonyl group; see, for example: a) W. Tao, L. J. Silverberg, A. L. Rheingold, R. F. Heck, *Organometallics* **1989**, *8*, 2550–2559; b) R. C. Larock, X. Han, M. J. Doty, *Tetrahedron Lett.* **1998**, *39*, 5713–5716.
- [7] For recent reviews, see: a) G. C. Lloyd-Jones, Angew. Chem.
  2002, 114, 995-998; Angew. Chem. Int. Ed. 2002, 41, 953-956;
  b) D. A. Culkin, J. F. Hartwig, Acc. Chem. Res. 2003, 36, 234-245.
- [8] a) W. A. Moradi, S. L. Buchwald, J. Am. Chem. Soc. 2001, 123, 7996-8002; b) S. Lee, N. A. Beare, J. F. Hartwig, J. Am. Chem. Soc. 2001, 123, 8410-8411; c) M. Jorgensen, S. Lee, X. Liu, J. P. Wolkowski, J. F. Hartwig, J. Am. Chem. Soc. 2002, 124, 12557-12565; d) T. Hama, X. Liu, D. A. Culkin, J. F. Hartwig, J. Am. Chem. Soc. 2003, 125, 11176-11177; e) X. Liu, J. F. Hartwig, J. Am. Chem. Soc. 2004, 126, 5182-5191.
- [9] O. Gaertzen, S. L. Buchwald, J. Org. Chem. 2002, 67, 465-475.
- [10] a) A. B. Daruwala, J. E. Gearien, W. J. Dunn III, P. S. Benoit, L. Bauer, J. Med. Chem. 1974, 17, 819–824; b) K. Nishijima, T. Shinkawa, Y. Yamashita, N. Sato, H. Nishida, K. Kato, Y. Onuki, M. Mizota, K. Ohmoto, S. Miyano, Eur. J. Med. Chem. 1998, 33, 267–277.
- [11] The reduction is a common side reaction in such processes; see, for example, refs. [3b,c,5d].
- [12] The Pd<sup>0</sup>-catalyzed intramolecular coupling of amino-tethered vinyl halides with esters also proceeds without any interference from nucleophilic attack at the ester group; the addition of a phenoxide to the reaction mixture is necessary to promote the reaction at the  $\alpha$  position; see: D. Solé, X. Urbaneja, J. Bonjoch, *Adv. Synth. Catal.* **2004**, *346*, 1646–1650.
- [13] For a related rhodium(I)-catalyzed transformation, see: T. Miura, T. Sasaki, H. Nakazawa, M. Murakami, J. Am. Chem. Soc. 2005, 127, 1390-1391.
- [14] For the intramolecular coordination of the ester carbonyl group to Pd<sup>II</sup>, see: S. Gosiewska, M. Huis in't Veld, J. J. M. de Pater, P. C. A. Bruijnincx, M. Lutz, A. L. Spek, G. van Koten, R. J. M. Klein Gebbink, *Tetrahedron: Asymmetry* **2006**, *17*, 674–686.
- [15] The involvement of other reductants (e.g. Et<sub>3</sub>N) can not be excluded completely; see: S.-I. Murahashi, T. Hirano, T. Yano, J. Am. Chem. Soc. 1978, 100, 348–350.