




Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/lncn20>

DESIGN AND SYNTHESIS OF NOVEL FLUOROCYCLOPROPANOID NUCLEOSIDES

Young Rang Lee ^a, Ju-Hyun Park ^a, Raok Jeon ^a, Lak Shin Jeong ^b, Moon Woo Chun ^c & Hee-Doo Kim ^c

^a College of Pharmacy, Sookmyung Women's University, Seoul, 140-742, Korea

^b College of Pharmacy, Ewha Womans University, Seoul, 120-750, Korea

^c College of Pharmacy, Seoul National University, Seoul, 151-742, Korea

Published online: 07 Feb 2007.

To cite this article: Young Rang Lee, Ju-Hyun Park, Raok Jeon, Lak Shin Jeong, Moon Woo Chun & Hee-Doo Kim (2001) DESIGN AND SYNTHESIS OF NOVEL FLUOROCYCLOPROPANOID NUCLEOSIDES, *Nucleosides, Nucleotides and Nucleic Acids*, 20:4-7, 677-679

To link to this article: <http://dx.doi.org/10.1081/NCN-100002349>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

DESIGN AND SYNTHESIS OF NOVEL FLUOROCYCLOPROPANOID NUCLEOSIDES

Young Rang Lee,¹ Ju-Hyun Park,¹ Raok Jeon,¹ Lak Shin Jeong,²
Moon Woo Chun,³ and Hee-Doo Kim³

¹College of Pharmacy, Sookmyung Women's University,
Seoul 140-742, Korea

²College of Pharmacy, Ewha Womans University, Seoul 120-750, Korea

³College of Pharmacy, Seoul National University, Seoul 151-742, Korea

ABSTRACT

Novel fluorocyclopropanoid nucleosides were designed, synthesized and evaluated their antiviral activities against poliovirus, HSV, HIV, and HBV.


Despite the intensive synthetic efforts, cyclopropanoid nucleosides have not been of interest until A-5021 was discovered recently (1). A-5021, (1'S,2'R)-9-[[1',2'-bis(hydroxymethyl)cycloprop-1'-yl]methyl]guanine, was reported more active than acyclovir or penciclovir against HSV, HCMV and VZV (2). In an effort to search for the chemically and enzymatically stable carbonucleoside, we designed (*E*)-(1'-fluoro-2'-hydroxymethylcyclopropylmethyl)purines and pyrimidines. The underlying concept for our design is to seek relatively conformationally-locked compound with minimal structural disturbance from acyclic carbonucleoside such as acyclovir or penciclovir.

To meet such a requirement, we need to introduce cyclopropane and fluorine moiety. Due to its hybrid character of cyclic and acyclic molecules, cyclopropyl group could render the conformational rigidity to the target molecule. It is also envisioned that fluorine could play a role to control the conformation by *gauche* effect between fluorine and nitrogen in nucleoside base moiety (3). As shown in Scheme 1, N-alkylation of purine or pyrimidine bases with the mesylate **7** or the iodide **8** under basic condition (NaH, DMF, 80°C) was chosen as a simple and convenient approach to the synthesis of the fluorocyclopropanoid nucleoside **9**. The key



Figure 1.

synthetic intermediate **7** or **8** were synthesized starting from allyl alcohol. Protection of allyl alcohol with TBDPSCl, followed by ozonolysis afforded the aldehyde **2** in good yield. Introduction of fluorine and double bond for the installation of cyclopropyl group was achieved by Horner-Wadsworth-Emmons reaction of the aldehyde **3** with triethyl 2-fluoro-2-phosphonoacetate using *n*-BuLi in THF. The α -fluoro- α , β -unsaturated ester **3** with (*E*)-stereochemistry was formed preferentially over **4** in a ratio of 95 to 5. Unusual preference of (*E*)-isomer is well-known phenomenon in Horner-Wadsworth-Emmons reaction of aldehydes with triethyl 2-fluoro-2-phosphonoacetate (4). The similar stereoselectivity was also observed when using *Pr*MgBr in THF at 0°C. The relatively unstable ester **3** was directly subjected to DIBAL-H reduction in CH₂Cl₂ at -78°C to room temperature to furnish the allylic alcohol **5** in 71% yield. Conventional cyclopropanation of the allylic



Scheme 1.



alcohol **5** using Simmons-Smith or Furukawa modification method proceeded too slowly, and provided the desired product **6** in low yield. However, encouraged with the previous reports on Lewis acid-catalyzed cyclopropanation (**5**), we had tried several kinds of Lewis-acid, then, finally selected ZnI_2 . The tendency of alkylzinc alkoxides (ROZnEt) to aggregate in solution (**6**) could be circumvented by sonification. By the dual action of sonification and ZnI_2 , chemical yield and the reaction rate were greatly improved.

Unseparable mixture of **5** and **6** by conventional chromatography could be isolated using radial chromatography. The fluorinated cyclopropyl alcohol **6** was, then, converted to the corresponding mesylate **7** or iodide **8**. The coupling **7** or **8** with purine or pyrimidine bases under basic conditions, followed by removal of TBDPS group to afford the desired nucleosides **10a** (81% from **8**), **10b** (46% from **7**), **10c** (40% from **8**), and **10d** (40% from **8**), respectively. Treatment of **10d** with 2-mercaptoethanol and sodium methoxide in methanol, followed by hydrolysis with glacial acetic acid gave **10e** in 14% yield. The synthesized nucleosides (**10a~e**) were evaluated for their antiviral activity against poliovirus, HSV-1, HSV-2, HBV and HIV. However, all compounds were found to be inactive in the assay.

ACKNOWLEDGMENT

This research was supported by the grant from the Good Health R & D project (HMP-98-D-4-0057), Ministry of Health and Welfare, Korea.

REFERENCES

1. Rife, J.; Ortuno, R. M. *Org. Lett.*, **1999**, 1, 1221–1223.
2. Sekiyama, T.; Hatsuya, S.; Tanaka, Y.; Uchiyama, M.; Ono, N.; Iwayama, S.; Oikawa, M.; Suzuki, K.; Okunishi, M.; Tsuji, T. *J. Med. Chem.*, **1998**, 41, 1284–1298.
3. Pankiewicz, K. W. *Carbohydr. Res.*, **2000**, 327, 87–105.
4. Sano, S.; Ando, T.; Yokoyama, K.; Nagao, Y. *Synlett*, **1998**, 777–779.
5. Charette, A. B.; Brochu, C. *J. Am. Chem. Soc.*, **1995**, 117, 11367–11368.
6. Denmark, S.E.; O'Connor S. P. *J. Org. Chem.*, **1997**, 62, 584–594.



Request Permission or Order Reprints Instantly!

Interested in copying and sharing this article? In most cases, U.S. Copyright Law requires that you get permission from the article's rightsholder before using copyrighted content.

All information and materials found in this article, including but not limited to text, trademarks, patents, logos, graphics and images (the "Materials"), are the copyrighted works and other forms of intellectual property of Marcel Dekker, Inc., or its licensors. All rights not expressly granted are reserved.

Get permission to lawfully reproduce and distribute the Materials or order reprints quickly and painlessly. Simply click on the "Request Permission/Reprints Here" link below and follow the instructions. Visit the [U.S. Copyright Office](#) for information on Fair Use limitations of U.S. copyright law. Please refer to The Association of American Publishers' (AAP) website for guidelines on [Fair Use in the Classroom](#).

The Materials are for your personal use only and cannot be reformatted, reposted, resold or distributed by electronic means or otherwise without permission from Marcel Dekker, Inc. Marcel Dekker, Inc. grants you the limited right to display the Materials only on your personal computer or personal wireless device, and to copy and download single copies of such Materials provided that any copyright, trademark or other notice appearing on such Materials is also retained by, displayed, copied or downloaded as part of the Materials and is not removed or obscured, and provided you do not edit, modify, alter or enhance the Materials. Please refer to our [Website User Agreement](#) for more details.

[Order now!](#)

Reprints of this article can also be ordered at

<http://www.dekker.com/servlet/product/DOI/101081NCN100002349>