Cite this: Org. Biomol. Chem., 2012, 10, 970

Dynamic Article Links 🕟



The reaction of β-lactam carbenes with 3,6-dipyridyltetrazines: switch of reaction pathways by 2-pyridyl and 4-pyridyl substituents of tetrazines[†]

Xiao-Rong Wang, Juan Xing, Cai-Xia Yan and Ying Cheng*

Received 19th September 2011, Accepted 26th October 2011 DOI: 10.1039/c1ob06595e

The reactions of β -lactam carbenes with both 3,6-di(2-pyridyl)tetrazine and 3,6-di(4-pyridyl)tetrazine were studied. It was found that β -lactam carbenes reacted with 3,6-di(2-pyridyl)tetrazine to produce 5-triazolo[1,5-*a*]pyridylpyrrol-2-ones in good yields, while with 3,6-di(4-pyridyl)tetrazine, they afforded pyrido[*c*]cyclopenta[*b*]pyrrol-2-ones in moderate yields. Both reactions were proposed to follow cascade mechanisms containing a 3,6*a*-dipyridylpyrrolo[3,2-*c*]pyrazol-5-one intermediate. The different pathways of the transformation of pyrrolo[3,2-*c*]pyrazol-5-ones were switched by the 2- and 4-pyridyl substituents. This work not only provided a simple and efficient strategy for the construction of novel triazolo[1,5-*a*]pyridine and pyrido[*c*]cyclopenta[*b*]pyrrole derivatives, respectively, but also revealed two different thermal transformation patterns of 3*H*-pyrazole compounds.

Introduction

β-Lactam carbenes, namely 2-azetidinone-4-ylidenes, were firstly generated from thermolysis of spiro[β-lactam-4,2'-oxadiazolines] by Warkentin and co-workers in the 1990's.^{1,2} They investigated the reactions of β -lactam carbenes with both electron-rich and electron-deficient alkenes or alkynes to produce β-lactam-spirocyclopropane or β -lactam-spiro-cyclopropene derivatives, and demonstrated the β -lactam carbenes to be ambiphilic carbenes but with pronounced electrophilic activity.^{3,4} We have been interested in the chemistry of β -lactam carbenes, not only in their ambiphilic reactivity but more importantly in their synthetic applications.⁵⁻⁹ We demonstrated that β -lactam carbenes are unique intermediates in the construction of mono-, spiro- or fused heterocyclic compounds with or without a β -lactam moiety. For example, β lactam carbenes underwent nucleophilic reaction toward aryl isocyanates to produce β-lactam-spiro-indole-2-ones.⁵ On the other hand, β -lactam carbenes underwent electrophilic addition to aryl isonitriles to afford high yields of 2-(\beta-lactam-4-ylidene)indoles that rearranged into δ -carbolin-2-ones in the presence of an acid.⁶ While the same carbenes reacted with alkyl isonitriles, 4-cyano or 4-carbamoyl substituted β -lactams were obtained depending on the alkyl groups of isonitriles.⁷ Very recently, we found that the reaction of β -lactam carbenes with 3,6-diphenyl-1,2,4,5-tetrazines proceeded in a five-step cascade process to produce indeno[2,1b]pyrrol-2-ones in good yields.8 We envisioned that, by varying the aryl substituents of 3,6-diaryltetrazines, the reaction between

Downloaded by Pennsylvania State University on 02 July 2012 Published on 26 October 2011 on http://pubs.rsc.org | doi:10.1039/C1OB06595E

β-lactam carbenes and 3,6-diaryltetrazines might be a general method for the synthesis of aryl-fused cyclopenta[*b*]pyrrol-2-one compounds. Thus, we undertook the current study on the reaction of β-lactam carbenes with 3,6-di(2-pyridyl)tetrazines and 3,6-di(4-pyridyl)tetrazines. Interestingly, the similar reactions of β-lactam carbenes with di(2-pyridyl)- and with di(4-pyridyl)tetrazines produced totally different products, 5-triazolo[1,5-*a*]pyridylpyrrol-2-ones or pyrido[*c*]cyclopenta[*b*]pyrrol-2-ones, *via* two different transformations of pyrrolo[3,2-*c*]pyrazol-5-one intermediates. Herein, we report our results.

Results and discussion

We first studied the reaction of β -lactam carbenes with di(2pyridyl)tetrazines. In practice, all β -lactam carbenes 2 are generated in situ by thermolysis of spiro[β -lactam-4,2'-oxadiazolines] 1, which were prepared on the basis of Warkentin's method^{1,2} and our previous report.^{5,9} Since the optimal temperature for the generation of carbenes 2 from spiro-oxadiazolines 1 is around 100-110 °C according to Warkentin's reports¹⁻⁴ and our experience,⁵⁻⁹ the reaction between 3,3-dimethyl-1-(p-methoxyphenyl)-2-azetidinone-4-ylidene 2c and 3,6-di(2-pyridyl)tetrazine 3 (2c:3 = 1:1) was initially examined in refluxing 1,4-dioxane, which produced 3,6adi(2-pyridyl)pyrrolo[3,2-c]pyrazol-5-one 4c as a major product in 40% yield along with 7% of minor product 5-triazolo[1,5*a*]pyridylpyrrol-2-one **5c**. The spectroscopic analysis indicated that 4c and 5c were isomeric products derived from 1 + 1 addition of carbene 2c with tetrazine 3 with the loss of one N_2 molecule. It was found that the reaction of β -lactam carbene 2c with di(2-pyridyl)tetrazine 3 firstly yielded product 4c, which was converted into 5c in prolonged time under heating. Thus, the reaction conditions for the predominant formation of 4c or 5c were optimized by varying the ratio of starting materials,

College of Chemistry, Beijing Normal University, Beijing, 100875, China. E-mail: ycheng2@bnu.edu.cn

[†] Electronic supplementary information (ESI) available: NMR data and crystal structure data for compounds **4c**, **5c** and **8i**. CCDC reference numbers 844397–844399. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c1ob06595e

 Table 1
 Optimization of reaction conditions for the predominant formation of 4c or 5c from interaction of 3,3-dimethyl-1-(p-methoxyphenyl)-2azetidinone-4-ylidene 2c with 3,6-di(2-pyridyl)tetrazine 3



	Reactio	Yield				
Entry	1c:3	Solvent	Temp.	Time (h)	4c	5c
1	1:1	1,4-Dioxane	100	9	40	7
2	1.5:1	1,4-Dioxane	100	9	60	10
3	2:1	1,4-Dioxane	100	5	70	15
4	1.5:1	<i>n</i> -Heptane	100	12	22	49
5	1.5:1	Toluene	100	9	50	11
6	1.5:1	Toluene	110	16	28	60
7	1.5:1	1,1,2-Trichloroethane	110	24	_	43
8	1.5:1	Xylene	140	5	_	92

reaction temperature and solvents that were chemically inert towards carbenes 2 (Table 1). By using excess amount of carbene precursor 1c (1c: 3 = 2:1) to promote the interaction of carbene with tetrazine and controlling both reaction time and temperature to inhibit the transformation of 4c to 5c, 70% yield of 4c was obtained from the reaction in 1,4-dioxane at 100 °C (Table 1, entry 3). On the other hand, it was noted that increasing reaction temperature in toluene significantly improved the yield of 5c (Table 1, entries 5 and 6). While further elevating reaction temperature in xylene, to our delight, product 5c was obtained in 92% yield from the reaction conducted in refluxing xylene (Table 1, entry 8).

The generality of the reaction was studied, respectively, under the optimized conditions for the selective formation of products 4 or 5. The reaction of β -lactam carbones that were attached by different substituents with di(2-pyridyl)tetrazine 3 was firstly examined at 100 °C in 1,4-dioxane. As summarized in Table 2, the *N*-aryl group of β -lactam carbenes **2** showed little effects on the reaction. By keeping reaction temperature at 100 °C and time in 3.5-5 h, all reactions of N-phenyl, N-(4-methylphenyl), N-(4-methoxyphenyl), N-(4-chlorophenyl) and N-(4-bromophenyl) substituted β -lactam carbenes 2a-2e with di(2-pyridyl)tetrazine 3 produced pyrrolo[3,2-c]pyrazol-5-ones 4a-4e as major products in 70-83% yields, along with a small amount of byproducts 5a-5e (Table 2, entries 1–5). However, the 3,3-alkyl substituents of carbene reactants have influence on the efficiency of both the reaction between carbenes 2 and tetrazine 3, and the transformation from products 4 to 5. For example, the reaction between 3,3-cyclohexyl- β -lactam carbene 2h and tetrazine 3 proceeded somewhat more slower than that of methyl or/and ethyl substituted carbenes, probably due to the steric hindrance between two reactants (Table 2, entry 8). While all 3,3-dimethyl-β-lactam carbenes 2a-2e reacted with tetrazine 3 to afford 4a-4e as major products in good yields, the reaction of 3,3-diethyl- and 3-ethyl-3-methyl substituted carbenes 2f and 2g with 3 afforded 5f and 5g, respectively, as major products under the similar conditions, which indicated the isomerization of products 4f and 4g being faster than that of 4a-4e. To selectively obtain compounds 5, the reaction of β -

$X \xrightarrow{\begin{tabular}{c} N \\ N \\ N \\ 1 \\ 1 \\ 3 \\ \begin{tabular}{c}{3}{1} \\ 1.4 \\ -1.4$					
				Yield (%)
Entry	1	$\mathbf{X}, \mathbf{R}^1, \mathbf{R}^2$	Time (h)	4	5
1 2 3 4 5 6 7 8	1a 1b 1c 1d 1e 1f 1g 1h	$\begin{array}{c} H, CH_3, CH_3\\ CH_3, CH_3, CH_3\\ OCH_3, CH_3, CH_3\\ Cl, CH_3, CH_3\\ Br, CH_3, CH_3\\ CH_3, C_2H_5, C_2H_5\\ CH_3, CH_3, C_2H_5\\ CH_3, CH_3, C_2H_5\\ CH_3, (CH_2)_5 \end{array}$	4 5 3.5 4 3.5 4 6	4a: 83 4b: 77 4c: 70 4d: 82 4e: 73 4f: 25 4g: 37 ^b 4h: 88	5a : 6 5b : 10 5c : 15 — ^{<i>a</i>} 5f : 55 5g : 44 — ^{<i>a</i>}

Table 2 The reaction of spiro[β -lactam-4,2'-oxadiazolines] 1 and di(2-

pyridyl)tetrazine 3 in 1,4-dioxane at 100 °C

^{*a*} A little amount of byproduct **5** was detected by TLC without isolation. ^{*b*} **4g** is a mixture of diastereomers that could not be separated by column chromatography.

Table 3 The reaction of spiro[β -lactam-4,2'-oxadiazolines] 1 and di(2-pyridyl)tetrazine 3 in refluxing xylene

x-{[$ \begin{array}{c} $		2
				Yield (%)
Entry	1	$\mathbf{X}, \mathbf{R}^1, \mathbf{R}^2$	Time (h)	5
1 2 3 4 5 6 7 8	1a 1b 1c 1d 1e 1f 1g 1h	$\begin{array}{c} H, CH_3, CH_3\\ CH_3, CH_3, CH_3\\ OCH_3, CH_3, CH_3\\ Cl, CH_3, CH_3\\ Br, CH_3, CH_3\\ CH_3, C_2H_5, C_2H_5\\ CH_3, CH_3, C_2H_5\\ CH_3, CH_3, C_2H_5\\ CH_3, CH_2)_5 \end{array}$	5 4 4 4 4 4 3 12	5a: 87 5b: 83 5c: 92 5d: 91 5e: 80 5f: 82 5g: 90 5h: 75

lactam carbenes **2** with di(2-pyridyl)tetrazine **3** was then conducted under the optimized conditions for the formation of 5-triazolo[1,5*a*]pyridylpyrrol-2-ones **5**. As indicated in Table 3, in refluxing xylene, all carbenes **2a–2h** bearing different substituents reacted with tetrazine **3** to afford pyrrol-2-ones **5** in good to excellent yields, although the reaction between 3,3-cyclohexyl substituted β -lactam carbene **2h** and tetrazine **3** proceeded more slowly than others.

Following the reaction of β -lactam carbenes 2 with di(2pyridyl)tetrazine 3, the behavior of carbenes 2 towards di(4pyridyl)tetrazine 6 was also investigated. The optimized conditions for the reaction of carbenes 2 with di(2-pyridyl)tetrazine 3 were firstly employed in the reaction of β -lactam carbene 2c with 6

7

8

 $1 \cdot 1$

1.5:1

 $1.5 \cdot 1$

Toluene

Toluene

1,1,2-Trichloroethane



Table 4 Reaction of β -lactam carbene **2c** with 3,6-di(4-pyridyl)tetrazine

^a When two reactants were employed in the ratio of 1:1, partial tetrazine 6 was not consumed.

110

110

110

di(4-pyridyl)tetrazine 6. It was found that the reaction of 2c with 6 (2c:6 = 1.5:1) produced two compounds 7c and 8c with low selectivity in both refluxing 1,4-dioxane and xylene (Table 4, entries 1 and 3). After the spectroscopic analyses, product 7c was identified as 3,3-dimethyl-1-(p-methoxyphenyl)-8-(4-pyridyl)pyrido[c]cyclopenta[b]pyrrol-2-one, and 8c was a Nβ-lactam substituted derivative of 7c. Based on the structures of two products, product $\mathbf{8c}$ was apparently resulted from the N–H insertion of β -lactam carbene 2c to product 7c. We considered that if product 7c could be removed from the reaction mixture during the reaction process, the formation of 8c would be limited and then the yield of 7c might be improved. It was noted that 7c has very low solubility in many common organic solvents, such as heptane, toluene, dioxane, chloroform, ethyl acetate, acetone and ethanol. The use of low polarity solvents led to the precipitation of 7c from solvent during reaction. Thus, the reaction of carbene 2c with di(4-pyridyl)tetrazine 6 was examined in refluxing toluene and *n*-heptane, respectively. Although the reaction in toluene afforded a mixture of 7c and 8c in a ratio about 2:1 (Table 4, entry 7), product 7c was obtained in 65% yield from the reaction in *n*-heptane and almost no 8c was isolated (Table 4, entry 5). Comparing the reactions of carbene 2c with tetrazine 6(2c:6 =1.5:1) in dioxane, toluene and *n*-heptane, it was found that the reaction in dioxane or toluene gave higher total yields of products (7c + 8c) than that in *n*-heptane, but with lower selectivity in the formation of 7c versus 8c. On the other hand, the reactions in toluene and *n*-heptane preferred to the formation of 7c, while 8c was isolated as major product in dioxane. The lower reactivity and higher selectivity of the reaction in *n*-heptane can be explained by the solubility of reactants or products in solvent. While the low solubility of di(4-pyridyl)tetrazine 6 in *n*-heptane decreased the reaction efficiency between carbene 2c and tetrazine 6, the non-solubility of product 7c in *n*-heptane inhibited the reaction of carbene 2c with 7c. The reversed selectivity of reaction in dioxane and toluene was also attributed to the different solubilities of 7c in these two solvents (Table 4, entries 1 and 7). Since 1,4dioxane was the optimal solvent for the formation of product 8c and it was derived from 2 + 1 addition of carbene 2c with tetrazine 6, two equivalents of 2c were then employed to react with 6 in refluxing 1,4-dioxane, which improved the yield of 8c to 78%.

The generality of the reaction between β -lactam carbenes 2 bearing different substituents and di(4-pyridyl)tetrazine 6 was studied in refluxing n-heptane and 1,4-dioxane, respectively. It was found that the substituents of carbenes showed small influence on the reaction. As shown in Table 5, all carbenes 2 bearing different aryl and alkyl substituents on the β -lactam rings reacted with di(4-pyridyl)tetrazine 6 in refluxing *n*-heptanes to afford pyrido[c]cyclopenta[b]pyrrol-2-ones 7 in moderate yields. While

Table 5 Reaction of β -lactam carbenes 2 with 3,6-di(4-pyridyl)tetrazine 6 in refluxing *n*-heptane or 1,4-dioxane

10

10

12

624

65

36

31

5

	$X \longrightarrow \begin{bmatrix} R^{1} \\ N \\ N \\ N \\ N \\ N \\ N \\ 1 \\ 6 \\ \end{bmatrix} \begin{pmatrix} R^{1} \\ N \\ N \\ R^{2} \\ N \\ N \\ N \\ R^{2} \\ N \\ N \\ R^{2} \\ N \\ N \\ N \\ R^{2} \\ N \\ $							
							Yield (%)	
Entry	1	$\mathbf{X}, \mathbf{R}^1, \mathbf{R}^2$	1:6	Solvent	Temp.	Time (h)	7	8
1	1a	H, CH_3, CH_3	1.5:1	<i>n</i> -Heptane	Reflux	8	7a : 65	
2	1b	CH ₃ , CH ₃ , CH ₃	1.5:1	n-Heptane	Reflux	8	7b : 63	_
3	1c	OCH ₃ , CH ₃ , CH ₃	1.5:1	n-Heptane	Reflux	8	7c: 65	
4	1d	Cl, CH ₃ , CH ₃	1.5:1	n-Heptane	Reflux	8	7d : 60	
5	1f	CH_3, C_2H_5, C_2H_5	1.5:1	n-Heptane	Reflux	8	7f : 56	
6	1g	CH_3, CH_3, C_2H_5	1.5:1	n-Heptane	Reflux	8	7g: 51	
7	1a	H, CH_3, CH_3	2:1	1,4-Dioxane	Reflux	13	7a : 8	8a : 57
9	1b	CH ₃ , CH ₃ , CH ₃	2:1	1,4-Dioxane	Reflux	13	7 b: 7	8b : 70
9	1c	OCH ₃ , CH ₃ , CH ₃	2:1	1,4-Dioxane	Reflux	13	7c: 12	8c: 78
10	1d	Cl, CH ₃ , CH ₃	2:1	1,4-Dioxane	Reflux	10	7d : 14	8d : 45
11	1i	OCH_3, C_2H_5, C_2H_5	2:1	1,4-Dioxane	Reflux	11	_	8i : 79
12	1j	OCH_3 , $(CH_2)_4$	2:1	1,4-Dioxane	Reflux	18	7j : 5	8j : 65

...



Scheme 1 The proposed mechanisms for the reactions of β -lactam carbenes with 3,6-di(2-pyridyl)tetrazine and 3,6-di(4-pyridyl)tetrazine, respectively.

in refluxing 1,4-dioxane, the reaction of carbenes **2** with tetrazine **6** produced N- β -lactam substituted pyrido[c]cyclopenta[b]pyrrol-2-ones **8** in 45–79% yields, along with a small amount of products **7**.

The structures of all products 4, 5, 7 and 8 were ascertained by spectroscopic methods. ¹H NMR, MS and microanalysis indicated that both products 4 and 5 were the 1 + 1 adducts of β -lactam carbenes 2 and di(2-pyridyl)tetrazine 3 with the loss of one N₂ molecule, while products 7 were derived from the 1 + 1 addition of a carbene 2 and a di(4-pyridyl)tetrazine 6 with the loss of two N₂ molecules and products 8 were the 1 + 1 adducts of products 7 and carbenes 2. Since the spectroscopic data did not allow full verification of the structures, the structures of 4c, 5c and 8i were determined unambiguously by single crystal X-ray diffraction analysis[†].

According to literature, the reactions of nucleophilic or ambiphilic carbenes with 3,6-disubstituted tetrazines generally proceed *via* a [4 + 1] cycloaddition followed by a retro [4 + 2] Diels–Alder cycloaddition to produce pyrazole derivatives.¹⁰⁻¹³ In our previous study, we have demonstrated that the reaction of β -lactam carbenes with 3,6-diphenyltetrazines produce indeno[2,1-*b*]pyrrol-2-ones *via* the isolable β -lactam-spiro-pyrazole and pyrrolo[3,2-c]pyrazol-5-one intermediates.⁸ However, it is surprising that the reactions of β -lactam carbenes **2** with 3,6-di(2-pyridyl)tetrazine **3** and 3,6-di(4-pyridyl)tetrazine **6** produce totally different types of products, the bicyclic 5-triazolo[1,5-*a*]pyridylpyrrol-2-ones **5** and tricyclic pyrido[*c*]cyclopenta[*b*]pyrrol-2-ones **7**, respectively. On the basis of our knowledge of the interaction between β -lactam carbenes **2** and 3,6-diphenyltetrazine, two cascade mechanisms

are proposed, respectively, for the reactions of β -lactam carbenes 2 with 3.6-di(2-pyridyl)tetrazine 3 and 3.6-di(4-pyridyl)tetrazine 6. As illustrated in Scheme 1, a [4 + 1] cycloaddition of β -lactam carbenes 2 with 3,6-di(2-pyridyl)tetrazine 3 or 3,6di(4-pyridyl)tetrazine 6 forms the bridged compounds 9 or 10, which are converted to β -lactam-spiro-pyrazoles 11 or 12 by a retro-Diels-Alder reaction. Under the heating condition, spiropyrazoles 11 or 12 undergo a 1,5-sigmatropic rearrangement to produce isolable 3,6a-di(2-pyridyl)pyrrolo[3,2-c]pyrazol-5-ones 4 or 3,6a-di(4-pyridyl)pyrrolo[3,2-c]pyrazol-5-ones 13, respectively. The isomerization of pyrrolo[3,2-c]pyrazol-5-ones 4 to triazolo[1,5-a]pyridines 5 most probably proceeds via the breaking of pyrazole ring of 4 to form dipolar intermediates 14. Intramolecular cyclization of 14 between the diazo species and the nearby nitrogen atom of pyridyl affords triazolo[1,5-a]pyridines 5. In the reaction of carbenes 2 with 3,6-di(4-pyridyl)tetrazine 6, the intermediates 3,6a-di(4-pyridyl)pyrrolo[3,2-c]pyrazol-5-ones 13 are unstable and have not been isolated. Decomposition of the pyrazole rings of pyrrolo[3,2-c]pyrazol-5-ones 13 with the loss of one N₂ molecule probably forms the carbene intermediates 16B, which undergo an intramolecular insertion to the C-H bond of the pyridyl to afford pyrido[c]cyclopenta[b]pyrrol-2-ones 7.Insertion of β -lactam carbenes 2 to the N–H bond of products 7 produced N- β -lactam substituted pyrido[c]cyclopenta[b]pyrrol-2-ones 8. The different transformations of pyrrolo[3,2-c]pyrazol-5-one intermediates 4 and 13 can be explained by the different stabilities and structures of the dipolar intermediates 14 and 15 derived from breaking of pyrazole ring of 4 and 13. Since 2pyridyl has a stronger electron-withdrawing inductive effect than 4-pyridyl group, the anions of intermediates 14 might be more stable than 15. Meanwhile, the anions of 14 can be delocalized on the nitrogen atom of 2-pyridyl attached to the same carbon with the diazo group, and then catches the diazo species to form the triazole rings of products 5. On the contrary, the less stable dipolar intermediates 15 releases a N_2 molecule to form carbone intermediates 16B, which undergoes C–H insertion to the pyridyl to form the cyclopentadiene moieties of products 7.

Conclusions

In summary, we have studied the reactions of β -lactam carbenes with both 3,6-di(2-pyridyl)tetrazine and 3,6-di(4-pyridyl)tetrazine. Under the heating conditions, B-lactam carbenes reacted with 3,6di(2-pyridyl)tetrazine to produce 5-triazolo[1,5-a]pyridylpyrrol-2-ones in good yields, while with 3,6-di(4-pyridyl)tetrazine, pyrido[c]cyclopenta[b]pyrrol-2-ones were obtained in moderate yields. Both reactions proceeded via a 3,6a-dipyridylpyrrolo[3,2c]pyrazol-5-one intermediate. The transformation pathways of pyrrolo[3,2-c]pyrazol-5-one intermediates were switched by the 2pyridyl or 4-pyridyl substituents, which led to the formation of totally different types of products. This work not only provided a simple and efficient strategy for the construction of novel triazolo[1,5*a*]pyridine derivatives and the new pyrido[*c*]cyclopenta[*b*]pyrrole ring system, respectively, but also revealed two different thermal transformation patterns of 3H-pyrazole compounds that were regulated by their substituents.

Experimental

Melting points are uncorrected. ¹H NMR (400 MHz) and ¹³C NMR (100 MHz) were recorded in the indicated solvents. J values are reported in Hz. IR spectra were recorded using an AVATAR 360 FT-IR spectrometer. Mass spectra were recorded on a Surveyor MSQ Plus (ESI) instrument and elemental analyses were performed on a GMBH Vario EL instrument. Column chromatography was performed using 200–300 mesh silica gel.

General procedure for the reaction of β-lactam carbenes with 3,6-di(2-pyridyl)tetrazine in 1,4-dioxane

Under nitrogen atmosphere, the mixture of spiro[β -lactam-4,2'oxadiazolines] 1^{1,2,5,9} (2 mmol) and 3,6-di(2-pyridyl)tetrazine 3¹⁴ (1 mmol) was heated in slightly refluxing 1,3-dioxane (50 mL) for 3.5–6 h (the time for each reaction was listed in Table 2). After removal of the solvent under vacuum, the residue was subject to chromatography on a silica gel column eluting with a mixture of petroleum ether and ethyl acetate from 10:1 to pure ethyl acetate to give products 3,6*a*-di(2-pyridyl)pyrrolo[3,2-*c*]pyrazol-5-ones 4 in 25–88% yields and 5-triazolo[1,5-*a*]pyridylpyrrol-2ones 5 in 6–55% yields. Products 4 or 5 were further purified by recrystallization in ethyl acetate and petroleum ether.

6,6-Dimethyl-4-phenyl-3,6*a*-di(2-pyridyl)-6,6*a*-dihydropyrrolo-[**3,2**-*c*]pyrazol-5-one 4a. 83%, mp 152–153 °C; IR ν (cm⁻¹) 1757, 1635, 1583, 1500; ¹H NMR (400 MHz, CDCl₃) δ (ppm) 8.46 (d, J = 4.1 Hz, 1H), 8.11 (d, J = 4.2 Hz, 1H), 7.81 (d, J = 8.0 Hz, 1H), 7.67 (td, J = 7.8, 1.8 Hz, 1H), 7.56–7.61 (m, 2H), 7.38–7.40 (m, 3H), 7.36 (d, J = 7.7, 3.1 Hz, 2H), 7.20 (dd, J = 8.5, 4.8 Hz, 1H), 7.01 (dd, J = 7.4, 4.9 Hz, 1H), 1.39 (s, 3H), 1.14 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ (ppm) 183.0, 161.3, 154.9, 149.7, 149.12, 149.07, 137.0, 136.3, 135.9, 134.0, 128.3, 127.3, 125.7, 123.6, 122.1, 121.7, 107.1, 55.04, 21.24, 18.95; MS (EI): 309 (55), 337 (100), 380 (45, M-1), 382 (35%, M+1). Anal. Calcd for C₂₃H₁₉N₅O: C 72.42, H 5.02, N 18.36; Found: C72.13, H 4.87, N 18.30.

6,6-Dimethyl-3,6*a***-di(2-pyridyl)-4-***p***-tolyl-6,6***a***-dihydropyrrolo-[3,2-***c*]**pyrazol-5-one 4b.** 77%, mp 119–120 °C; IR ν (cm⁻¹) 1755, 1638, 1583, 1514; ¹H NMR (400 MHz, CDCl₃) δ (ppm) 8.46 (d, J = 4.0 Hz, 1H), 8.17 (d, J = 4.2 Hz, 1H), 7.74 (d, J = 8.0 Hz, 1H), 7.66 (dt, J = 7.6, 1.8 Hz, 1H), 7.59 (d, J = 7.8 Hz, 1H), 7.55 (dd, J = 7.8, 1.7 Hz, 1H), 7.14–7.21 (m, 5H), 7.02 (dd, J = 6.8, 5.0 Hz, 1H), 2.41 (s, 3H), 1.38 (s, 3H), 1.13 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ (ppm) 183.1, 161.9, 154.9, 149.4, 149.0, 148.8, 137.4, 137.0, 136.3, 133.7, 133.3, 129.2, 129.0, 125.5, 123.6, 122.3, 121.8, 107.2, 55.1, 21.3, 21.2, 18.9; HRMS (ESI): 396.1823 (M+1), Anal. Calcd for C₂₄H₂₂N₃O: 396.1824 (M+1).

4-(4-Methoxyphenyl)-6,6-dimethyl-3,6*a***-di(2-pyridyl)-6,6***a***-dihydropyrrolo[3,2-***c***]pyrazol-5-one 4c. 70%, mp 162–163 °C; IR \nu (cm⁻¹) 1773, 1628, 1582, 1512; ¹H NMR (400 MHz, CDCl₃) \delta (ppm) 8.47 (d, J = 4.1 Hz, 1H), 8.19 (d, J = 4.6 Hz, 1H), 7.77 (d, J = 8.0 Hz, 1H), 7.66 (td, J = 7.8, 1.6 Hz, 1H), 7.58 (d, J = 7.8 Hz, 1H), 7.56 (td, J = 7.8, 1.6 Hz, 1H), 7.20–7.22 (m, 3H), 7.02 (dd, J = 7.4, 4.9 Hz, 1H), 6.90 (d, J = 8.9 Hz, 2H), 3.85 (s, 3H), 1.37 (s, 3H), 1.12 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) \delta (ppm) 183.4, 161.8, 158.8, 155.0, 149.8, 149.2, 149.0, 137.0, 135.9, 133.9, 129.5, 127.1, 123.6, 122.1, 121.7, 113.6, 107.0, 55.5, 54.9, 21.3, 18.9; MS (ESI): 412 (M+1). Anal. Calcd for C₂₄H₂₁N₅O₂: C 70.06, H 5.14, N 17.02; Found: C 69.87, H 5.51, N 16.95.**

4-(4-Chlorophenyl)-6,6-dimethyl-3,6*a***-di(2-pyridyl)-6,6***a***-dihydropyrrolo[3,2-***c***]pyrazol-5-one 4d. 82%, mp 112–113 °C; IR** *ν* **(cm⁻¹) 1762, 1635, 1583, 1493; ¹H NMR (400 MHz, CDCl₃)** *δ* **(ppm) 8.36 (d,** *J* **= 4.1 Hz, 1H), 8.02 (d,** *J* **= 4.5 Hz, 1H), 8.00 (d,** *J* **= 8.0 Hz, 1H), 7.58–7.61 (m, 2H), 7.54 (d,** *J* **= 7.9 Hz, 1H), 7.27 (d,** *J* **= 8.7 Hz, 2H), 7.16 (d,** *J* **= 8.7 Hz, 2H), 7.12 (dd,** *J* **= 6.3, 1.0 Hz, 1H), 6.96 (dd,** *J* **= 7.0, 5.0 Hz, 1H), 1.29 (s, 3H), 1.04 (s, 3H); ¹³C NMR (100 MHz, CDCl3)** *δ* **(ppm) 182.7, 160.7, 154.9, 149.6, 149.1, 149.1, 137.0, 136.1, 135.0, 134.2, 132.6, 128.2, 126.9, 123.8, 123.6, 122.0, 121.9, 107.2, 55.0, 21.2, 19.0; MS (CI): 416 (M+1); Anal. Calcd for C₂₃H₁₈ClN₅O: C 66.43, H 4.36, N 16.84; Found: C 66.53, H 4.79, N 16.75.**

4-(4-Bromophenyl)-6,6-dimethyl-3,6*a***-di(2-pyridyl)-6,6***a***-dihydropyrrolo[3,2-***c***]pyrazol-5-one 4e. 73%, mp 145–146 °C; IR ν (cm⁻¹) 1770, 1627, 1583, 1488; ¹H NMR (400 MHz, CDCl₃) δ (ppm) 8.43 (dd, J = 4.7, 0.8 Hz, 1H), 8.12 (d, J = 4.7 Hz, 1H), 8.06 (d, J = 8.0 Hz, 1H), 7.68 (td, J = 8.0, 1.8 Hz, 2H), 7.61 (d, J = 7.9 Hz, 1H), 7.49 (dt, J = 8.7, 2.8 Hz, 2H), 7.20–7.22 (m, 1H), 7.18 (dt, J = 8.7, 2.9 Hz, 2H), 7.27 (ddd, J = 7.5, 4.8, 1.0 Hz, 1H), 1.37 (s, 3H), 1.11 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ (ppm) 182.6, 160.7, 154.8, 149.4, 149.1, 148.9, 137.1, 136.4, 135.4, 134.0, 131.2, 127.2, 123.8, 123.7, 122.1, 122.0, 120.6, 107.2, 55.0, 21.20, 19.0; MS (EI): 44 (100), 417 (40), 459 (25%, M⁺); Anal. Calcd for C₂₃H₁₈BrN₅O: C 60.01, H 3.94, N 15.21; Found: C 59.89, H 3.95, N 15.18.**

6,6-Diethyl-3,6*a***-di(2-pyridyl)-4-***p***-tolyl-6,6***a***-dihydropyrrolo[3, 2-***c***]pyrazol-5-one 4f. 25%, mp 145–146 °C; IR ν (cm⁻¹) 1765, 1632, 1582; ¹H NMR (400 MHz, CD₃COCD₃) δ (ppm) 8.47 (dq,** *J* = 4.7, 1.0 Hz, 1H), 8.07 (dt, *J* = 8.0, 1.0, Hz, 1H), 7.94 (dq, *J* = 4.8, 1.0 Hz, 1H), 7.81 (dt, *J* = 8.0, 1.9, Hz, 1H), 7.73 (dt, *J* = 7.5, 1.8 Hz, 1H), 7.58 (dt, *J* = 8.0, 0.9 Hz, 1H), 7.35 (ddd, *J* = 7.6, 4.8, 1.1 Hz, 1H), 7.15 (d, *J* = 8.6 Hz, 2H), 7.03–7.06 (m, 3H), 2.35 (s, 3H), 1.95–2.02 (m, 1H), 1.77–1.85 (m, 1H), 1.38–1.47 (m, 1H), 1.11 (t, *J* = 7.6 Hz, 3H), 0.90 (t, *J* = 5.5 Hz, 3H), 0.87–0.93 (m, 1H); ¹³C NMR (100 MHz, CD₃COCD₃) δ (ppm) 180.9, 162.6, 155.2, 150.4, 149.0, 148.8, 137.5, 136.8, 136.1, 134.6, 133.3, 128.4, 126.0, 124.0, 123.2, 121.6, 121.4, 107.0, 63.0, 21.6, 20.3, 19.4, 7.8, 7.3; HRMS (TOF-ESI): 424.2125 (M+1); Anal. Calcd for C₂₆H₂₆N₅O: 424.2137 (M+1).

3',6*a*'-**Di**(2-pyridyl)-4'*-p*-tolyl-4'H-spiro[cyclohexane-1,6'-pyrrolo[3,2-*c*]pyrazol]-5'-one 4h. 88%, 156–157 °C; IR ν (cm⁻¹) 1752, 1639, 1581; ¹H NMR (400 MHz, CD₃COCD₃) δ (ppm) 8.47 (dq, J = 4.8, 1.0 Hz, 1H), 8.05 (dd, J = 8.0, 1.0, Hz, 1H), 7.95 (dq, J = 4.7, 1.0 Hz, 1H), 7.83 (dt, J = 7.9, 1.8, Hz, 1H), 7.72 (dt, J = 7.6, 1.8 Hz, 1H), 7.58 (dt, J = 8.0, 1.0 Hz, 1H), 7.35 (ddd, J = 7.6, 4.8, 1.1 Hz, 1H), 7.15 (d, J = 8.6 Hz, 2H), 7.08 (d, J = 8.6 Hz, 2H), 7.05 (ddd, J = 7.6, 4.8, 1.1 Hz, 1H), 1.34–1.55 (m, 5H); ¹³C NMR (100 MHz, CD₃COCD₃) δ (ppm) 181.5, 161.9, 155.2, 150.3, 149.1, 149.0, 137.4, 136.7, 136.1, 134.6, 133.2, 128.4, 126.0, 124.0, 123.7, 121.6, 121.5, 107.0, 60.4, 30.6, 25.5, 22.6, 22.03, 22.01, 20.4; HRMS (TOF-ESI): 436.2132 (M+1); Anal. Calcd for C₂₇H₂₆N₅O: 436.2137 (M+1).

General procedure for the reaction of β -lactam carbenes 2 with 3,6-di(2-pyridyl)tetrazine 3 in xylene

Under nitrogen atmosphere, the mixture of spiro[β -lactam-4,2'oxadiazolines] **1** (1.5 mmol) and 3,6-di(2-pyridyl)tetrazine **3** (1 mmol) was heated in refluxing xylene (50 mL) for 3–12 h. After removal of the solvent under vacuum, the residue was chromatographed on a silica gel column eluting with a mixture of petroleum ether and ethyl acetate from 10:1 to pure ethyl acetate to give products 5-triazolo[1,5-*a*]pyridylpyrrol-2-ones **5** in 75–92% yields. Products **5** were further purified by recrystallization in ethyl acetate and petroleum ether.

5-(3-[1,2,3]Triazolo[1,5-*a***]pyridyl)-3,3-dimethyl-1-phenyl-4-(2pyridyl)pyrol-2-one 5a.** 87%, mp 192–193 °C; IR v (cm⁻¹) 1716, 1622, 1582, 1562, 1518; ¹H NMR (400 MHz, CD₃COCD₃) δ (ppm) 8.86 (d, J = 7.0 Hz, 1H), 8.56 (dt, J = 3.9 Hz, 1H), 7.43 (d, J = 7.8 Hz, 1H), 7.34 (dt, J = 7.7, 1.9 Hz, 1H), 7.27 (dd, J = 9.0, 6.6 Hz, 1H), 7.19–7.23 (m, 2H), 7.10–7.16 (m, 4H), 7.07 (ddd, J = 7.5, 4.8 Hz, 1H), 6.82 (d, J = 8.1 Hz, 1H), 1.68 (s, 6H); ¹³C NMR (100 MHz, CDCl₃) δ (ppm) 182.1, 153.3, 149.0, 136.0, 135.0, 132.4, 130.8, 129.5, 128.7, 128.3, 127.2, 127.1, 126.4, 125.4, 124.1, 121.5, 117.4, 115.6, 49.48, 23.75; MS (ESI): 382(M+1); Anal. Calcd for C₂₃H₁₉N₅O: C 72.42, H 5.02, N 18.36; Found: C 72.04, H 4.68, N 18.17.

5-(3-[1,2,3]Triazolo[1,5-*a***]pyridyl)-3,3-dimethyl-4-(2-pyridyl)-1-**(*p*-tolyl)**pyrrol-2-one 5b.** 83%, mp 184–185 °C; IR v (cm⁻¹) 1733, 1619, 1584, 1513; ¹H NMR (400 MHz, CDCl₃) δ (ppm) 8.61 (d, J = 7.0 Hz, 1H), 8.56 (d, J = 4.2 Hz, 1H), 7.19 (d, J = 8.9 Hz, 1H), 7.07 (d, J = 6.8 Hz, 1H), 7.04 (d, J = 4.0 Hz, 2H), 7.00 (d, J = 4.1 Hz, 2H), 6.91 (t, J = 6.6 Hz, 1H), 6.85 (brs, 1H), 2.24 (s, 3H), 1.69 (s, 6H); ¹³C NMR (100 MHz, CDCl₃) δ (ppm) 182.2, 153.5, 149.1, 137.0, 135.8, 132.4, 132.4, 130.8, 129.3, 128.5, 127.0, 126.2, 125.4, 124.0, 121.4, 117.5, 115.5, 49.4, 23.7, 21.1, 21.0; MS (EI): 180 (100), 230 (100), 396 (3%, M+1); Anal. Calcd for $C_{24}H_{21}N_5O$: C 72.89, H 5.35, N 17.71; Found: C 72.68, H 5.30, N 17.82.

5-(3-[1,2,3]Triazolo[1,5-*a***]pyridyl)-1-(4-methoxyphenyl)-3,3-dimethyl-4-(2-pyridyl)pyrrol-2-one 5c.** 92%, mp 202–203 °C; IR ν (cm⁻¹) 1712, 1638, 1620, 1587, 1554, 1512; ¹H NMR (400 MHz, CDCl₃) δ (ppm) 8.61 (d, J = 7.0 Hz, 1H), 8.56 (d, J = 4.1 Hz, 1H), 7.26–7.24 (m, 1H), 7.19 (d, J = 9.0 Hz, 1H), 7.06–7.08 (m, 1H), 7.03 (dd, J = 8.9, 1.9 Hz, 2H), 6.99 (t, J = 6.8 Hz, 1H), 6.91 (td, J = 6.9, 1.0 Hz, 1H), 6.83 (d, J = 7.7 Hz, 1H), 6.73 (d, J = 8.9 Hz, 2H), 3.71 (s, 3H), 1.69 (s, 6H); ¹³C NMR (100 MHz, CDCl₃) δ (ppm) 182.5, 158.5, 153.5, 149.3, 135.6, 132.4, 130.8, 129.6, 128.5, 127.8, 126.3, 125.4, 123.8, 121.3, 117.4, 115.5, 114.0, 55.3, 49.3, 23.7; MS (ESI): 412 (M+1); Anal. Calcd for C₂₄H₂₁N₅O₂: C 70.06, H 5.14, N 17.02; Found: C 69.91, H 5.34, N 17.07.

5-(3-[1,2,3]Triazolo[1,5-*a***]pyridyl)-1-(4-chlorophenyl)-3,3-dimethyl-4-(2-pyridyl)pyrrol-2-one 5d.** 91%, mp 208–209 °C; IR *ν* (cm⁻¹) 1720, 1620, 1583, 1526; ¹H NMR (400 MHz, CDCl₃) *δ* (ppm) 8.63 (d, *J* = 7.1 Hz, 1H), 8.58 (d, *J* = 4.1 Hz, 1H), 7.27 (brs, 1H), 7.19 (d, *J* = 8.6 Hz, 2H), 7.16 (d, *J* = 9.0 Hz, 1H), 7.09 (d, *J* = 6.6 Hz, 1H), 7.05 (d, *J* = 8.7 Hz, 2H), 7.03 (brs, 1H), 6.94 (td, *J* = 6.9, 1.1 Hz, 1H), 6.84 (brs, 1H), 1.68 (s, 6H); ¹³C NMR (100 MHz, CDCl₃) *δ* (ppm) 182.0, 153.3, 149.4, 135.8, 133.7, 132.8, 132.3, 130.1, 129.9, 128.8, 128.4, 128.1, 126.6, 125.5, 124.0, 121.5, 117.3, 115.7, 49.5, 23.7; MS (CI): 416 (M+1); Anal. Calcd for C₂₃H₁₈ClN₅O: C 66.43, H 4.36, N 16.84; Found: C 66.52, H 4.60, N 16.84.

5-(3-[1,2,3]Triazolo[1,5-*a***]pyridyl)-1-(4-bromophenyl)-3,3-dimethyl-4-(2-pyridyl)pyrrol-2-one 5e.** 80%, mp 205–206 °C; IR *ν* (cm⁻¹) 1720, 1635, 1583, 1561; ¹H NMR (400 MHz, CDCl₃) δ (ppm) 8.56 (d, *J* = 7.0 Hz, 1H), 8.50 (d, *J* = 4.4 Hz, 1H), 7.27 (d, *J* = 8.6 Hz, 2H), 7.20 (bs, 1H), 7.08 (d, *J* = 8.9 Hz, 1H), 7.01(t, *J* = 6.6 Hz, 1H), 6.95 (brs 1H), 6.92 (d, *J* = 8.6 Hz, 2H), 6.87 (t, *J* = 6.9 Hz, 1H), 6.76 (brs 1H), 1.61 (s, 6H); ¹³C NMR (100 MHz, CDCl₃) δ (ppm) 181.9, 153.2, 149.3, 135.9, 134.2, 132.3, 131.8, 130.1, 129.8, 128.7, 128.0, 126.6, 125.6, 124.0, 121.6, 120.9, 117.3, 115.7, 49.5, 23.7; MS (ESI): 460 (M+1); Anal. Calcd for C₂₃H₁₈BrN₅O: C 60.01, H 3.94, N 15.21; Found: C 59.75, H 3.87, N 15.18.

5-(3-[1,2,3]Triazolo[1,5-*a***]pyridyl)-3,3-diethyl-4-(2-pyridyl)-1-(***p***-tolyl)pyrrol-2-one 5f. 82%; mp 182–183 °C, IR \nu (cm⁻¹) 1712, 1623, 1585, 1514; ¹H NMR (400 MHz, CDCl₃) \delta (ppm) 8.53 (d, J = 7.0 Hz, 1H), 8.48 (d, J = 4.3 Hz, 1H), 7.15 (td, J = 7.8, 1.8 Hz, 1H), 7.11 (d, J = 7.9 Hz, 1H), 7.00 (ddd, J = 8.8, 6.6, 0.8 Hz, 1H), 6.88–6.98 (m, 5H), 6.83 (td, J = 6.9, 1.2 Hz, 1H), 6.79 (d, J = 8.0 Hz, 1H), 2.21–2.26 (m, 2H), 2.16 (s, 3H), 1.92–1.97 (m, 2H), 0.91 (t, J = 7.3 Hz, 6H); ¹³C NMR (100 MHz, CDCl₃) \delta (ppm) 181.2, 153.8, 149.5, 137.1, 135.5, 133.0, 132.5, 132.4, 129.3, 128.7, 127.2, 126.3, 125.4, 125.1, 123.6, 121.1, 117.3, 115.4, 59.9, 30.1, 21.1, 9.3; MS (CI): 424 (M+1); Anal. Calcd for C₂₆H₂₅N₅O: C 73.74, H 5.95, N 16.54; Found: C 73.49, H 6.29, N 16.35.**

5-(3-[1,2,3]Triazolo[1,5-*a***]pyridyl)-3-ethyl-3-methyl-4-(2-pyridyl)-1-(***p***-tolyl)pyrrol-2-one 5g. 90%, mp 161–162 °C; IR \nu (cm⁻¹) 1713, 1635, 1619, 1582, 1514; ¹H NMR (400 MHz, CDCl₃) \delta (ppm) 8.53 (d, J = 7.0 Hz, 1H), 8.48 (dd, J = 4.8, 0.8 Hz, 1H), 7.16 (d, J = 7.6, 1H), 7.11 (d, J = 9.0 Hz, 1H), 7.00 (td, J = 7.4, 0.7 Hz, 1H), 6.89–6.94 (m, 5H), 6.83(td, J = 7.8, 1.1 Hz, 1H), 6.77**

(d, J = 7.9 Hz, 1H), 2.29–2.34 (m, 1H), 2.16 (s, 3H), 1.96–2.01 (m, 1H), 1.56 (s, 3H), 0.92 (t, J = 7.4 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ (ppm) 181.7, 153.6, 149.4, 137.0, 135.6, 132.4, 131.9, 129.3, 128.6, 127.3, 127.1, 126.3, 125.4, 125.4, 123.7, 121.2, 117.3, 115.5, 54.3, 30.6, 23.1, 21.0, 9.52; MS (CI): 410 (M+1); Anal. Calcd for C₂₅H₂₃N₅O: C 73.33, H 5.66, N 17.10; Found: C 73.45, H 5.44, N 17.16.

3'-(3-[1,2,3]Triazolo[1,5-*a***]pyridyl)-4'-(2-pyridyl)-2-(***p***-tolyl)-spiro[cyclohexane-1,3'-pyrrol]-5'-one 5h. 75%, mp 203–204 °C; IR v (cm⁻¹) 1704, 1624, 1584, 1512; ¹H NMR (400 MHz, CDCl₃) \delta (ppm) 8.79 (d,** *J* **= 4.1 Hz, 1H), 8.58 (dt,** *J* **= 4.8, 0.9 Hz, 1H), 7.37 (dt,** *J* **= 9.0, 1.0 Hz, 1H), 7.33 (td,** *J* **= 7.8, 1.9 Hz, 1H), 7.22 (ddd,** *J* **= 8.9, 6.8, 0.8 Hz, 1H), 7.04–7.09 (m, 2H), 7.01 (d,** *J* **= 8.7 Hz, 2H), 6.98 (d,** *J* **= 8.7 Hz, 2H), 6.82 (dd,** *J* **= 8.0, 0.9 Hz, 1H), 2.53 (td,** *J* **= 13.3, 4.2 Hz, 2H), 2.27 (tt,** *J* **= 12.9, 3.7 Hz, 2H), 2.20 (s, 3H), 1.90 (d,** *J* **= 13.2 Hz, 2H), 1.72–1.77 (m, 1H), 1.57 (dt,** *J* **= 13.2, 3.6 Hz, 2H), 1.32 (qt,** *J* **= 12.7, 3.7 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) \delta (ppm) 180.9, 154.2, 149.1, 136.7, 135.9, 132.7, 132.4, 131.1, 129.3, 128.6, 126.9, 126.0, 125.6, 125.3, 125.3, 121.5, 117.6, 115.4, 52.0, 31.4, 25.20, 21.1, 20.3; MS (CI): 436 (M+1); Anal. Calcd for C₂₇H₂₅N₅O: C 74.46, H 5.79, N 16.08; Found: C 74.79, H 6.06, N 16.08.**

General procedure for the reaction of β -lactam carbenes 2 with 3,6-di(4-pyridyl)tetrazine 3 in *n*-heptane

Under nitrogen atmosphere, the mixture of spiro[β -lactam-4,2'oxadiazolines] **1** (1.5 mmol) and 3,6-di(4-pyridyl)tetrazine **6** (1 mmol) was heated in refluxing *n*-heptane (50 mL) for 8 h. The yellow colored pyrido[*c*]cyclopenta[*b*]pyrrol-2-ones products **7** precipitated form the solvent during the reaction process. The crude products were then filtered and washed with chloroform, ethyl acetate, acetone and methanol to remove the soluble impurities. Products **7** can be further purified by recrystallization in DMSO and methanol.

3,3-Dimethyl-1-phenyl-8-(4-pyridyl)pyrido[c]cyclopenta[b]pyrrol-2-one 7a. 65%, mp 307–308 °C; IR ν (cm⁻¹) 3241, 1703, 1622, 1590; ¹H NMR (400 MHz, DMSO-d₆) δ (ppm) 11.97 (s, 1H), 8.22 (s, 1H), 8.04 (d, J = 5.9 Hz, 2H), 7.39 (d, J = 5.9 Hz, 2H), 7.26–7.28 (m, 4H), 7.16–7.18 (m, 2H), 6.76 (d, J = 6.0 Hz, 2H), 1.51 (s, 6H); ¹³C NMR (100 MHz, DMSO-d₆) δ (ppm) 184.2, 148.2, 144.0, 142.3, 136.4, 128.6, 127.0, 126.3, 125.8, 124.2, 123.3, 122.9, 121.0, 112.0, 107.2, 94.0, 42.2, 25.0; HRMS (TOF-ESI): 354.1611 (M+1), Anal. Calcd for C₂₃H₂₀N₃O: 354.1606 (M+1).

3,3-Dimethyl-8-(4-pyridyl)1-(*p***-tolyl)pyrido[***c***]cyclopenta**[*b***]pyrrol-2-one 7b.** 62%; mp 312–313 °C; IR ν (cm⁻¹) 3244, 1714, 1619, 1597; ¹H NMR (400 MHz, DMSO-d₆) δ (ppm) 11.94 (s, 1H), 8.21 (d, *J* = 4.4 Hz, 1H), 8.05 (d, *J* = 5.4 Hz, 2H), 7.39 (t, *J* = 4.1 Hz, 1H), 7.26 (d, *J* = 6.6 Hz, 1H), 7.07 (d, *J* = 8.2 Hz, 2H), 7.04 (d, *J* = 8.3 Hz, 2H), 6.77 (d, *J* = 5.4 Hz, 2H), 2.29 (s, 3H), 1.49 (s, 6H); ¹³C NMR (100 MHz, DMSO-d₆) δ (ppm) 184.3, 148.1, 144.2, 142.3, 136.4, 133.9, 129.0, 126.2, 125.7, 124.2, 123.3, 122.9, 121.0, 112.0, 107.2, 93.9, 42.1, 25.0, 20.6; MS (ESI): 368 (M+1). Anal. Calcd for C₂₄H₂₁N₃O: C 78.45, H 5.76, N 11.44; Found: C 78.13, H 5.56, N 11.19.

3,3-Dimethyl-1-(4-methoxyphenyl)-8-(4-pyridyl)pyrido[*c*]cyclopenta[*b*]pyrrol-2-one 7c. 65%; mp 302–303 °C; IR ν (cm⁻¹) 3244, 1698, 1621, 1592; ¹H NMR (400 MHz, DMSO-d₆) δ (ppm) 11.92 (s, 1H), 8.18 (s, 1H), 8.05 (d, J = 6.0 Hz, 2H), 7.37 (d, J = 6.7 Hz, 1H), 7.24 (d, J = 6.6 Hz, 1H), 7.06 (d, J = 8.8 Hz, 2H), 6.81 (d, J = 8.8 Hz, 2H), 6.77 (d, J = 6.0 Hz, 2H) 3.72 (s, 3H), 1.47 (s, 6H); ¹³C NMR (100 MHz, DMSO-d₆) δ (ppm) 184.5, 158.1, 148.1, 144.6, 142.2, 129.3, 127.8, 125.7, 124.1, 123.3, 122.9, 121.0, 113.9, 111.9, 107.1, 93.9, 55.4, 42.0, 25.0; MS (ESI): 384 (M+1). Anal. Calcd for C₂₄H₂₁N₃O₂: C 75.18, H 5.52, N 10.96; Found: C 74.98, H 5.91, N 10.72.

1-(4-Chlorophenyl)-3,3-dimethyl-8-(4-pyridyl)pyrido[*c*]cyclopenta[*δ*]pyrrol-2-one 7d. 60%, mp 318–319 °C; IR ν (cm⁻¹) 3246, 1716, 1620, 1598; ¹H NMR (400 MHz, DMSO-d₆) δ (ppm) 12.01 (s, 1H), 8.24 (d, *J* = 6.1 Hz, 1H), 8.11 (d, *J* = 6.0 Hz, 2H), 7.41 (t, *J* = 5.9 Hz, 1H), 7.32 (dd, *J* = 6.7, 2.0 Hz, 2H), 7.29 (d, *J* = 6.6 Hz, 1H), 7.20 (dd, *J* = 6.8, 2.0 Hz, 2H), 6.81 (dd, *J* = 6.1, 1.4 Hz, 2H), 1.50 (s, 6H); ¹³C NMR (100 MHz, DMSO-d₆) δ (ppm) 184.3, 148.2, 143.5, 142.3, 135.2, 131.1, 128.5, 127.9, 126.0, 124.1, 123.4, 123.0, 121.1, 111.9, 107.4, 93.9, 42.3, 25.0; HRMS (TOF-ESI): 388.1213, Anal. Calcd for C₂₃H₁₉CIN₃O: 388.1217 (M+1).

3,3-Diethyl-8-(4-pyridyl)1-(*p***-tolyl)pyrido**[*c*]cyclopenta[*b*]pyrrol-**2-one 7f.** 56%, mp 303–304 °C; IR ν (cm⁻¹) 1717, 1619, 1595; ¹H NMR (400 MHz, DMSO-d₆) δ (ppm) 11.97 (s, 1H), 8.19 (t, *J* = 3.3 Hz, 1H), 8.01 (d, *J* = 5.6 Hz, 2H), 7.36 (t, *J* = 5.7 Hz, 1H), 7.14 (d, *J* = 6.8 Hz, 1H), 7.05 (d, *J* = 8.2 Hz, 2H), 6.96 (d, *J* = 8.2 Hz, 2H), 6.76 (d, *J* = 6.0 Hz, 2H), 2.25 (s, 3H), 1.93–2.01 (m, 2H), 1.75–1.83 (m, 2H), 0.63 (t, *J* = 7.3 Hz, 6H); ¹³C NMR (100 MHz, DMSO-d₆) δ (ppm) 183.3, 148.1, 146.2, 142.2, 136.4, 133.9, 129.1, 126.0, 125.5, 124.6, 123.9, 123.2, 121.1, 107.6, 107.4, 93.8, 52.9, 30.2, 20.6, 9.3; HRMS (TOF-ESI): 396.2072 (M+1), Anal. Calcd for C₂₆H₂₆N₃O: 396.2076 (M+1).

3-Ethyl-3-methyl-8-(4-pyridyl)1-(*p***-tolyl)pyrido**[*c*]cyclopenta[*b*]**pyrrol-2-one 7g.** 51%, mp 298–300 °C; IR ν (cm⁻¹) 1718, 1619, 1596; ¹H NMR (400 MHz, DMSO-d₆) δ (ppm) 11.90 (s, 1H), 8.14 (d, *J* = 6.0 Hz, 1H), 7.98 (d, *J* = 6.0 Hz, 2H), 7.32 (t, *J* = 5.8 Hz, 1H), 7.14 (d, *J* = 6.6 Hz, 1H), 7.01 (d, *J* = 8.2 Hz, 2H), 6.95 (d, *J* = 8.2 Hz, 2H), 6.71 (dd, *J* = 4.5, 1.4 Hz, 2H), 2.22 (s, 3H), 1.86–1.95 (m, 1H), 1.73–1.81 (m, 1H), 1.40 (s, 3H), 0.63 (t, *J* = 7.4 Hz, 3H); ¹³C NMR (100 MHz, DMSO-d₆) δ (ppm) 183.9, 148.1, 145.2, 142.3, 136.4, 133.9, 129.1, 126.1, 125.6, 124.4, 123.3, 121.5, 121.1, 109.8, 107.3, 93.9, 47.2, 31.6, 23.3, 20.62, 20.57, 9.5; HRMS (TOF-ESI): 382.1928 (M+1), Anal. Calcd for C₂₅H₂₄N₃O: 382.1919 (M+1).

General procedure for the reaction of β -lactam carbenes 2 with 3,6-di(4-pyridyl)tetrazine 3 in 1,4-dioxane

Under nitrogen atmosphere, the mixture of spiro[β -lactam-4,2'oxadiazolines] **1** (2 mmol) and 3,6-di(4-pyridyl)tetrazine **6** (1 mmol) was heated in refluxing 1,4-dioxane (50 mL) for 11–18 h. After the reaction mixture was cooled to room temperature, the small amount of yellow solids were filtered and washed with ethyl acetate, acetone and methanol to give the byproducts **7** in 5–14% yield. The filtrates were combined to remove the solvents under vacuum. The residue was chromatographed on a silica gel column eluting with a mixture of petroleum ether, ethyl acetate and triethyl amine (10:5:1) to give products **8** in 45–79% yields. **3,3-Dimethyl-6-(3,3-dimethyl-4-oxo-1-phenylazetidin-2-yl)-1-phenyl-8-(4-pyridyl)pyrido**[*c*]cyclopenta[*b*]pyrrol-2-one **8a.** 57%; mp 289–290 °C; IR ν (cm⁻¹) 1772, 1725, 1627, 1596; ¹H NMR (400 MHz, DMSO-d₆) δ (ppm) 8.25 (br, 1H), 7.97 (s, 2H), 7.43 (d, *J* = 6.6 Hz, 1H), 7.32 (d, *J* = 8.2 Hz, 2H), 7.26 (d, *J* = 7.5 Hz, 2H), 7.18–7.22 (m, 4H), 7.06–7.11 (m, 3H), 6.66 (br, 2H), 6.34 (s, 1H), 1.44 (s, 6H), 1.41 (s, 3H), 0.89 (s, 3H); ¹³C NMR (100 MHz, DMSO-d₆) δ (ppm) 184.0, 169.3, 148.2, 148.1, 145.4, 141.6, 136.1, 136.0, 129.5, 128.6, 127.2, 126.5, 125.1, 124.7, 123.5, 121.8, 116.9, 114.0, 107.6, 95.0, 77.7, 57.3, 42.1, 24.8, 21.2, 15.1; HRMS (TOF-ESI): 527.2458 (M+1), Anal. Calcd for C₁₄H₁₁N₄O₂: 527.2447 (M+1).

6-(3,3-Dimethyl-4-oxo-1-(*p***-tolyl)azetidin-2-yl)-3,3-dimethyl-8-(4-pyridyl)1-(***p***-tolyl)pyrido**[*c***]cyclopenta**[*b***]pyrrol-2-one 8b.** 70%; mp 180–181 °C; IR *v* (cm⁻¹) 1774, 1725, 1626, 1592; ¹H NMR (400 MHz, DMSO-d₆) δ (ppm) 8.28 (br, 1H), 8.03 (s, 2H), 7.44 (d, *J* = 5.8 Hz, 1H), 7.16–7.26 (m, 5H), 7.07 (d, *J* = 8.1 Hz, 2H), 7.00 (d, *J* = 8.1 Hz, 2H)), 6.72 (br, 2H), 6.36 (s, 1H), 2.28 (s, 3H), 2.24 (s, 3H), 1.48 (s, 6H), 1.45 (s, 3H), 0.93 (s, 3H); ¹³C NMR (100 MHz, DMSO-d₆) δ (ppm) 184.0, 169.1, 148.1, 145.6, 141.6, 136.7, 134.0, 133.7, 133.6, 129.8, 129.1, 126.4, 125.1, 123.6, 121.8, 116.9, 114.0, 107.5, 94.9, 77.7, 57.2, 42.0, 24.8, 21.2, 20.6, 20.4, 15.2; HRMS (TOF-ESI): 555.2764 (M+1), Anal. Calcd for C₃₆H₃₅N₄O₂: 555.2760 (M+1).

3,3-Dimethyl-6-[1-(4-methoxyphenyl)-3,3-dimethyl-4-oxo-azetidin-2-yl]-1-(4-methoxyphenyl)-8-(4-pyridyl)pyrido[c]cyclopenta[*b***]-pyrrol-2-one 8c.** 78%; mp 240–241 °C; IR ν (cm⁻¹) 1763, 1721, 1626, 1591; ¹H NMR (400 MHz, DMSO-d₆) δ (ppm) 8.25 (br, 1H), 8.05 (d, *J* = 3.5 Hz, 2H), 7.44 (d, *J* = 7.0 Hz, 1H), 7.26 (d, *J* = 9.0 Hz, 2H), 7.26 (br, 1H), 7.05 (d, *J* = 8.8 Hz, 2H), 6.95 (d, *J* = 8.9 Hz, 2H), 6.81 (d, *J* = 8.9 Hz, 2H), 6.72 (br, 2H), 6.35 (s, 1H), 3.73 (s, 3H), 3.69 (s, 3H), 1.48 (s, 6H), 1.46 (s, 3H), 0.93 (s, 3H); ¹³C NMR (100 MHz, DMSO-d₆) δ (ppm) 184.2, 168.8, 158.4, 156.3, 148.2, 146.0, 141.6, 129.3, 128.9, 127.9, 125.1, 123.6, 121.8, 118.4, 114.7, 113.9, 107.4, 94.8, 77.9, 57.2, 55.4, 55.3, 41.9, 24.8, 21.2, 15.2; HRMS (TOF-ESI): 587.2662 (M+1), Anal. Calcd for C₃₆H₃₅N₄O₄: 587.2658 (M+1).

1-(4-Chlorophenyl)-6-[1-(4-chlorophenyl)-3,3-dimethyl-4-oxoazetidin-2-yl]-3,3-dimethyl-8-(4-pyridyl)pyrido[c]cyclopenta[*b***]py-rrol-2-one 8d.** 45%; mp 180–181 °C; IR ν (cm⁻¹) 1778, 1724, 1626, 1592; ¹H NMR (400 MHz, DMSO-d₆) δ (ppm) 8.30 (br, 1H), 8.06 (s, 1H), 7.45 (d, J = 6.9 Hz, 1H), 7.39 (d, J = 8.7 Hz, 2H), 7.27 (d, J = 9.1 Hz, 2H), 7.23 (d, J = 7.8 Hz, 2H), 7.21 (br, 1H), 7.11 (d, J = 8.6 Hz, 2H), 6.75 (br, 2H), 6.34 (s, 1H), 1.43 (s, 6H), 1.41 (s, 3H), 0.89 (s, 3H); ¹³C NMR (100 MHz, DMSO-d₆) δ (ppm) 183.9, 169.3, 148.3, 144.9, 141.6, 135.0, 134.9, 131.5, 129.4, 128.6, 128.5, 128.1, 125.1, 123.7, 121.8, 118.6, 114.0, 107.7, 95.0, 77.9, 57.6, 42.2, 24.8, 21.2, 15.1; HRMS (TOF-ESI): 595.1647 (M+1), Anal. Calcd for C₃₄H₂₉Cl₂N₄O₂: 595.1668 (M+1).

6-[3,3-Diethyl-1-(4-methoxyphenyl)-4-oxo-azetidin-2-yl]-3,3-diethyl-1-(4-methoxyphenyl)-8-(4-pyridyl)pyrido[c]cyclopenta[b]pyrrol-2-one 8i. 79%; mp 168–169 °C; IR ν (cm⁻¹) 1765, 1722, 1625, 1591; ¹H NMR (400 MHz, DMSO-d₆) *δ* (ppm) 8.32 (br, 1H), 8.06 (s, 2H), 7.47 (d, *J* = 6.9 Hz, 1H), 7.28 (d, *J* = 8.2 Hz, 2H), 7.15 (s, 1H), 6.99 (d, *J* = 8.6 Hz, 2H), 6.96 (d, *J* = 8.6 Hz, 2H), 6.81 (d, *J* = 8.9 Hz, 2H), 6.73 (br, 1H), 6.45 (s, 1H), 3.72 (s, 3H), 3.69 (s, 3H), 1.91–2.01 (m, 3 H), 1.77–1.85 (m, 3 H), 1.53 (br, 1H), 1.34 (br, 1H), 1.02 (t, *J* = 7.2 Hz, 3H), 0.68–0.66 (m, 9H); ¹³C NMR (100 MHz, DMSO-d₆) *δ* (ppm) 183.1, 168.2, 158.4, 156.4, 148.2, 147.8, 141.5, 129.0, 127.7, 125.6, 123.5, 122.8, 118.6, 114.7, 114.0, 109.8, 107.6, 94.7, 76.0, 64.4, 55.3, 55.2, 52.6, 30.0, 23.3, 19.6, 9.4, 8.3, 7.8; HRMS (TOF-ESI): 643.3291 (M+1), Anal. Calcd for $C_{40}H_{41}N_4O_4$: 643.3284 (M+1).

6'-[1''-(4-Methoxyphenyl)-spiro[cyclopentane-1,3''-(4''-oxo-aze-tidin-2''-yl)]-1'-(4-methoxyphenyl)-8'-(4-pyridyl)spiro[cyclopenta-ne-1,3'-pyrido]c]cyclopenta[b]pyrrol]-2'-one 8j. 65%; mp 192–193 °C; IR ν (cm⁻¹) 1766, 1721, 1625, 1591; ¹H NMR (400 MHz, DMSO-d₆) δ (ppm) 8.38 (br, 1H), 8.08 (s, 2H), 7.40 (s, 1H), 7.22 (d, J = 9.0 Hz, 2H), 7.10 (d, J = 6.2 Hz, 1H), 7.04 (d, J = 8.7 Hz, 2H), 6.93 (d, J = 8.8 Hz, 2H), 6.80 (d, J = 8.8 Hz, 2H), 6.73 (br, 2H), 6.47 (s, 1H), 3.72 (s, 3H), 3.68 (s, 3H), 2.25–2.31 (m, 1H), 1.92–2.10 (m, 10H), 1.41–1.73 (m, 4H), 1.04–1.09 (m, 1H); ¹³C NMR (100 MHz, DMSO-d₆) δ (ppm) 185.1, 169.6, 158.8, 156.7, 148.8, 147.2, 142.1, 129.8, 129.4, 128.4, 126.0, 124.2, 121.5, 118.8, 115.2, 115.0, 114.4, 108.2, 95.4, 78.9, 67.4, 55.9, 55.8, 52.4, 38.59, 38.56, 33.5, 27.9, 26.7, 26.6; HRMS (TOF-ESI): 639.2964 (M+1), Anal. Calcd for C₄₀H₃₉N₄O₄: 639.2971 (M+1).

Acknowledgements

This work was supported by the National Natural Science Foundation of China (No. 20832006 and 20972017), Beijing Municipal Commission of Education and the Fundamental Research Funds for the Central Universities (2009SC-1).

Notes and references

- 1 M. Zoghbi and J. Warkentin, J. Org. Chem., 1991, 56, 3214-3215.
- 2 M. Zoghbi and J. Warkentin, Can. J. Chem., 1993, 71, 912-918.
- 3 M. Zoghbi and J. Warkentin, Can. J. Chem., 1992, 70, 2967-2971.
- 4 M. Zoghbi, S. E. Horne and J. Warkentin, J. Org. Chem., 1994, 59, 4090–4095.
- 5 Y. Cheng, B. Wang and L.-Q. Cheng, J. Org. Chem., 2006, 71, 4418-4427.
- 6 Y. Cheng and L.-Q. Cheng, J. Org. Chem., 2007, 72, 2625-2630.
- 7 L.-Q. Cheng and Y. Cheng, Tetrahedron, 2007, 63, 9359-9364.
- 8 J. Xing, X.-R. Wang, C.-X. Yan and Y. Cheng, J. Org. Chem., 2011, 76, 4746–4752.
- 9 N. Shao, G.-X. Pang, C.-X. Yan, G.-F. Shi and Y. Cheng, J. Org. Chem., 2011, 76, 7458–7465.
- 10 A. Kümmell and G. Seitz, Tetrahedron Lett., 1991, 32, 2743-2746.
- 11 G. Frenzen, A. Kümmell, C. Meyer-Dulheuer and G. Seitz, Chem. Ber.,
- 1994, **127**, 1803–1806. 12 C. Gerninghaus, A. Kümmell and G. Seitz, *Chem. Ber.*, 1993, **126**, 733–738.
- 13 P. Imming, A. Kümmell and G. Seitz, *Heterocycles*, 1993, 35, 299– 304.
- 14 Y. F. Suen, H. Hope and M. H. Nantz, J. Org. Chem., 2005, 70, 8468-8471.