

Tetrahedron Letters 40 (1999) 5187-5190

TETRAHEDRON LETTERS

## Asymmetric synthesis of amino sugars. Part 1: Stereoselective synthesis of (2S,3S,4R,5S)-2-amino-1,3,4,5,6-hexanepentol derivatives and their conversion to L-mannosamine derivatives

Ludmila Ermolenko, N. André Sasaki \* and Pierre Potier

Institut de Chimie des Substances Naturelles, CNRS, Avenue de la Terrasse, 91198 Gif-sur-Yvette, France

Received 13 April 1999; accepted 13 May 1999

## Abstract

A novel approach for asymmetric synthesis of amino sugars is developed, starting from readily available chiral building blocks 1 and 2,3-O-isopropylideneglyceraldehyde 2, via the Julia olefination and subsequent dihydroxylation as key steps.  $\bigcirc$  1999 Published by Elsevier Science Ltd. All rights reserved.

Keywords: amino sugar; L-mannosamine; Julia olefination; dihydroxylation.

Amino sugars are constituents of various biologically active compounds such as anticancer antibiotics<sup>1</sup> and biopolymers<sup>2</sup> and consequently these molecules have been the targets of syntheses for many years. The traditional methods have involved carbohydrates as principal intermediates and usually required multistep synthetic transformations. Recently, several groups have focused on the asymmetric synthesis of amino sugars from non-carbohydrate precursors.<sup>3,4</sup> Among the methods available, the common approaches are direct amination of carbocycles,<sup>5a-c</sup> transformation of isoxazolines<sup>5d-f</sup> and 2-thiazolyl *N*-alkylhydroxylamines,<sup>5g,h</sup> the use of hetero Diels–Alder reactions,<sup>5i,j</sup> [3+2] cycloaddition of nitrones with vinylene carbonate<sup>5k</sup> and the utilization of readily available natural products such as amino acids<sup>51,m</sup> and lactic acid<sup>5n-p</sup> as chiral pool materials for the construction of amino sugar skeletons. One of the most interesting methodologies is the transformation of serine and threonine into various types of amino sugars.<sup>5q</sup> However, some of these reactions suffer from complete lack of enantioselectivity, lengthy reaction sequences and limited versatility.

We describe here a novel and potentially versatile methodology for asymmetric synthesis of amino sugars either in the D- or L-configuration based on the transformation of the readily available chiral building blocks  $1^6$  and 2 into a fully protected derivative of 2-amino polyol 3 (Fig. 1).

Starting from either (R)- or (S)-1 and either (2R)- or (2S)-2,3-O-isopropylideneglyceraldehyde 2 and using the Julia olefination process followed by dihydroxylation, one can obtain a priori any one

<sup>\*</sup> Corresponding author. E-mail: andre.sasaki@icsn.cnrs-gif.fr



of four diastereomers of the 2-amino hexanepentol derivative 3 as a major compound with the four asymmetric centers of desired stereochemistry at C-2, C-3, C-4 and C-5. This depends on three factors: the absolute configuration of 1 and 2, the stereochemistry of the newly created olefin bond (E or Z) and the course of stereoselection during dihydroxylation. Intramolecular cyclization between C-1 and the OH function at C-5 of 2-aminopolyol 3 provides four 2-amino sugars in pyranose-form: L- or D-mannosamine and L- or D-glucosamine. Likewise, the cyclization between C-1 and the OH function at C-4 affords the furanose-form of these amino sugars. In order to demonstrate our novel approach in this preliminary communication, we have chosen L-mannosamine as a target molecule which is difficult to obtain in enantio- and diastereomerically pure form.<sup>7</sup> The synthesis of 1,4,6-tri-O-acetyl-2-N-(tert-butoxycarbonyl)amino-2-deoxy-3-O-tert-butyldimethylsilyl-L-mannopyranose 12 is shown in Scheme 1.



Scheme 1. Reagents and conditions: (a) NaH, THF, 0°C, 30 min, BnBr, Bu<sub>4</sub>NI, 0°C, 24 h, 86%; (b) 2 equiv. BuLi, THF, (2S)-2,3-O-isopropylideneglyceraldehyde 2, -78°C, 2 h, then rt, 1 h, 85%; (c) 6% Na-Hg, Na<sub>2</sub>HPO<sub>4</sub>, MeOH, 0°C, 3 h, 80%; (d) OsO<sub>4</sub>, NMO, 'BuOH:H<sub>2</sub>O (1:4), rt, 2 h, 95%; (e) TBDMSCl, imidazole, DMF, rt, 24 h, 95%; (f) Ac<sub>2</sub>O, Et<sub>3</sub>N, DMAP, 50°C, 8 h, 80%; (g) H<sub>2</sub> (1 atm), 10% Pd/C, EtOAc, rt, 6 h, 98%; (h) Py·SO<sub>3</sub>, Et<sub>3</sub>N, DMSO, 0°C, 2 h, 90%; (i) HCl, MeOH, rt, 8 h, then Ac<sub>2</sub>O, Et<sub>3</sub>N, DMAP, CH<sub>2</sub>Cl<sub>2</sub>, rt, 3 h, 80%

Coupling of the dilithiate of 4, prepared from (S)-1,<sup>8</sup> with (2S)-2,3-O-isopropylideneglyceraldehyde  $2^{9,10}$  afforded a diastereomeric mixture of  $\beta$ -hydroxysulfone 5 in good yield.<sup>11</sup> Treatment of the  $\beta$ -hydroxysulfone 5 with 6% Na–Hg furnished the olefin 6 in 80% yield as a mixture of *E*- and *Z*-isomers (4:1) which were easily separated by column chromatography. Dihydroxylation of the *E*-olefin 6 was first investigated using the Sharpless chiral reagent AD-mix- $\beta$ .<sup>12</sup> After 48 h, 40% conversion of the olefin to a mixture of diols 7 and 13 was attained with a 9:1 ratio, respectively.<sup>13</sup> On the contrary, application of the osmium-catalyzed procedure without any chiral auxiliary<sup>14</sup> afforded a mixture of diols 7 and 13

in 95% yield with a 77:23 ratio, respectively. These diols are easily separable by preparative column chromatography.

The stereochemistry of diols 7 and 13 was assigned by NMR studies of N,O-bisacetyl oxazolidinone 14 and 15<sup>15</sup> obtained from 7 and 13, respectively, by treatment with NaH followed by acetylation. The  $J_{2,3}$  coupling constants are 7.0 Hz for the 2,3-*anti*-isomer 14 and 3.1 Hz for the 2,3-*syn*-isomer 15. These values are in agreement with corresponding data for related compounds<sup>5g,16</sup> and are also consistent with the dihedral angles determined from molecular mechanics calculations (approximately 10° for 14 and 115° for 15). Furthermore, an intense NOE observed between H-2 and H-3, and between H-1 and H-4 in 2D NOESY experiments for oxazolidinone 14 is in accordance with the stereochemistry depicted in Scheme 2.



Scheme 2. Reagents and conditions: (a) NaH, THF, 50°C, 2 h, 80%; (b) Ac<sub>2</sub>O, Et<sub>3</sub>N, DMAP, CH<sub>2</sub>Cl<sub>2</sub>, 95%

The protection of the hydroxy functionalities of the diol 7 was found to be more difficult than expected. Attempts to obtain 3,4-O-bis protected diols were successful only in the case of acetyl and isopropylidene protecting groups. The presence of an acetyl group at the C-3 position of diol 7 resulted in undesired  $\alpha$ ,  $\beta$ -elimination during the oxidation of the C-1 primary alcohol to an aldehyde (step h). Fortunately, 3-O-mono TBDMS ether 8 was obtained in 95% yield by employing conventional conditions (TBDMSCl, imidazole, DMF, rt, 24 h). Subsequent acetylation at the C-4 position of 8 furnished the fully protected 2-amino hexanepentol 9 which was subjected to hydrogenolysis to give  $\alpha$ -amino alcohol 10. Oxidation of this amino alcohol by Py·SO<sub>3</sub> in DMSO<sup>17</sup> provided the desired  $\alpha$ -amino aldehyde 11 in 90% yield. Hydrolysis of acetonide 11 in MeOH in the presence of a catalytic amount of HCl followed by acetylation afforded the fully protected L-mannosamine 12 as a mixture of  $\alpha$ - and  $\beta$ -anomers approximately in 2:1 ratio, respectively, and in 80% yield.<sup>18</sup>

In summary, we have demonstrated an efficient methodology for the construction of amino sugars with good stereochemical control as exemplified by the synthesis of an enantiopure L-mannosamine derivative. Further applications of this new approach to the synthesis of other amino sugars of both Land D-configurations are in progress.

## Acknowledgements

We thank Dr. Gabor Lukacs for his helpful suggestions in assigning the  ${}^{1}$ H and  ${}^{13}$ C chemical shifts of 12.

## References

- 1. Pelyvás, I. V.; Monneret, C.; Herczegh, P. In Synthetic Aspects of Aminodeoxy Sugars of Antibiotics; Springer: Berlin, 1988; pp. 123-162.
- 2. Lindberg, B. Adv. Carbohydr. Chem. Biochem. 1990, 48, 279-318.
- 3. Hudlicky, T.; Entwistle, D. A.; Pitzer, K. K.; Thorpe, A. J. Chem. Rev. 1996, 96, 1195-1220.
- 4. Otsomaa, L. A.; Koskinen, A. M. P. In Progress in the Chemistry of Organic Natural Products; Herz, W.; Kirby, G. W.; Moore, R. E.; Steglich, W.; Tamm, C., Eds.; Springer: Wien, 1998; pp. 197-255.
- For some selected recent articles see: (a) Lehmann, J.; Moritz, A. Liebigs Ann. Chem. 1991, 937-940; (b) Pitzer, K.; Hudlicky, T. Synlett 1995, 803-805; (c) Bérubé, G.; Luce, E.; Jankowsky, K. Bull. Soc. Chim. Fr. 1983, II, 109-111; (d) Jäger, V.; Schohe, R. Tetrahedron 1984, 40, 2199-2210; (e) Wade, P. A.; Price, D. T. Tetrahedron Lett. 1989, 30, 1185-1189; (f) Jäger, V.; Müller, I.; Schohe, R.; Frey, M.; Ehrler, R.; Häfele, B.; Schröter, D. Lect. Heterocycl. Chem. 1985, 8, 79-98 and references cited therein; (g) Dondoni, A.; Fantin, G.; Fogagnolo, M.; Pedrini, P. J. Org. Chem. 1990, 55, 1439-1446; (h) Dondoni, A.; Franco, S.; Junquera, F.; Merchán, F. L.; Merino, P.; Tejero, T.; Bertolasi, V. Chem. Eur. J. 1995, I, 505-520 and references cited therein; (i) Tietze, L. Q.; Hartfield, U. Tetrahedron Lett. 1990, 31, 1697-1700; (j) Vogel, P. Bull. Soc. Chim. Belg. 1990, 99, 395-439; (k) Xu, Z.; Johannes, C. W.; Houri, A. F.; La, D. S.; Cogan, D. A.; Hofilena, G. E.; Hoveyda, A. H. J. Am. Chem. Soc. 1997, 119, 10302-10316; (l) Jurczak, J.; Golebiowski, A. Chem. Rev. 1989, 89, 149-164; (m) Coppola, G. M.; Schuster, H. F. Asymmetric Synthesis: Construction of Chiral Molecules Using Amino Acids; Wiley: New York, 1987; (n) Hamada, Y.; Kawai, A.; Shioiri, T. Tetrahedron Lett. 1984, 25, 5413-5414; (o) Hamada, Y.; Kawai, A.; Matsui, T.; Hara, O.; Shioiri, T. Tetrahedron 1990, 46, 4823-4846; (p) Hamada, Y.; Kawai, A.; Shioiri, T. Tetrahedron Lett. 1984, 25, 5409-5412; (q) Golebiowski, A.; Jurczak, J. Synlett 1993, 241-245.
- 6. Sasaki, N. A.; Hashimoto, C.; Potier, P. Tetrahedron Lett. 1987, 28, 6069-6072. Both (R)- and (S)-1 are commercially available in THP protected form from Acros Organics, Belgium.
- 7. O'Neill, A. N. Can. J. Chem. 1959, 37, 1747-1753.
- 8. Dockner, M.; Sasaki, N. A.; Rich, C.; Potier, P. Liebigs Ann/Recueil 1997, 1267-1272.
- 9. Andrews, G. C.; Crawford, T. C.; Bacon, B. E. J. Org. Chem. 1981, 46, 2976-2977.
- 10. Hubschwerlen, C. Synthesis 1986, 962-964.
- 11. Julia, M.; Paris, J.-M. Tetrahedron Lett. 1973, 4833-4836.
- Sharpless, K. B.; Amberg, W.; Bennani, Y. L.; Crispino, G. A.; Hartung, J.; Jeon, K.-S.; Kwong, H.-L.; Morikawa, K.; Wang, Z.-M.; Xu, D.; Zhang, X.-L. J. Org. Chem. 1992, 57, 2768–2771.
- 13. Determined by HPLC on a Hypersil silica column (5  $\mu$ ), (heptane:propanol-2, 90:10).
- 14. VanRheenen, V.; Kelly, R. C.; Cha, D. Y. Tetrahedron Lett. 1976, 1973-1976.
- The signals due to H-2 and H-3 in the <sup>1</sup>H NMR (CDCl<sub>3</sub>) spectra of 14 and 15 are as follows. Compound 14: δ 4.69 (dd, J<sub>3,4</sub>=J<sub>2,3</sub>=7.0 Hz, 1H, H-3), 4.59 (ddd, J=7.0, 4.8, 2.0 Hz, 1H, H-2). Compound 15: δ 4.66 (dd, J<sub>3,4</sub>=4.7 Hz, J<sub>2,3</sub>=3.1 Hz, 1H, H-3), 4.45-4.55 (m, 1H, H-2).
- (a) Futagawa, S.; Inui, T.; Shiba, T. Bull. Chem. Soc. Jpn. 1973, 46, 3308-3310; (b) Rama Rao, A. V.; Murali Dhar, T. G.; Chakraborty, T. K.; Gurjar, M. K. Tetrahedron Lett. 1988, 29, 2069-2072; (c) Kano, S.; Yokomatsu, T.; Shibuya, S. J. Org. Chem. 1989, 54, 513-515.
- 17. Clayden, J.; McCarthy, C.; Cumming, J. G. Tetrahedron: Asymmetry 1998, 9, 1427-1440.
- 18. Compound 12: IR (neat) 3449, 3031, 2959, 1748, 1714, 1504, 1392, 1369, 1233, 1165, 1140, 1059, 967, 840 cm<sup>-1</sup>; <sup>1</sup>H NMR (250 MHz, CHCl<sub>3</sub>) δ 6.15 (br. s, α-anomer H-1) and 5.85 (d,  $J_{1,2}$ =2.4 Hz, β-anomer H-1), 1H; 5.05–4.8 (m, α-H-4 and β-H-4, α-NH and β-NH), 2H; 4.45–4.30 (m), 4.28–4.10 (m), 4.04 (dd, J=12.3, 2.5 Hz), 3.95–3.85 (m), 3.82–3.75 (m), 5H; 2.14 (s), 2.10 (s), 2.09 (s), 2.08 (s), 9H; 1.44 (s), 9H; 0.89 (s), 0.86 (s), 9H; 0.12 (s), 0.10 (s), 0.09 (s), 6H; <sup>13</sup>C NMR (62.9 MHz, CHCl<sub>3</sub>) δ 170.8, 170.6, 169.8, 169.6, 169.2, 168.4 (CH<sub>3</sub>CO); 155.7 (Me<sub>3</sub>COCO); 92.3 (C-1, α-anomer), 91.6 (C-1, β-anomer), 80.2 (Me<sub>3</sub>COCO, α), 80.0 (Me<sub>3</sub>COCO, β), 72.9 (C-3-β or C-5-β interchangeable attribution), 70.1 (C-3-α or C-5-α), 69.3 (C-3-β or C-5-β), 68.8 (C-3-α or C-5-α), 68.4 (C-4-β), 68.1 (C-4-α), 62.9 (C-6-β), 62.3 (C-6-α), 53.9 (C-2-β), 51.1 (C-2-α), 28.3 (Me<sub>3</sub>COCO), 25.0 (Me<sub>3</sub>CSiMe<sub>2</sub>), 21.0 and 20.8 (CH<sub>3</sub>CO), 17.9 (Me<sub>3</sub>CSiMe<sub>2</sub>), -4.8 (Me<sub>3</sub>CSiMe<sub>2</sub>, β), -5.0 (Me<sub>3</sub>CSiMe<sub>2</sub>, α); MS (CI) m/z 537 [M+NH<sub>4</sub>], 436.