

# Total synthesis of (±)-dihydrospininiferin-1 via a polyfluoro alkanosulfonyl fluoride induced tandem carbonium ion rearrangement reaction

Ling Chen, Kai Ding and Wei-Sheng Tian\*

Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 354 Fenglin Road, Shanghai, 200032, China. E-mail: wstian@pub.sioc.ac.cn; Fax: 86-21-64166128; Tel: 086-021-64163300-83225

Received (in Corvallis, OR, USA) 2nd January 2003, Accepted 18th February 2003

First published as an Advance Article on the web 28th February 2003

A novel polyfluoroalkanosulfonyl fluoride induced carbonium ion rearrangement reaction of  $\gamma$ -hydroxymethyl cyclohexenone has been used for the total synthesis of (±)-dihydrospininiferin 1.


Spiniferin-1 (**1**) is a structurally unique furanosesquiterpene, isolated by Cimino *et al.* in 1975 from the Mediterranean sponge *Pleraplysilla spinifera*, which is present in the Bay of Naples.<sup>1</sup> In 1978, its structure was reformulated as **12** and supported by Marshall's rational synthesis of (±)-dihydrospininiferin-1 (**2**) (Fig. 1).<sup>3</sup> Racemic **1** had also been synthesized<sup>4</sup> by Marshall by a similar strategy. Spiniferin-1 is the first and unique natural product with the skeleton of 1,6-methano[10]annulene up to now, although Vogel *et al.* prepared this kind of annulene in 1964.<sup>5</sup> The physical and chemical properties of 1,6-methano[10]annulene have been described in detail since it and its analogues were synthesized, but to the best of our knowledge no biological information has been reported. The biological secrets of spiniferin-1 make us interested in its total synthesis.

As a part of our research on the reaction of poly(per)-fluoroalkanosulfonyl fluorides and their application in the synthesis of medicines and natural products,<sup>6</sup> we found a novel polyfluoroalkanosulfonyl fluoride induced carbonium ion rearrangement reaction<sup>7</sup> of  $\gamma$ -hydroxymethyl cyclohexenone recently. This rearrangement reaction provides a new and efficient synthetic method for dihydrospininiferin-1, spiniferin-1, as well as their analogues. In this paper, we will describe the total synthesis of (±)-dihydrospininiferin-1 via a novel polyfluoroalkanosulfonyl fluoride induced carbonium ion rearrangement reaction of  $\gamma$ -hydroxymethyl cyclohexenone.


Our synthetic plan relied on the construction of 1,6-methano[10]annulene **3**, a key intermediate in Marshall's synthesis,<sup>3</sup> from polyhydro-4a,8a-methanonaphthalene **4**, which would be obtained by the rearrangement of  $\gamma$ -hydroxymethyl cyclohexenone **5** induced by polyfluoroalkanosulfonyl fluoride. Unsaturated ketone **5** was simplified into a readily available Robinson annulation product **6** and an  $\alpha$ -halogen acetaldehyde

afford **10** in 80% yield upon heating with methanolic potassium carbonate<sup>4</sup> (Scheme 2).

We planned to use  $\text{BrCH}_2\text{CH}(\text{OEt})_2$ , the equivalent of  $\alpha$ -halogen acetaldehyde, to introduce the side chain. Kinetic controlled enolation of enone **10** could be realized by using  $\text{LDA}^{10}$  as the base at low temperature ( $-70^\circ\text{C}$ ). But the subsequent alkylation did not occur even with the iodide due to the poor reactivity of enol anion. Fortunately, the alkylation of allylic bromide instead of  $\text{XCH}_2\text{CH}(\text{OEt})_2$  ( $\text{X} = \text{Br}$  or  $\text{I}$ ) gave two isomers **11a** and **11b** in 63% and 9% yields respectively. Treatment of **11a** with  $\text{LiAlH}_4$  followed by selective oxidation of the allylic hydroxyl with DDQ<sup>11</sup> afforded hydroxymethyl enone **13**. Ozonolysis of **13** directly furnished the undesired ketal **14**, which supported the structural assignment of **11a** (Scheme 3).



Scheme 1 Retrosynthetic analysis.



Scheme 2 Reagents and conditions: (a) NaH, n-BuLi, 4-bromo-2-methyl-2-butene, THF,  $0^\circ\text{C}$ , 79%; (b)  $\text{SnCl}_4$ ,  $\text{CH}_2\text{Cl}_2$ , r.t., 81%; (c)  $\text{KOCH}_3$ , MVK,  $0^\circ\text{C}$ ;  $\text{K}_2\text{CO}_3$ ,  $\text{CH}_3\text{OH}$ , reflux, 80%; (d) LDA, allylic bromide,  $-78^\circ\text{C} \rightarrow \text{r.t.}$ , 63% (**11a**), 9% (**11b**); (e)  $\text{LiAlH}_4$ , THF, r.t., 76%; (f) DDQ, benzene, r.t., 70%.






Fig. 1 Structures of spiniferin-1 and dihydrospininiferin-1.



Scheme 3 The confirmation of relative configuration of **11a**.



**Scheme 4** Reagents and conditions: (a)  $\text{HCF}_2\text{CF}_2\text{OCF}_2\text{CF}_2\text{SO}_2\text{F}$ , DBU, THF, 0 °C;  $\text{NEt}_3$ , reflux, 82%; (b)  $\text{O}_3$ , -70 °C,  $\text{CH}_3\text{OH}$ ;  $\text{Me}_2\text{S}$ , r.t., 67%; (c) 12 N HCl, THF, r.t. 69%.

Using a novel poly(per)fluoroalkanosulfonyl fluoride induced tandem carbonium rearrangement which was recently found by us,  $\gamma$ -hydroxymethyl cyclohexenone **13** was transformed into our desired cyclopropyl enone **15** in 82% yield<sup>†</sup>. Selective ozonolysis<sup>12</sup> of compound **15** furnished aldehyde **4**. The latter in the presence of 12 N HCl rearranged to cycloheptatriene **3**, a unstable intermediate, which immediately cyclized to form ( $\pm$ )-**2** as a colorless oil (Scheme 4). Our synthetic material has identical  $^1\text{H}$  and  $^{13}\text{C}$  NMR data<sup>‡</sup> as reported by Cimino<sup>1,2</sup> and Mashall.<sup>3,4</sup>

## Notes and references

<sup>†</sup> Synthesis of key intermediate **15**: To a stirred, cooled solution of 99 mg of enone **13** (0.4 mmol) in 2.5 ml of dry THF was added 119  $\mu\text{l}$  (2 eq) of DBU and 144  $\mu\text{l}$  (2 eq) of  $\text{HCF}_2\text{CF}_2\text{OCF}_2\text{CF}_2\text{SO}_2\text{F}$ . The solution was stirred for 30 min and was filtered through a short silica gel column with the aid of 2.5 ml of THF. The combined solution was added 280  $\mu\text{l}$  (5 eq.)  $\text{NEt}_3$  and was heated to reflux for 8 h. The solvent was removed under reduced pressure and the remainder was purified by column chromatography on silica gel to afford 75 mg (82%) of **15** as a colorless oil. IR (film):  $\nu$  1666, 1640  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  7.08 (s, 1H), 5.77 (m, 1H), 5.01 (m, 2H), 2.92 (m, 1H), 2.84 (d,  $J$  = 8.6 Hz, 1H), 2.51 (d,  $J$  = 8.6 Hz, 1H), 1.11 (s, 3H), 1.10 (s, 3H), 0.51 (d,  $J$  = 4.0 Hz, 3H); EIMS:  $m/e$  230 ( $\text{M}^+$ , 100%)

<sup>‡</sup> Procedure for the synthesis of ( $\pm$ )-**2**: To a stirred solution of 62 mg **4** (0.27 mmol) in 3 ml THF was added 0.6 ml 12 N HCl. The solution was stirred for 1 h and was extracted with ether. The combined organic layers was

washed with brine and dried. The solvent was removed under reduced pressure and the remainder was purified by column chromatography on silica gel to afford 40 mg (69%) of ( $\pm$ )-**2** as a colorless oil.  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  7.33 (d,  $J$  = 1.8 Hz, 1H), 6.56 (d,  $J$  = 1.5 Hz, 1H), 6.29 (s, 1H), 6.28 (s, 1H), 3.14 (d,  $J$  = 10.8 Hz, 1H), 2.2–2.5 (m, 2H), 1.7–1.9 (m, 2H), 1.3–1.5 (m, 1H), 1.32 (s, 3H), 1.14 (d,  $J$  = 10.8 Hz, 1H), 1.05 (s, 3H), 0.6–0.8 (m, 1H);  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ )  $\delta$  153.2, 140.3, 132.5, 129.4, 125.3, 112.2, 110.0, 109.1, 42.0, 37.8, 36.7, 35.0, 29.2, 27.1, 25.8.

- G. Cimino, S. De Stefano, L. Minale and E. Trivellone, *Tetrahedron Lett.*, 1975, **43**, 3727.
- G. Cimino, S. De Stefano, L. Minale and E. Trivellone, *Experientia*, 1978, **34**, 1425.
- J. A. Marshall and R. E. Conrow, *J. Am. Chem. Soc.*, 1980, **102**, 4274.
- J. A. Marshall and R. E. Conrow, *J. Am. Chem. Soc.*, 1983, **105**, 5679.
- (a) E. Vogel and H. D. Roth, *Angew. Chem., Int. Ed. Engl.*, 1964, **3**, 228; (b) E. Vogel and W. A. Böll, *Angew. Chem., Int. Ed. Engl.*, 1964, **9**, 642; (c) E. Vogel, W. Pretzer and W. A. Böll, *Angew. Chem., Int. Ed. Engl.*, 1964, **9**, 642.
- (a) Z. Zhu, W. S. Tian and Q. J. Liao, *Tetrahedron Lett.*, 1996, **37**, 8553; (b) Z. Zhu, W. S. Tian, Q. J. Liao and Y. K. Wu, *Bioorg. Med. Chem. Lett.*, 1998, **8**, 1949; (c) X. S. Feng, W. S. Tian and Q. Y. Chen, *Bioorg. Med. Chem. Lett.*, 1997, **7**, 3113; (d) X. S. Feng, W. S. Tian and Q. Y. Chen, *J. Chem. Soc., Perkin Trans. 1*, 1998, **6**, 1139; (e) Z. H. Yan, Ph D dissertation, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 1999.
- (a) W. S. Tian, Z. Lei, L. Chen and Y. Huang, *J. Fluorine Chem.*, 2000, **101**, 305; (b) Z. Lei, Masters dissertation, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 1997; (c) L. Chen, PhD dissertation, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 2000.
- S. N. Huckin and L. Weiler, *J. Am. Chem. Soc.*, 1974, **96**, 1082.
- (a) M. T. Reetz, I. Chatziiosifidis and K. Schwellnus, *Angew. Chem., Int. Ed. Engl.*, 1981, **20**, 687; (b) F. W. Sum and L. Weiler, *Can. J. Chem.*, 1979, **57**, 1431.
- G. Stork and R. L. Danheiser, *J. Org. Chem.*, 1973, **38**, 1775.
- (a) D. Burn, V. Perrow and G. O. Weston, *Tetrahedron Lett.*, 1960, **2**, 14; (b) D. Burn, J. W. Ducker, B. Ellis, A. K. Hiscock, A. P. Leftwick, C. M. Peach, V. Petrow and D. M. Williamson, *J. Chem. Soc.*, 1963, 4242.
- T. Veysoglu, L. A. Mitscher and J. K. Swayze, *Synthesis*, 1980, **10**, 807.