

Effect of Linker Length on DNA-binding Affinity, Cross-linking Efficiency and Cytotoxicity of C8-linked Pyrrolobenzodiazepine Dimers

D. Subhas Bose,^a Andrew S. Thompson,^a Melissa Smellie,^b Mark D. Berardini,^b John A. Hartley,^b Terence C. Jenkins,^c Stephen Neidle^c and David E. Thurston*^a

^a Division of Medicinal Chemistry, School of Pharmacy and Biomedical Sciences, University of Portsmouth, King Henry 1st Street, Portsmouth PO1 2DZ, UK

^b Department of Oncology, University College and Middlesex School of Medicine, 91 Riding House Street, London W1P 8BT, UK


^c Cancer Research Campaign Biomolecular Structure Unit, The Institute of Cancer Research, Sutton, Surrey SM2 5NG, UK

An efficient synthesis of a homologous series of C8-linked pyrrolobenzodiazepine dimers is reported; compounds with an odd number of methylenes ($n = 3$ or 5) in the linker show a higher affinity for DNA, enhanced cross-linking efficiency, and are more cytotoxic compared with compounds with either $n = 4$ or 6 .

It is generally accepted that a DNA interstrand cross-link is a highly cytotoxic lesion,^{1a} and many bifunctional alkylating agents such as the nitrogen mustards and cisplatin are used clinically as antitumour agents.^{1b} Although the basis for tumour-cell selectivity remains unknown, it is possible that cross-linked DNA adducts prove challenging to cellular repair mechanisms,² or that the agents target genes associated with the control of cell growth.³ We recently reported⁴ the design and synthesis of a novel irreversible guanine-specific DNA interstrand cross-linking agent, DSB-120 (**8a**; $n = 3$), based on the naturally occurring pyrrolo[2,1-c][1,4]benzodiazepine (PBD) antitumour antibiotic DC-81 (**10**),⁵ which bonds in the minor groove of DNA *via* covalent interaction between the C11-position and N2 of guanine. PBD monomers are known to be sequence selective with a preference for PuGpu triplets.⁶ NMR spectroscopy and modelling studies indicate that DSB-120 spans six DNA base-pairs (twice the binding-site size of DC-81) and actively recognises a 5'-GATC sequence.⁴ We report here the synthesis of a homologous series of C8-linked bifunctional DNA alkylating agents in order to probe the

effect of linker length on binding affinity, DNA cross-linking efficiency and cytotoxicity.

The overall synthetic strategy is shown in Scheme 1. A versatile approach has been developed to join two units of vanillic acid **1** with α,ω -dihaloalkanes of varying length to provide the dimer acids **2a–d**. Dibromo- and dichloro-alkanes gave inferior yields compared to the diiodo-compounds, and varying the chain length from $n = 3$ to 6 did not affect the efficiency of the reaction. By using this approach, the formation of mixtures of mono- and bis-alkylated products is avoided. The procedure involved refluxing **1** with the diiodoalkanes in the presence of aq. NaOH for 48 h to afford **2a–d** in 33–65% yields. However, all attempts to obtain the nitro acids of type **5** by direct nitration of **2a–d** failed using a variety of reaction conditions, including NaNO₂–H₂SO₄, SnCl₄–HNO₃ and H₂SO₄–HNO₃, owing to the insoluble nature of the dimer acids. Following conversion into the corresponding methyl esters **3a–d**, nitration with SnCl₄–HNO₃ proceeded smoothly to afford the nitro esters **4a–d** in high yield. An initial attempt to hydrolyse the methyl ester **4a**



Scheme 1 Reagents and conditions: i, I-(CH₂)_n-I, aq. NaOH, THF, reflux, 48 h; ii, dimethyl sulfate, K₂CO₃, acetone, reflux, 30 min; iii, SnCl₄, HNO₃, CH₂Cl₂, -20 °C, 15 min; iv, aq. NaOH, THF; v, oxalyl chloride, THF, Et₃N, H₂O, (2*S*)-pyrrolidine-2-carbaldehyde diethyl thioacetal,⁸ 4 h; vi, SnCl₂·2H₂O, MeOH, reflux, 40 min; vii, HgCl₂, CaCO₃, MeCN, H₂O, 2.5 h

Table 1 Thermal denaturation (ΔT_m values), DNA cross-linking efficiency, and *in vitro* cytotoxicity data for the C8-linked PBD dimers and DC-81


Compound	Linker <i>n</i> =	ΔT_m /°C ^a	C ₅₀ %/μmol dm ^{-3b}	IC ₅₀ /μmol dm ⁻³			
				L1210 ^c	ADJ/PC6 ^c	CH1 ^c	K ₅₆₂ ^d
8a	3	15.1	0.055	0.01	0.0005	0.003	0.2
8b	4	4.1	1.00	1.2	0.35	0.05	2.5
8c	5	8.1	0.070	0.0045	0.0004	0.00032	0.5
8d	6	7.0	0.750	0.34	0.002	0.002	1.0
DC-81 (10)	—	0.7	—	0.38	0.33	0.1	NA
Melphalan	—	—	20.0	3.0	0.02	2.0	35

^a Thermal denaturation studies with calf thymus DNA (see text). ^b Concentration of drug required for 50% cross-linking of pBR322 DNA (see text). ^c IC₅₀ is the drug dose that inhibits cell growth by 50% compared with solvent controls. Compounds were dissolved in DMSO to provide a final concentration of 0.05% DMSO. Incubation times (37 °C) were: L1210, 3 days; ADJ/PC6, 4 days; CH1, 9 days; ^d K₅₆₂ is a human leukaemia cell line in which IC₅₀ values were measured using an MTT assay following a 1 h exposure to drug. NA = Result not available.

by refluxing in aq. NaOH gave a high-melting solid identified as the diacid hydrolysis product **9** in which demethylation of the aromatic ether had occurred. The electron-withdrawing *p*-nitro groups may be responsible for this phenomenon.⁷ However, mild hydrolysis of the ester with aq. NaOH at room temperature for 6 h afforded **5a** in quantitative yield. Coupling of nitro acids **5a–d** with (2*S*)-pyrrolidine-2-carbaldehyde diethyl thioacetal⁸ afforded the bis(amides) **6a–d** in approximately 65% yield, which were subsequently reduced to the amino thioacetals **7a–d**. Cyclization⁸ with HgCl₂–CaCO₃ afforded the target C8-dimers **8a–d** in good yields.[†]

The general DNA-binding affinity of PBD dimers **8a–d** was examined by thermal denaturation studies.⁹ For a 5:1 mol ratio of DNA-ligand (calf thymus DNA concentration = 100 μmol dm⁻³, 10 mmol dm⁻³ phosphate buffer, pH 7.0) an increase in helix melting temperature (ΔT_m) is observed for each dimer compared with untreated control DNA following incubation at 37 °C for 18 h (Table 1). In the same experiment, the monomer DC-81 (**10**) gives only a small increase in T_m , consistent with the notion that the PBD dimers **8a–d** stabilise DNA through the formation of interstrand cross-links.

The DNA cross-linking efficiency of dimers **8a–d** was investigated using an assay involving linear double-stranded



DNA derived from the plasmid pBR322 (4362 bp, linearised with *Hind* III and then ³²P-end-labelled).¹⁰ Following complete denaturation to the single-stranded form, the presence of an interstrand cross-link results in renaturation to double-stranded DNA during electrophoresis in a neutral agarose gel (Fig. 1). Quantitation of the autoradiograph using laser densitometry allowed calculation of the concentration of each agent required to effect 50% cross-linking (Table 1).

Compounds **8a** (*n* = 3) and **8c** (*n* = 5) are broadly similar in cross-linking efficiency, whereas dimers **8b** (*n* = 4) and **8d** (*n* = 6) approximately 18- and 14-fold less efficient, respectively. This data is consistent with the determined ΔT_m values, which reflect the differences in ability of the compounds to stabilise DNA helix-coil transitions. Compound **8a** binds to double-stranded DNA more tightly than **8c** even though they have similar cross-linking efficiencies. This may reflect differences in their non-covalent binding, since they differ in isohelicity with respect to the DNA minor groove. It is also possible that mono adducts or intrastrand cross-links can be formed that would influence the thermal denaturation data but not the cross-linking assay.

Interestingly, *in vitro* cytotoxicity data in human K₅₆₂ and rodent ADJ/PC6 cell lines (Table 1) correlate with both the

[†] All compounds gave satisfactory spectral data: selected data for **8a**: ¹H NMR (CDCl₃, 270 MHz): δ 2.01–2.17 (m, 2H), 2.28–2.45 (m, 8H), 3.50–3.87 (m, 6H), 3.92 (s, 6H), 4.22–4.33 (m, 4H), 6.85 (s, 2H), 7.51 (s, 2H), 7.66 (d, 2H, *J* 4.4 Hz). ¹³C NMR (CDCl₃): δ 24.2, 28.8, 29.6, 46.7, 53.7, 56.1, 65.4, 110.7, 111.6, 120.3, 140.6, 147.8, 150.6, 162.4, 164.6.




Fig. 1 Autoradiograph of a neutral agarose gel showing DNA interstrand cross-linking by **8a-d** in linear ^{32}P -end-labelled pBR322 DNA. Drug reactions (2 h at 37°C) were in 25 mmol dm^{-3} triethanolamine/ 1 mmol dm^{-3} EDTA pH 7.2 buffer with 10 ng DNA in a final volume of 50 μl . Reaction was terminated by addition of an equal volume of 0.6 mol dm^{-3} sodium acetate, 20 mmol dm^{-3} EDTA and $100\text{ }\mu\text{g cm}^{-3}$ tRNA, and the DNA precipitated with ethanol. Dried pellets were taken up in strand separation buffer (30% DMSO in 1 mmol dm^{-3} EDTA). Denaturation for 2 min at 90°C was followed by immediate chilling in an ice-water bath. Electrophoresis was carried out on 0.8% submerged horizontal agarose gels at 40 V for 16 h with tris(acetate) running buffer. Double- (DS) and single-stranded (SS) DNA were quantitated by laser densitometry. O (nd) is the non-denatured double-stranded DNA control.

thermal denaturation data and the cross-linking efficiencies. There is a similar correlation for the rodent L1210 and human CH1 cell lines, except that the positions of **8a** and **8c** in the overall rank-order are transposed. As shown in Table 1, **8a** is approximately 300-fold more efficient at cross-linking DNA than the clinically-used major-groove cross-linking agent melphalan, and is significantly more cytotoxic across the four cell lines studied. Considering the lower activity of the monomer DC-81, this data suggests that interstrand cross-links probably represent the cytotoxic lesions leading to cell death.

In summary, an efficient synthesis of C8-linked pyrrolo-benzodiazepine dimers of varying linker length has been achieved. We find a significant correlation between DNA-binding affinity, cross-linking efficiency and cytotoxicity, with the $n = 3$ and $n = 5$ homologues **8a** and **8c** displaying optimal effects compared to the dimers with an even number of methylenes in the linker.

We acknowledge generous support for various aspects of this work from the SERC (GR/F52675) and the Cancer Research Campaign. Professor Ken Harrap and Phyllis Goddard (Institute of Cancer Research) are thanked for providing the cytotoxicity data.

Received, 9th July 1992; Com. 2103631B

References

- (a) K. W. Kohn, in *Molecular Aspects of Anticancer Drug Action*, ed. S. Neidle and M. J. Waring, Macmillan Press, London, 1983, p. 315-361; (b) W. B. Pratt and R. W. Ruddon, *The Anticancer Drugs*, OUP, New York, 1979.
- P. B. Hopkins, T. J. Millard, J. Woo, M. F. Weidner, J. J. Kirchner, S. Th. Sigurdsson and S. Raucher, *Tetrahedron*, 1991, **47**, 2475.
- W. B. Mattes, J. A. Hartley, K. W. Kohn and D. W. Matheson, *Carcinogenesis*, 1988, **9**, 2065.
- D. S. Bose, A. S. Thompson, J. Ching, J. A. Hartley, M. D. Berardini, T. C. Jenkins, S. Neidle, L. H. Hurley and D. E. Thurston, *J. Am. Chem. Soc.*, 1992, **114**, 4939.
- D. S. Bose, G. B. Jones and D. E. Thurston, *Tetrahedron*, 1992, **48**, 751, and references cited therein.
- L. H. Hurley, T. Reck, D. E. Thurston, D. R. Langley, K. G. Holden, R. P. Hertzberg, J. R. E. Hoover, G. Gallagher, Jr., L. F. Faucette, S.-M. Mong and R. K. Johnson, *Chem. Res. Toxicol.*, 1988, **1**, 258.
- M. Greenwood and R. Robinson, *J. Chem. Soc.*, 1932, 1370.
- D. R. Langley and D. E. Thurston, *J. Org. Chem.*, 1987, **52**, 91.
- G. B. Jones, C. L. Davey, T. C. Jenkins, A. Kamal, G. G. Kneale, S. Neidle, G. D. Webster and D. E. Thurston, *Anti-Cancer Drug Design*, 1990, **5**, 249.
- J. A. Hartley, M. D. Berardini and R. L. Souhami, *Anal. Biochem.*, 1991, **193**, 131.