## Scalable Total Synthesis of (–)-Berkelic Acid Using a Protecting-Group-Free Strategy\*\*

Francisco J. Fañanás,\* Abraham Mendoza, Tamara Arto, Baris Temelli, and Félix Rodríguez\*

(–)-Berkelic acid (1; Figure 1) is a biologically active and architecturally unique secondary metabolite that was isolated in 2006 by Stierle and co-workers from a extremophilic



*Figure 1.* Structure of (–)-berkelic acid (1).

Penicillium species obtained from the surface waters of Berkeley Pit Lake in Butte, Montana (USA).<sup>[1]</sup> This lake was formed when an abandoned copper mine was flooded with infiltrating ground water, thus resulting in the formation of a highly acidic (pH 2.5) and metal contaminated ecosystem where only extremophiles could survive. The microorganisms that tolerate these extreme growth conditions are a precious, although very scarce, source of architecturally nonconventional and potentially bioactive natural products.<sup>[2]</sup> A good example of this is (-)-berkelic acid (1), which has a unique structure and a diverse biological profile. This compound was reported to inhibit the cysteine protease caspase-1 (98 µм) and matrix metalloproteinase MMP-3 (1.87 µm), as well as displayed selective activity against the human ovarian cancer line OVCAR-3 (GI50 = 91 nM).<sup>[1]</sup> To date, three research groups have accomplished the total synthesis of **1**.<sup>[3,4]</sup> It should be noted that the strategies used in all these total syntheses only delivered very low quantities of the natural product (3-11 mg). Nevertheless, they provided material enough for

 [\*] Prof. Dr. F. J. Fañanás, Dr. A. Mendoza, T. Arto, Dr. B. Temelli,<sup>[+]</sup> Prof. F. Rodríguez Instituto Universitario de Química Organometálica "Enrique Moles", Unidad Asociada al C.S.I.C., Universidad de Oviedo Julián Clavería, 8, 33006 Oviedo (Spain) E-mail: fjfv@uniovi.es frodriguez@uniovi.es
[<sup>+</sup>] Permanent address: Hacettepe University Department of Chemistry, 06800 Ankara (Turkey)

[\*\*] We gratefully acknowledge financial support from the MICINN of Spain (grant CTQ2010-16790), MEC (FPU-predoctoral grant to A.M.), FICYT of Principado de Asturias (Severo Ochoa predoctoral grant to T. A.), and TUBITAK-BIDEB Programme of Turkey (postdoctoral fellowship to B.T.). We also thank Prof. Dr. José Barluenga (Universidad de Oviedo-Spain) for his support.

Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/anie.201109076.

some new bioactivity studies which led to some contradictory results when compared to those reported by the isolation team.<sup>[5]</sup> So, the necessity for an in depth study on the biological activity of  $\mathbf{1}$  as well as the more than likely future elimination of its natural source mandates a secured material supply.

Herein, we report a total synthesis of berkelic acid wherein all but the last step were executed on gram scale. This level of practicality was enabled by a highly stereoconvergent and modular synthetic strategy.

Recently, we developed a palladium(II)-catalyzed onepot three-component coupling reaction for the diastereoselective construction of chroman spiroacetals (Scheme 1 a).<sup>[6]</sup>



b)



**Scheme 1.** a) Synthesis of spiroacetals through a one-pot multicomponent cascade reaction (previous work; see Ref. [6]). b) Our strategy for the synthesis of 1 (this work).

The method relies on the in situ formation of an exocyclic enol ether and a *o*-quinonemethide, and the subsequent formal cycloaddition reaction between these two reagents. In principle, the reaction seems ideal for the construction of the berkelic acid skeleton; however, an easy way to introduce the additional pyran ring was initially far from evident. Upon

Angew. Chem. Int. Ed. 2012, 51, 1-5

© 2012 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim





considering the ability of  $\pi$ -acid catalysts to activate alkynes, thus facilitating the intramolecular nucleophilic addition of alcohols and aldehydes,<sup>[7]</sup> we hypothesized that ideally a unique metal complex could promote the cycloisomerization of alkyndiol **2** to give the enol ether **5** and also the cycloisomerization of aldehyde **3** to give **6** (Scheme 1 b). The formal cycloaddition reaction between the intermediates **5** and **6** would result in the formation of the core structure of berkelic acid **4** in an apparently very simple way.

Although the proposed reaction was risky in terms of stereoselectivity, because just one chiral center (that of alkyndiol 2) would induce the selective formation of the other three newly formed ones in 4 (including a desymmetrization in 2), the process was very attractive from the synthetic point of view as it avoided the synthesis of starting materials having multiple stereocenters and tedious protection/deprotection steps. Once the compound 4 was obtained we were confident about the possibilities of completion the total synthesis of (-)-berkelic acid (1), because only the installation of the lateral chain and a few conventional functional group manipulations would remain.

Preparation of the key building blocks, **2** and **3**, was achieved in a concise and scalable way from commercially available starting materials. Specifically, chiral fragment **2** was synthesized from (*S*)-(–)-3-butyn-2-ol (**7**; 98% *ee*) in just three conventional chemical transformations (Scheme 2). Thus, initial mesylation of alcohol **7** led to the corresponding derivative **8**. Displacement of the mesylate by diethylmalonate in the presence of cesium fluoride through an  $S_N^2$  reaction furnished the diester **9**. Finally, reduction with lithium aluminium hydride provided the desired diol **2** in 70% overall yield and 96% *ee* (2.8 g scale).<sup>[8]</sup> Synthesis of building block **3** was also achieved in three steps (Scheme 2). Namely, triflic anhydride reacted selectively with the hydroxy



**Scheme 2.** Synthesis of building blocks **2** and **3**. Reagents and conditions: a) MsCl, Et<sub>3</sub>N, CH<sub>2</sub>Cl<sub>2</sub>, 0°C; b) diethyl malonate, CsF, THF, 45°C; c) LAH, THF, 75°C; d) Tf<sub>2</sub>O, 2,6-lutidine, CH<sub>2</sub>Cl<sub>2</sub>, RT; e) potassium trifluoro(hept-1-yn-1-yl)borate, DIPEA, [PdCl<sub>2</sub>(dppf)] (5 mol%), MeOH, 65°C; f) formaldehyde, CaCl<sub>2</sub>·2 H<sub>2</sub>O, KOH, MeOH, RT and then MnO<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>, RT. DIPEA=*N*-ethyl-*N*,*N*-diisopropylamine, dppf=1,1'-bis(diphenylphosphino)ferrocene, LAH = lithium aluminium hydride, Ms = methanesulfonyl, Tf = trifluoromethylsulfonyl.

group in *para*-position of commercially available ester methyl 2,4,6-trihydroxybenzoate (**10**) to provide compound **11**.<sup>[3b]</sup> Reaction of this intermediate through a Sonogashira-type cross-coupling reaction employing the potassium trifluoroborate salt derived from 1-heptyne was successful, thus allowing the synthesis of alkyne **12** in high yield.<sup>[9]</sup> Finally, introduction of the aldehyde functionality was achieved using a sequence involving an initial hydroxymethylation with formaldehyde in the presence of calcium chloride and subsequent oxidation with manganese dioxide. Every reaction was performed on a gram scale, thus providing the fragment **3** in 60% overall yield from **10** (4.8 g scale).

At this stage, with substantial quantities of fragments 2 and 3 in hand, we turned our attention to the key reaction of our synthesis, that is, the catalytic cycloisomerization of these compounds and subsequent formal cycloaddition reaction between the in situ formed intermediates 5 and 6 to give product 4 (Scheme 1). Our experience in cycloisomerization reactions indicated that our desired transformation could be catalyzed by a metallic complex derived from typical carbophilic Lewis acids.<sup>[10]</sup> In fact, we found that the reaction proceeded in the presence of 5 mol% of AgOTf. To avoid possible decomposition of compound 4 as a result of the presence of the new reactive pyran ring, we directly performed the hydrogenation of the carbon–carbon double bond of this pyran ring under conventional conditions (Scheme 3).



**Scheme 3.** Synthesis of the central core of **1**. Reagents and conditions: a) AgOTf (5 mol%), THF, 0°C and then  $H_2$ , Pd/C (5 mol%), MeOH, -5°C. THF = tetrahydrofuran.

It should be remarked that in this global transformation four new chiral centers are formed and, surprisingly, only two diastereoisomers were observed in the crude reaction mixture (d.r. = 2:1). Even more pleasant was the confirmation that the structure of the major diastereoisomer corresponded to that of the desired product **13** with all chiral centers and functionalities resembling those of the natural product. The minor isomer could be easily separated at an advanced stage, as it will be shown. Notably, this reaction could be performed on a gram scale with 83 % yield (2.4 g of this material were synthesized). This new silver-catalyzed reaction allowed the assembly of the central core of the natural product, which contains four rings and five stereocenters, in just one step.

At this point we had achieved a formal synthesis of berkelic acid because intermediate **13** was an advanced intermediate in the total synthesis reported by Fürstner and co-workers.<sup>[3c]</sup> It should be also remarked that the introduction of the lateral chain at an advanced stage is beneficial and highly desirable because it makes the synthesis much more

www.angewandte.org © 2013 These are not the final page numbers! modular and convergent. Thus, the required building blocks are simpler and the synthesis of analogues of the natural product in the quest for new and relevant bioactive compounds is much easier. In this context, the syntheses reported by the groups of Fürstner and Snider are remarkable.<sup>[3a,c]</sup> The coupling of the lateral chain in both routes relies on the use of an aldol reaction which requires a final oxidation step. This late-stage oxidation was problematic, as the electron-rich aromatic ring of the central core was highly susceptible to oxidation and protection of the free phenol as an allyl or TBS ether was required.

Having in mind these problems and considering that a scalable total synthesis should avoid tedious protection/ deprotection steps, we devised a different strategy to introduce the lateral chain (Scheme 4). So, to avoid any redox



**Scheme 4.** Our devised strategy to introduce the lateral chain. TES = triethylsilyl.

process, we thought of using an umpolung alkylation reaction with iodide **14** and an appropriate acyl anion equivalent derived from aldehyde **15**. This reaction would deliver the berkelic acid methyl ester **16**, a direct precursor of the natural product **1**, in one step. Between the different acyl anion equivalents reported, we considered the use of the cyanohydrin **17** (Scheme 4).<sup>[11]</sup> We were able to synthesize this compound on a gram scale from commercially available dimethyl-D-malate (**18**) through the known hydroxyacid **19** (Scheme 5).<sup>[12]</sup> The cyanohydrin **17**,<sup>[13]</sup> obtained in 31 % overall yield (1.0 g scale) as an inconsequential mixture of isomers at the cyanohydrin center, could be stored for long periods of time without problems.

With alcohol **13** and cyanohydrin **17** in hand we tackled the completion of the synthesis (Scheme 6). Thus, treatment



**Scheme 5.** Synthesis of cyanohydrin **17**.Reagents and conditions: a) 1. Bu<sub>4</sub>NIO<sub>4</sub>, CHCl<sub>3</sub>, 65 °C; 2. TESCN, PNPCl, neat, RT. PNPCl=bis-(triphenylphosphoranylidene)ammonium chloride.

Angew. Chem. Int. Ed. 2012, 51, 1-5

© 2012 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim



**Scheme 6.** Final steps. Reagents and conditions: a)  $I_2$ , PPh<sub>3</sub>, imidazole, Et<sub>2</sub>O/CH<sub>3</sub>CN (3:1), RT; b) 1. **17**, LDA, DMPU, THF, -78 °C and then **14**; 2. TBAF-3 H<sub>2</sub>O, MeOH, RT; c) (Bu<sub>3</sub>Sn)<sub>2</sub>O, toluene, 115 °C. DMPU=1,3-dimethyl-3,4,5,6-tetrahydropyrimidin-2(1*H*)-one, LDA= lithium diisopropylamide, TBAF = tetrabutylammonium fluoride.

of alcohol 13 with PPh<sub>3</sub> and  $I_2$  gave the iodide 14 as a crystalline solid. This was especially convenient as a single recrystallization from a mixture of dichloromethane and ethanol removed the undesired minor diastereoisomer produced in the previous step, thus allowing the isolation of 14 in 52% yield and enantiomerically pure form (1.6 g of 14 were obtained).<sup>[14]</sup> Additional treatment of this iodide 14 with the anion derived from cyanohydrin 17 finally furnished the diester 16 in 88 % yield.<sup>[15]</sup> Again, this transformation could be performed on a gram scale (1.2 g of 16 was synthesized in one batch). X-ray crystallographic analysis of 16 confirmed it as the berkelic acid methyl ester.<sup>[16]</sup> It should be noted that diester 16 is a stable compound. In fact, we stored some of this product and we have not observed any decomposition. However as reported by other authors.<sup>[3c]</sup> **1** is not as stable as its ester 16. So, to keep in our hands enough material for transformation into the natural product when required, the last step of our total synthesis was conducted on a small scale using (Bu<sub>3</sub>Sn)<sub>2</sub>O for the selective cleavage the benzylic ester (Scheme 6). It should be noted that after a slight modification of the protocol described by De Brabander and co-workers to perform this last step,<sup>[3b]</sup> we were able to obtain 25 mg of compound 1 in one batch and we did not observe the formation of any other undesired product. The spectral data and optical rotation of synthetic 1 were in agreement with those for the natural sample and the synthetic (-)-berkelic acid reported in the literature.<sup>[1,3]</sup>

In summary, a practical and scalable route to (-)-berkelic acid (1) has been developed. The completed synthesis as it stands was accomplished with seven steps in the longest linear sequence and no more than 14 overall steps from three commercial materials (methyl 2,4,6-trihydroxybenzoate, (S)-(-)-3-butyn-2-ol, and dimethyl-D-malate). Notably, from (S)-(-)-3-butyn-2-ol, containing just one stereocenter, we

These are not the final page numbers!

www.angewandte.org



obtained the central core of the natural product, which contains five stereocenters, without the need of any other chiral reagent or catalyst. This transformation was achieved by a new silver-catalyzed reaction which allowed the construction of this central core in just one step. Also, a new strategy to introduce the lateral chain was applied. Moreover, all but the last step were conducted on a gram scale. As a result, we believe that the supply of **1** should no longer be an issue. It is also important to remark that the synthesis described herein does not require any protection/deprotection steps. Finally, the modular strategy should allow the easy production of several analogues of the natural product for our ongoing biological activity studies.

Received: December 22, 2011 Published online:

**Keywords:** antitumor agents · multicomponent reactions · natural products · spiro compounds · total synthesis

- [1] A. A. Stierle, D. B. Stierle, K. Kelly, J. Org. Chem. 2006, 71, 5357-5360.
- [2] Z. E. Wilson, M. A. Brimble, Nat. Prod. Rep. 2009, 26, 44-71.
- [3] a) X. Wu, J. Zhou, B. B. Snider, Angew. Chem. 2009, 121, 1309–1312; Angew. Chem. Int. Ed. 2009, 48, 1283–1286; b) C. F. Bender, F. K. Yoshimoto, C. L. Paradise, J. K. De Brabander, J. Am. Chem. Soc. 2009, 131, 11350–11351; c) T. N. Snaddon, P. Buchgraber, S. Schulthoff, C. Wirtz, R. Mynott, A. Fürstner, Chem. Eur. J. 2010, 16, 12133–12140.
- [4] For studies toward the synthesis of (-)-berkelic acid and model compounds, see: a) J. Zhou, B. B. Snider, Org. Lett. 2007, 9, 2071-2074; b) M. A. Marsini, Y. Huang, C. C. Lindsey, K.-L. Wu, T. R. R. Pettus, Org. Lett. 2008, 10, 1477-1480; c) Y. Huang, T. R. R. Pettus, Synlett 2008, 1353-1356; d) P. Buchgraber, T. N. Snaddon, C. Wirtz, R. Mynott, R. Goddard, A. Fürstner, Angew. Chem. 2008, 120, 8578-8582; Angew. Chem. Int. Ed. 2008, 47, 8450-8454; e) N. Stiasni, M. Hiersemann, Synlett 2009, 2133-2136; f) Z. E. Wilson, M. A. Brimble, Org. Biomol. Chem. 2010, 8, 1284-1286; g) T. A. Wenderski, M. A. Marsini, T. R. R. Pettus, Org. Lett. 2011, 13, 118-121.
- [5] X. Wu, J. Zhou, B. B. Snider, J. Org. Chem. 2009, 74, 6245-6252.
- [6] J. Barluenga, A. Mendoza, F. Rodríguez, F. J. Fañanás, Angew. Chem. 2009, 121, 1672–1675; Angew. Chem. Int. Ed. 2009, 48, 1644–1647. Yet in this paper we indicated the possibility of applying the reported strategy for the synthesis of berkelic acid. Later on, De Brabander and co-workers reported their total synthesis following a similar strategy (see Ref. [3b]).

- [7] A. Fürstner, P. W. Davies, Angew. Chem. 2007, 119, 3478–3519; Angew. Chem. Int. Ed. 2007, 46, 3410–3449.
- [8] To determine the enantiomeric excess, compound 2 was reacted through a Sonogashira reaction with phenylbromide and the enantiomeric excess was measured on the corresponding arylated product obtained (see the Supporting Information for details).
- [9] G. A. Molander, B. W. Katona, F. Machrouhi, J. Org. Chem. 2002, 67, 8416–8423.
- [10] a) J. Barluenga, J. Calleja, A. Mendoza, F. Rodríguez, F. J. Fañanás, *Chem. Eur. J.* 2010, *16*, 7110-7112; b) J. Barluenga, A. Fernández, A. Diéguez, F. Rodríguez, F. J. Fañanás, *Chem. Eur. J.* 2009, *15*, 11660-11667; c) J. Barluenga, A. Fernández, F. Rodríguez, F. J. Fañanás, *Chem. Eur. J.* 2009, *15*, 8121-8123; d) F. J. Fañanás, A. Fernández, D. Çevic, F. Rodríguez, *J. Org. Chem.* 2009, *74*, 932-934; e) J. Barluenga, A. Mendoza, F. Rodríguez, F. J. Fañanás, *Chem. Eur. J.* 2008, *14*, 10892-10895; f) J. Barluenga, A. Mendoza, F. Rodríguez, F. J. Fañanás, *Chem. Eur. J.* 2008, *14*, 10892-10895; f) J. Barluenga, A. Mendoza, F. Rodríguez, F. J. Fañanás, *Angew. Chem.* 2008, *120*, 7152-7155; *Angew. Chem. Int. Ed.* 2008, *47*, 7044-7047; g) J. Barluenga, A. Fernández, A. Satrústegui, A. Diéguez, F. Rodríguez, F. J. Fañanás, *Chem.* 2008, *14*, 4153-4156; h) J. Barluenga, A. Diéguez, F. Rodríguez, F. J. Fañanás, *Angew. Chem.* 2005, *117*, 128-130; *Angew. Chem. Int. Ed.* 2005, *44*, 126-128.
- [11] For some elegant examples on the use of cyanohydrins for alkylation processes, in the context of total synthesis, see: a) A. Fürstner, M. D. B. Fenster, B. Fasching, C. Godbout, K. Radkowski, Angew. Chem. 2006, 118, 5632-5636; Angew. Chem. Int. Ed. 2006, 45, 5506-5510; b) G. K. Packard, Y. Hu, A. Vescovi, S. D. Rychnovsky, Angew. Chem. 2004, 116, 2882-2886; Angew. Chem. Int. Ed. 2004, 43, 2822-2826.
- [12] The protocol followed is a slightly modified literature procedure (see the Supporting Information for details): P. Renaud, M. Hürzeler, D. Seebach, *Helv. Chim. Acta* **1987**, *70*, 292–298.
- [13] For the oxidation of α-hydroxyacids with Bu<sub>4</sub>NIO<sub>4</sub>, see: E. Santaniello, A. Manzocchi, C. Farachi, *Synthesis* **1980**, 563–565. For the silylcyanation of carbonyl compounds by the use of PNPCl, see: M.-A. Lacour, N. J. Rahier, M. Taillefer, *Chem. Eur. J.* **2011**, *17*, 12276–12279.
- [14] We thank Prof. A. Fürstner, Max-Planck-Institut für Kohlenforschung, Mülheim, Germany, for providing us with information on the structural determination of the iodide 14.
- [15] The use of DMPU and a careful control of the temperature in the reaction between the cyanohydrin 17 and iodide 14 are crucial for obtaining the coupling product in high yield. See: S. D. Rychnovsky, S. S. Swenson, *J. Org. Chem.* 1997, *62*, 1333–1340.
- [16] CCDC 856603 (16) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www. ccdc.cam.ac.uk/data\_request/cif.

www.angewandte.org

These are not the final page numbers!

## **Communications**



Natural Products

Scalable Total Synthesis of (–)-Berkelic Acid Using a Protecting-Group-Free Strategy



**Supply chain**: The polycyclic core of (–)berkelic acid (1) was constructed in just one step from very simple starting materials. The total synthesis of 1 involves a seven-step linear sequence. Protection/ deprotection steps were avoided and all but the last step were performed on a gram scale. This synthesis could solve the supply problem associated with the exhaustion of the natural source.

5