

Communication

Axially Chiral Dicarboxylic Acid Catalyzed Activation of Quinone Imine Ketals: Enantioselective Arylation of Enecarbamates

Takuya Hashimoto, Hiroki Nakatsu, Yuka Takiguchi, and Keiji Maruoka

J. Am. Chem. Soc., Just Accepted Manuscript • DOI: 10.1021/ja407501h • Publication Date (Web): 15 Oct 2013

Downloaded from http://pubs.acs.org on October 19, 2013

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



Journal of the American Chemical Society is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036 Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

Axially Chiral Dicarboxylic Acid Catalyzed Activation of Quinone Imine Ketals: Enantioselective Arylation of Enecarbamates

Takuya Hashimoto, Hiroki Nakatsu, Yuka Takiguchi and Keiji Maruoka*

Department of Chemistry, Graduate School of Science, Kyoto University, Sakyo, Kyoto 606-8502, Japan

Supporting Information Placeholder

ABSTRACT: The synthetic utility of quinone imine ketals in the context of asymmetric catalysis was disclosed for the first time. By expanding the utility of chiral Brønsted acid catalysis to the electrophilic activation of quinone imine ketals, we succeeded in the development of highly enantioselective arylation of encarbamates to give α -amino- β -aryl ethers wherein quinone imine ketals act as functionalized aromatic ring surrogate. Further transformations of the products were also examined to establish procedures to provide chiral β -aryl amines and α -aryl esters.

Quinones and quinone ketals, as well as cyclohexadienones, have long served as useful synthetic intermediates to construct densely functionalized aromatic rings and cyclohexanes.¹ Accordingly, their application in asymmetric catalysis, especially organocatalysis, has been on the rise as an expedient means to de-liver a range of optically enriched molecules.^{2,3} On the other hand, their nitrogen analogs, quinone imines and quinone imine ketals have rarely been utilized in asymmetric catalysis, despite their ready accessibility from the corresponding aniline derivatives⁴ and wide use in natural product synthesis.⁵⁻⁷ To the best of our knowledge, use of N-sulfonyl quinone imines in enamine catalysis as a way to perform α -arylation of aldehydes is the only example reported to date,⁸ and quinone imine ketals have never been applied in asymmetric catalysis. Moreover, regardless of the general fact that these imines can be electrophilically activated by a catalytic amount of an acid,^{6,7} chiral acid catalyzed asymmetric transformations have yet to be developed. Given our interest in the development of axially chiral dicarboxylic acid as chiral Brønsted acid catalyst and its capability to effectively activate imino functionalities,^{9,10} we became interested in its use to tap into this unexplored research area.

We report herein axially chiral dicarboxylic acid catalyzed highly enantioselective arylation of encarbamates^{11,12} using quinone imine ketals as electrophilic aryl group surrogate.^{2a,2b,8} As shown in Figure 1, we postulated the stereo-defining C-C bond formation would first occur by the nucleophilic addition of an enecarbamate to the chiral Brønsted acid-activated quinone imine ketal. From this initial product, one alkoxy group of the ketal would be expelled to generate an aromatic ring and the liberated alcohol sequentially would add to the transient imine to form an α -amino- β -aryl ether.^{13,14} In this study, we also carried out transformations of functionality-rich products into a variety of chiral building blocks, as well as a mechanistic investigation to provide more accurate understanding of the reaction.

Figure 1. Chiral Brønsted acid catalyzed reaction of quinone imine ketals and enecarbamates.



Initially, we selected N-Boc quinone imine ketal 1a and propanal-derived N-Boc enecarbamate (Z)-2a¹⁵ as model substrates and performed the reaction in the presence of 5 mol% of axially chiral dicarboxylic acid (*R*)-4a (Table 1, entry 1).^{9a} The desired α amino- β -aryl ether **3a** was obtained in good yield with high diastereo- and enantioselectivity. Prolongation of the reaction time to increase the product yield was found to be detrimental to the diastereoselectivity, as the epimerization of the hemiaminal ether moiety proceeded gradually by the reversible addition of methanol to the imine under the acidic reaction condition (entry 2).¹ Further screening of solvents revealed the effectiveness of toluene to suppress this epimerization (entries 3 and 4). Although other representative catalysts 4b and 4c also exhibited fairly good catalytic activity, 9b,9c none of them outperformed (R)-4a (entries 5 and 6). While this optimization was carried out in a test tube without any precaution against air and oxygen, we opted for the use of molecular sieves in the further study to prevent a variable degree of hydrolysis of the substrate by the adventitious water. In addition, the amount of (Z)-2a was reduced to 1.2 equiv without affecting the yield. Under the optimized reaction conditions, 3a was obtained diastereoselectively in 83% yield with 97% ee (entry 7). At this stage, we confirmed the scalability of this reaction by carrying out the reaction on 1.0 mmol scale with a reduced 2 mol% catalyst loading (entry 8).

Table 1. Optimization of the Reaction Conditions^a



^{*a*} Performed with **1a** (0.10 mmol), (*Z*)-**2a** (0.15 mmol) and (*R*)-**4** (0.005 mmol). ^{*b*} Combined yield of diastereomers. ^{*c*} Determined by ¹H NMR analysis of the crude material. ^{*d*} Determined by chiral HPLC analysis. ^{*c*} (*Z*)-**2a** (0.12 mmol) and MS4A. ^{*f*} Performed on 1.0 mmol scale with 2 mol% catalyst loading.

With the optimized reaction conditions in hand, enecarbamate substrate scope was examined (Table 2). Use of enecarbamates (Z)-2 derived from butyraldehyde and hydrocinnamaldehyde gave 3b and 3c in high yields with high stereoselectivities. Enecarbamates having an alkenyl, a silyloxy, an ester or a protected amino group could be used to give 3d-g, underlining the functional group tolerance. Due to the low reactivity of the sterically hindered enecarbamate derived from isovaleraldehyde, the reaction was carried out at rt to give 3h in 89% yield without severely compromising the selectivities. Use of the enecarbamate with a benzyloxy group gave 3i having a vicinal amino alcohol structure, although the reaction must be carried out at rt with 5.0 equiv of the enecarbamate to compensate for its low reactivity. The reaction could also be extended to the use of N-Cbz enecarbamate and N-Bz enamide to give 3j and 3k respectively, offering a way to give α -amino- β -aryl ethers having orthogonal protecting groups on two nitrogen atoms.

We then investigated the quinone imine ketal substrate scope (Table 3). A variety of 3-substituted N-Boc quinone imine ketals gave the corresponding products **31-30** in good yields with high stereoselectivities. The quinone imine ketal having a vinyl group could be applied to give **3p** despite the susceptibility of the terminal alkene to the nucleophilic attack. By use of an N-Bz quinone imine ketal, the orthogonally protected product **3q** was obtained with high enantioselectivity albeit with lower diastereoselectivity. The stereochemistry of **3m** was unambiguously determined by X-ray crystallographic analysis (see Supporting Information). In this study, it became obvious that 2-substituted quinone imine ketals have a completely different reactivity from 3-substituted ones, giving indolines as a major product.¹⁷⁻¹⁹

Table 2. Enecarbamate Scope^a



^{*a*} Performed with **1a** (0.10 mmol), (*Z*)-**2** (0.12 mmol) and (*R*)-**4a** (0.005 mmol). ^{*b*} Combined yield of diastereomers. ^{*c*} Determined by ¹H NMR analysis of the crude material. ^{*d*} Determined by chiral HPLC analysis. ^{*e*} (*Z*)-**2** (0.50 mmol).

Table 3. Quinone Imine Ketal Scope⁴



^{*a*} Performed with **1** (0.10 mmol), (*Z*)-**2a** (0.12 mmol) and (*R*)-**4a** (0.005 mmol). ^{*b*} Combined yield of diastereomers. ^{*c*} Determined by ¹H NMR analysis of the crude material. ^{*d*} Determined by chiral HPLC analysis.

Next, transformations of the products into a variety of chiral building blocks were examined (Scheme 1). Hydride reduction of **3a** and **3k** led to the formation of N-Boc β -aryl amine **7a** and N-Bz β -aryl amine **7b** in good yields with no loss of the enantiose-lectivities, respectively. To convert the hemiaminal ether moiety into an ester, **3a** was hydrolyzed into the aldehyde and subse-

quently subjected to Pinnick oxidation to give the carboxylic acid. It was necessary to carry out these two steps in one pot to minimize the deterioration of the enantioselectivity. The carboxylic acid was then transformed into methyl ester 8 in 82% overall yield. The hemiaminal ether moiety could also be exploited as a masked imine.^{12d} In the presence of a catalytic amount of (R)-4a, the addition of 1-methylindole to 3a proceeded to give 9 with high diastereoselectivity, ^{20,21} thus paving a way for one-pot sequential transformation of quinone imine ketal to 9 (see Supporting Information). We then focused on the derivatization of the aromatic rings of β -aryl amines 7a and 7b. Oxidation of 7a by PhI(OAc)₂ in aqueous solution led to the direct conversion to quinone 10. Bromination of 7a proceeded at the ortho-position of the NHBoc group to give **11**, thereby making up for the inability to use 2-substituted quinone imine ketals.¹⁷ By taking advantage of the orthogonally protected amino groups of 7b, the NHBoc moiety was selectively deprotected to give an aniline, which was then converted to iodoarene 12 by the diazotization and iodination.

Scheme 1. Transformations of α-Amino-β-Aryl Ethers



Conditions: (a) **3**, LiAlH₄ (1.5 equiv), THF, -20 °C. (b) (i) **3a**, 3N HCl, THF, 0 °C, then NaClO₂ (3 equiv), (CH₃)₂C=CHCH₃ (10 equiv), 'BuOH, rt. (ii) TMSCHN₂ (2 equiv), benzene, MeOH, rt. (c) **3a**, 1-Me-indole (2 equiv), (*R*)-**4a** (5 mol%), toluene, rt, MS4A. (d) **7a**, PhI(OAc)₂ (1.2 equiv), 'BuOH, H₂O, 0 °C. (e) **7a**, Br₂ (1.6 equiv), THF, -40 °C. (f) (i) **7b**, conc. HCl, dioxane, rt. (ii) I₂ (1.1 equiv), isoamyl nitrite (2.0 equiv), benzene, 85 °C.

To get insight on the reaction mechanism, we performed three additional experiments (Scheme 2). One was the use of (E)-2a in place of (Z)-isomer to investigate the effect of the E/Z geometry on the reactivity and selectivity. Interestingly, this study resulted in the selective formation of the same diastereomer 3a with high enantioselectivity, while the sense of asymmetric induction was opposite (eq. 1).²² Therefore, this method provides a unique example in which both enantiomers can be accessed by switching the geometry of a substrate while using a single enantiomer catalyst. In terms of the reactivity, an apparent deceleration of the reaction was observed with (E)-2a. Next, we conducted a reaction by using N-methyl encarbamate 13 (eq. 2). It turned out to be completely unreactive, implying the importance of the hydrogenbonding between the N-H hydrogen of the enecarbamate and the catalyst to facilitate the reaction. Finally, the mechanism of the transfer of the methoxy group was examined since not only the intermolecular pathway as we initially proposed in Figure 1, but the intramolecular pathway is also conceivable. A crossover experiment using a 1:1 mixture of the dimethyl ketal **3a** and diethyl ketal **14** revealed no formation of the crossover products (eq. 3), thereby confirming the unique intramolecularity of the process.

Scheme 2. Exploration of the Reaction Mechanism



From these observations, we postulated the reaction mechanism using (*Z*)- and (*E*)-enecarbamates as follows (Figure 2). In both reactions, the first step is the highly diastereo- and enantioselective C-C bond formation between two substrates hydrogen bonded by the carboxylic acid.^{9d,10} It is assumed that the catalyst recognizes the identical prochiral face of the quinone imine ketal,²³ and the diastereomeric intermediates I and I' are generated depending on the geometry of the enecarbamate.²⁴ The steric repulsion between the methyl group of the (*E*)-enecarbamate and the methoxy group of the quinone imine ketal may explain its lower reactivity. From each intermediate, the intramolecular methoxy group transfer occurs in the opposite sense of diastereoselectivity while minimizing the 1,3-allylic strain. As a consequence of two diastereodivergent processes, the same diastereomers with the opposite sense of enantioselectivity are finally obtained.²⁵

Figure 2. Plausible reaction mechanism.



In summary, we have developed the first asymmetric transformation using quinone imine ketals with enecarbamates to give chiral α -amino- β -aryl ethers, by shedding light on the reactivity of quinone imine ketals in chiral acid catalysis. Transformations of these products were examined to secure their use as a synthetic

60

handle to deliver a variety of enantiomerically enriched molecules. Further research to use quinone imine ketals in other asymmetric transformations is currently underway in this laboratory.

Supporting Information

Experimental details and characterization data. This material is available free of charge via the Internet at http://pubs.acs.org.

AUTHOR INFORMATION

Corresponding Author

maruoka@kuchem.kyoto-u.ac.jp

ACKNOWLEDGMENT

This work was partially supported by a Grant-in-Aid for Scientific Research from the MEXT (Japan). H.N. thank a Grant-in-Aid for the Research Fellowship of JSPS for Young Scientists.

REFERENCES

(1) For general reviews, see: (a) Nicolaou, K. C.; Snyder, S. A.; Montagnon, T.; Vassilikogiannakis, G. Angew., Chem. Int. Ed. 2002, 41, 1668. (b) Magdziak, D.; Meek, S. J.; Pettus, T. R. R. Chem. Rev. 2004, 104, 1383. (c) Roche, S. P.; Porco Jr, J. A. Angew., Chem. Int. Ed. 2011, 50, 4068. (d) Zhuo, C. X.; Zhang, W.; You, S. L. Angew., Chem. Int. Ed. 2012, 51, 12662

(2) For selected examples using quinones and quinone ketals, see: (a) Alemán, J.; Richter, B.; Jørgensen, K. A. Angew. Chem., Int. Ed. 2007, 46, 5515. (b) Angew. Chem., Int. Ed. 2007, 46, 5520. (c) Aleman, J.; Jacobsen, C. B.; Frisch, K.; Overgaard, J.; Jørgensen, K. A. Chem. Commun. 2008, 632. (d) Mikami, K.; Motoyama, Y.; Terada, M. J. Am. Chem. Soc. 1994, 116, 2812. (e) Evans, D. A.; Wu, J. J. Am. Chem. Soc. 2003, 125, 10162. (f) Ryu, D. H.; Zhou, G.; Corey, E. J. J. Am. Chem. Soc. 2004, 126, 4800. (g) Jarvo, E. R.; Lawrence, B. M.; Jacobsen, E. N. Angew. Chem., Int. Ed. 2005, 44, 6043. (h) Wang, C.; Chen, X.-H.; Zhou, S.-M.; Gong, L.-Z. Chem. Commun. 2010, 46, 1275. (i) Han, Z.-Y.; Chen, D.-F.; Wang, Y.-Y.; Guo, R.; Wang, P.-S.; Wang, C.; Gong, L.-Z. J. Am. Chem. Soc. 2012, 134, 6532.

(3) For selected examples using cyclohexadienones, see: (a) Liu, Q.; Rovis, T. J. Am. Chem. Soc. 2006, 128, 2552. (b) Rubush, D. M.; Morges, M. A.; Rose, B. J.; Thamm, D. H.; Rovis, T. J. Am. Chem. Soc. 2012, 134, 13554. (c) Vo, N. T.; Pace, R. D. M.; O'Hara, F.; Gaunt, M. J. J. Am. Chem. Soc. 2008, 130, 404. (d) Gu, Q.; You, S. L. Chem. Sci. 2011, 2, 1519. (e) Gu, Q.; You, S. L. Org. Lett. 2011, 13, 5192. Gu, Q.; Rong, Z.-Q.; Zheng, C.; You, S.-L. J. Am. Chem. Soc. 2010, 132, 4056. (f) Takizawa, S.; Nguyen, T. M.-N.; Grossmann, A.; Enders, D.; Sasai, H. Angew. Chem., Int. Ed. 2012, 51, 5423. (g) Wu, W.; Li, X.; Huang, H.; Yuan, X.; Lu, J.; Zhu, K.; Ye, J. Angew. Chem., Int. Ed. 2013, 52, 1743. (h) Corbett, M. T.; Johnson, J. S. Chem. Sci. 2013, 4, 2828.

(4) (a) Carreño, M. C.; Ribagorda, M. J. Org. Chem. 2000, 65, 1231. (b) Bodipati, N.; Peddinti, R. K. Org. Biomol. Chem., 2012, 10, 4549.

(5) For selected applications in total synthesis, see: (a) Chou, C. T.; Swenton, J. S. J. Am. Chem. Soc. 1987, 109, 6898. (b) Shair, M. D.; Yoon, T. Y.; Mosny, K. K.; Chou, T. C.; Danishefsky, S. J. J. Am. Chem. Soc. 1996, 118, 9509. (c) Myers, A. G.; Tom, N. J.; Fraley, M. E.; Cohen, S. B.; Madar, D. J. J. Am. Chem. Soc. 1997, 119, 6072. (d) Tohma, H.; Harayama, Y.; Hashizume, M.; Iwata, M.; Kiyono, Y.; Egi, M.; Kita, Y. J. Am. Chem. Soc. 2003, 125, 11235. (e) Jackson, S. K.; Banfield, S. C.; Kerr, M. A. Org. Lett. 2005, 7, 1215. (f) Ganton, M. D.; Kerr, M. A. J. Org. Chem. 2007, 72, 574. (g) Lebold, T. P.; Kerr, M. A. Org. Lett. 2007, 9, 1883. (h) Leduc, A. B.; Kerr, M. A. Eur. J. Org. Chem. 2007, 237. (i) Lebold, T. P.; Kerr, M. A. Org. Lett. 2008, 10, 997. (j) T. Chuang, K. V.; Navarro, R.; Reisman, S. E. Angew. Chem., Int. Ed. 2011, 50, 9447. (j) T. Chuang, K. V.; Navarro, R.; Reisman, S. E. Chem. Sci., 2011, 2, 1086. (k) Navarro, R.; Reisman, S. E. Org. Lett. 2012, 14, 4354.

(6) For a review on the use of quinone imines, see: Nair, V.; Dhanya, R.; Rajesh, C.; Devipriya, S. Synlett 2005, 2407.

56 (7) For other representative developments, see: (a) Chen, C.-P.; Chou, C.-

57

T.; Swenton, J. S. J. Am. Chem. Soc. **1987**, 109, 946. (b) Swenton, J. S.; Bonke, B. R.; Chen, C.-P.; Chou, C.-T. J. Org. Chem. **1989**, 54, 51. (c)

58 Swenton, J. S.; Bonke, B. R.; Clark, W. M.; Chen, C.-P.; Martin, K. V. J. 59

Org. Chem. 1990, 55, 2027. (d) Banfield, S. C.; England, D. B.; Kerr, M. A. Org. Lett. 2001, 3, 3325. (e) Banfield, S. C.; Kerr, M. A. Can. J. Chem. 2004, 82, 131.

(8) Jensen, K. L.; Franke, P. T.; Nielsen, L. T.; Daasbjerg, K.; Jørgensen, K. A. Angew., Chem. Int. Ed. 2010, 49, 129

(9) (a) Hashimoto, T.; Maruoka, K. J. Am. Chem. Soc. 2007, 129, 10054. (b) Hashimoto, T.; Uchiyama, N.; Maruoka, K. J. Am. Chem. Soc. 2008,

130, 14380. (c) Hashimoto, T.; Kimura, H.; Kawamata, Y.; Maruoka, K. Nat. Chem. 2011, 3, 642. (d) Hashimoto, T.; Kimura, H.; Nakatsu, H.; Maruoka, K. J. Org. Chem. 2011, 76, 6030.

(10) For general reviews on chiral Brønsted acid catalysis, see: (a) Akiyama, T. Chem. Rev. 2007, 107, 5744. (b) Terada, M. Synthesis 2010, 1929.

(11) For a review, see: Matsubara, R.; Kobayashi, S. Acc. Chem. Res. 2008, 41, 292.

(12) For selected examples on the use of enecarbamates and enamides as nucleophile in chiral Brønsted acid catalysis, see: (a) Terada, M.; Machioka, K.; Sorimachi, K. Angew., Chem. Int. Ed. 2006, 45, 2254. (b) Terada, M.; Machioka, K.; Sorimachi, K. J. Am. Chem. Soc. 2007, 129, 10336. (c) Terada, M.; Soga, N.; Momiyama, N. Angew. Chem., Int. Ed. 2008, 47, 4122. (d) Terada, M.; Machioka, K.; Sorimachi, K. Angew. Chem., Int. Ed. 2009, 48, 2553. (e) Liu, H.; Dagousset, G.; Masson, G.; Retailleau, P.; Zhu, J. J. Am. Chem. Soc. 2009, 131, 4598. (f) Dagousset, G.; Drouet, F.; Masson, G.; Zhu, J. Org. Lett. 2009, 11, 5546. (g) Dagousset, G.; Zhu, J.; Masson, G. J. Am. Chem. Soc. 2011, 133, 14804. (h) Alix, A.; Claudia, L.; Retailleau, P.; Masson, G. J. Am. Chem. Soc. 2012, 134, 10389. (i) Lu, M.; Lu, Y.; Zhu, D.; Zeng, X.; Li, X.; Zhong, G. Angew. Chem., Int. Ed. 2010, 49, 8588.

(13) For the chiral Brønsted acid catalyzed enantioselective synthesis of hemiaminal ethers, see: (a) Li, G.; Fronczek, F. R.; Antilla, J. C. J. Am. Chem. Soc. 2008, 130, 12216. (b) Vellalath, S.; Čorić, I.; List, B. Angew. Chem., Int. Ed. 2010, 49, 9749.

(14) Honjo, T.; Phipps, R. J.; Rauniyar, V.; Toste, F. D. Angew. Chem., Int. Ed. 2012, 51, 9684.

(15) (a) Pansare, S. V.; Vederas, J. C. J. Org. Chem. 1989, 54, 2311. (b) Mecozzi, T.; Petrini, M. Synlett, 2000, 73.

(16) For the use of stronger chiral Brønsted acids, see SI.

(17) For example, use of 2-methoxy quinone imine ketal 5 gave rise to 2amino-3-methylindoline 6 as the only isolable compound with low enantioselectivity



(18) Avdeenko, A. P.; Konovalova, S. A.; Mikhailichenko, O. N.; Yusia, A. L.; Santalova, A. A.; Palamarchyk, G. V.; Zubatyuk, R. I.; Shishkin, O. V.; Burmistrov, K. S. Russ. J. Org. Chem., 2011, 47, 1169.

(19) (a) Dalidowicz, P.; Swenton, J. S. J. Org. Chem. 1993, 58, 4802. (b) Engler, T. A.; Scheibe, C. M. J. Org. Chem. 1998, 63, 6247. (c) Hu, Y.; Kamitanaka, T.; Mishima, Y.; Dohi, T.; Kita, Y. J. Org. Chem. 2013, 78, 5530

(20) Terada, M.; Sorimachi, K. J. Am. Chem. Soc. 2007, 129, 292.

(21) The diastereoselectivity was essentially governed by the substrate, and the use of a weaker acid resulted in a cleaner conversion.

(22) Chang, L.; Kuang, Y.; Qin, B.; Zhou, X.; Liu, X.; Lin, L.; Feng, X. Org. Lett. 2010, 12, 2214.

(23) VT NMR experiments revealed facile interconversion of the C=N double bond of the quinone imine ketal.

(24) (a) Matsubara, R.; Nakamura, Y.; Kobayashi, S. Angew., Chem. Int. Ed. 2004, 43, 3258. (b) Matsubara, R.; Kawai, N.; Kobayashi, S. Angew., Chem. Int. Ed. 2006, 45, 3814.

(25) For the discussion on the transition state model, see SI.

