

# CHEMISTRY

## A European Journal

A Journal of



### Accepted Article

**Title:** Selective inhibitors of a human prolyl hydroxylase (OGFOD1) involved in ribosomal decoding

**Authors:** Cyrille C Thinnes, Christopher T Lohans, Martine I Abboud, Tzu-Lan Yeh, Anthony Tumber, Radosław P Nowak, Martin Attwood, Matthew E Cockman, Udo Oppermann, Christoph Loenarz, and Christopher Schofield

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

**To be cited as:** *Chem. Eur. J.* 10.1002/chem.201804790

**Link to VoR:** <http://dx.doi.org/10.1002/chem.201804790>

Supported by  
**ACES**

WILEY-VCH

## FULL PAPER

# Selective inhibitors of a human prolyl hydroxylase (OGFOD1) involved in ribosomal decoding


Cyrille C. Thinnès,<sup>[a]</sup> Christopher T. Lohans,<sup>[a]</sup> Martine I. Abboud,<sup>[a]</sup> Tzu-Lan Yeh,<sup>[a]</sup> Anthony Tumber,<sup>[a,b]</sup> Radosław P. Nowak,<sup>[b]</sup> Martin Attwood,<sup>[c]</sup> Matthew E. Cockman,<sup>[c]</sup> Udo Oppermann,<sup>[b]</sup> Christoph Loenarz<sup>[a]</sup> and Christopher J. Schofield\*<sup>[a]</sup>

**Abstract:** Human prolyl hydroxylases are involved in the modification of transcription factors, procollagen, and ribosomal proteins, and are current medicinal chemistry targets. To date, there are few reports on inhibitors selective for the different types of prolyl hydroxylases. Here we report a structurally informed template-based strategy for the development of inhibitors selective for the human ribosomal prolyl hydroxylase OGFOD1. These inhibitors did not target the other human oxygenases tested, including the structurally-similar hypoxia-inducible transcription factor prolyl hydroxylase, PHD2.

## Introduction

In humans and other animals, prolyl hydroxylases (PHs) play critical roles in collagen biosynthesis and hypoxia sensing.<sup>[1–3]</sup> The PHs are Fe(II) and 2-oxoglutarate (2OG) dependent oxygenases, which normally produce succinate and CO<sub>2</sub> as coproducts.<sup>[4]</sup> The procollagen PHs (CP3H and CP4H) hydroxylate either C3 or C4 of prolyl residues; the latter is essential for maintenance of the collagen triple helix secondary structure (Figure 1A).<sup>[5,6]</sup> In humans, the prolyl hydroxylase domain enzymes (PHD1, 2, and 3) act as oxygen sensors in the chronic response to hypoxia by catalysing oxygen-limited hydroxylation of prolyl residues in the hypoxia-inducible factor- $\alpha$  (HIF $\alpha$ ) subunits of HIF transcription factors, leading to HIF $\alpha$  degradation in aerobic conditions (Figure 1A).<sup>[1]</sup>

The human 2OG oxygenase OGFOD1 has been recently shown to hydroxylate Pro-62 of the ribosomal protein RPS23.<sup>[7–10]</sup> Pro-62<sub>RPS23</sub> is situated in the ribosomal decoding site, which is responsible for ensuring the fidelity of mRNA codon recognition by tRNA and release factor proteins during protein synthesis.<sup>[11,12]</sup> While the role of this hydroxylation in human and animal cells is not yet understood, the *Saccharomyces cerevisiae* OGFOD1 homologue Tpa1p is proposed to catalyze dihydroxylation of the corresponding prolyl residue, and to regulate translational accuracy in an mRNA sequence context-dependent manner.<sup>[8,13,14]</sup>



**Figure 1. Prolyl hydroxylase reactions and structures.** (A) Regio- and stereoselectivity of hydroxylations catalyzed by different types of prolyl hydroxylases. Each hydroxylation is coupled to the conversion of 2-oxoglutarate (2OG) and O<sub>2</sub> into succinate and CO<sub>2</sub>. OGFOD1 acts on ribosomal proteins, the CPs act on procollagen, and the PHDs act on the hypoxia inducible factor (HIF) transcription factors.<sup>[5]</sup> (B) Structures of 2OG and the 2OG analogue N-oxalylglycine (NOG). (C, D) Views from the crystallographically observed active sites of OGFOD1 (PDB 4NHX)<sup>[10]</sup> and PHD2 (PDB 5L9R)<sup>[15]</sup> showing the interactions between active site residues, the bound metal [Mn(II)] substituting for Fe(II), and NOG.

Of the ~60–70 human 2OG oxygenases, some are current medicinal chemistry targets, including enzymes involved in chromatin modification and lipid metabolism.<sup>[5,16]</sup> Inhibition of the procollagen hydroxylases is under consideration as a target to limit the overproduction of collagen associated with certain cancers and fibrotic diseases.<sup>[17]</sup> The PHDs are presently being targeted for the treatment of hypoxia-related diseases, with inhibitors in late-stage clinical trials for anaemia.<sup>[18]</sup> If OGFOD1 is indeed involved in mRNA codon recognition, as suggested based on studies of yeast homologues,<sup>[8,13]</sup> small molecule-mediated inhibition of ribosomal hydroxylation could prove useful for the treatment of diseases such as muscular dystrophy that are caused by premature stop-codons through nonsense suppression.<sup>[19]</sup> However, due to the uncertainty regarding the specific roles of OGFOD1 and OGFOD1-catalysed hydroxylation in animals, it is unclear how exactly its inhibition might manifest. Thus, such OGFOD1 inhibitors are also of interest as chemical probes to decipher the biological role of RPS23 hydroxylation, as

[a] Dr. C. C. Thinnès, Dr. C. T. Lohans, Dr. M. I. Abboud, Dr. T.-L. Yeh, Dr. A. Tumber, Prof. Dr. C. Loenarz, Prof. Dr. C. J. Schofield  
Department of Chemistry, University of Oxford  
Oxford, OX1 3TA (UK)

[b] Dr. A. Tumber, Dr. R. P. Nowak, Prof. Dr. U. Oppermann  
Structural Genomics Consortium, University of Oxford  
Headington, OX3 7DQ (UK)

[c] Dr. M. Attwood, Dr. M. E. Cockman  
Centre for Cellular and Molecular Physiology, University of Oxford  
Oxford, OX3 7BN (UK)

Supporting information for this article is given via a link at the end of the document.


## FULL PAPER

well as those of other recently reported ribosome-associated hydroxylations.<sup>[20–23]</sup>

As the 2OG oxygenases are involved in diverse biological processes, developing inhibitors selective for a particular oxygenase is an important therapeutic consideration. The available biophysical evidence, principally from crystallography, implies that key features in the active sites of the different types of human PHs are substantially, but not completely, conserved.<sup>[4]</sup> Therefore, there is the potential that inhibitors targeting OGFOD1 could also interfere with hypoxia sensing and collagen biosynthesis through inhibition of other PHs (or vice versa). To date, no detailed evidence for inhibitors selective for the different types of human PHs have been reported. Here, we establish the viability of a structure-guided template-based approach for the development of selective OGFOD1 inhibitors which do not target the other human oxygenases tested, including the human hypoxia-sensing enzyme PHD2.

## Results

In order to assess the viability of developing inhibitors selective for particular PHs, with a focus on OGFOD1, we first compared crystal structures of OGFOD1 and PHD2.<sup>[10,24,25]</sup> Although there are differences in the OGFOD1 and PHD2 active sites, the binding modes of 2OG [and 2OG analogue *N*-oxalylglycine (NOG)] are conserved (Figure 1B–1D). The 2-oxoacid group of 2OG (or NOG) binds to the metal in a bidentate manner, while the 2OG C-5 carboxylate is positioned to interact with conserved tyrosine and arginine residues (Tyr169 and Arg230 in OGFOD1, Tyr329 and Arg383 in PHD2; Figure 1C, 1D). These comparisons, along with those shown for other human oxygenases,<sup>[4]</sup> suggest that inadvertent inhibition of the PHDs may represent a challenge in developing selective OGFOD1 inhibitors, and vice versa.



**Figure 2. Oxygenase inhibitors and their binding modes.** (A) Structures of inhibitors of 4-hydroxyphenylpyruvate dioxygenase (HPPD) and PHD2. Like OGFOD1 and PHD2, HPPD is an Fe(II)-dependent 2-oxoacid oxygenase, but from a different structural family. (B) Comparison of the common chelation motifs of sulcotrione **2**, GSK1278863 **6**, and FG2216 **7**. The structures of sulcotrione **2** and GSK1278863 **6** were modelled onto the crystallographically observed structure of PHD2 with bound FG2216 **7** (PDB 3HQJ).<sup>[25]</sup> A larger version is shown in Figure S5. (C) *In silico* binding model of GSK1278863 **6** with OGFOD1 (PDB 4NHX)<sup>[10]</sup> and PHD2 (PDB 5L9R).<sup>[15]</sup> The tricarbonyl is expected to interact with the bound metal, while the *N*-cyclohexyl groups likely occupy the substrate binding position.

Given the conserved elements of the OGFOD1 and PHD2 active sites,<sup>[10]</sup> and evidence for induced fit and conformational movements in prolyl hydroxylase catalysis,<sup>[15]</sup> we contemplated a structurally informed template-derivatization approach for developing OGFOD1 inhibitors.<sup>[26]</sup> We first considered the 4-hydroxyphenylpyruvate dioxygenase (HPPD) inhibitor nitisinone **1** (which is clinically used for the treatment of tyrosinaemia)<sup>[27]</sup> and related plant growth inhibitors sulcotrione **2**, mesotrine **3**, prohexadione-calcium **4**, and trinexapac-ethyl **5** (Figure 2A).<sup>[28]</sup> These compounds are related to the ‘tricarbonyl’ chemotype found in the PHD inhibitor GSK1278863 **6**, which is currently in clinical trials for the treatment of anaemia.<sup>[18]</sup> In addition to GSK1278863 **6**, we tested PHD inhibitors FG2216 **7**, FG4592 **8**, and IOX2 **9**.<sup>[29,30]</sup>

These inhibitors were screened for binding to OGFOD1 and PHD2 by differential scanning fluorimetry (DSF) and for inhibition using matrix-assisted laser desorption ionisation time-of-flight mass spectrometry (MALDI-TOF MS) (Figure S1–S4).<sup>[31,32]</sup> As expected based on previous studies,<sup>[33]</sup> PHD inhibitors FG2216 **7**, FG4592 **8**, and IOX2 **9** inhibited OGFOD1 (Figure S1). Notably, triketone-based HPPD and plant growth inhibitors **1–5** (Figure 2A),


## FULL PAPER

related to GSK1278863 **6**, displayed moderate inhibition of OGFOD1, while not inhibiting the activity of PHD2 within our limits of detection (Figure S2). Additionally, these inhibitors increased the apparent thermal stability of OGFOD1, as observed by DSF (Figure S3, S4). The inhibition of OGFOD1 by prohexadione-calcium **4** and trinexapac-ethyl **5** (Figure S2) suggests that further investigation of these agrochemicals is warranted, given their potential interactions with other human 2OG oxygenases.

GSK1278863 **6** and sulcotrione **2** were modelled into the crystallographically-observed OGFOD1 and PHD2 active sites, with consideration for potential metal-chelating properties, salt bridge interactions, and crystallographic studies of PHD2 with inhibitors (such as FG2216 **7**; PDB 3HQY) (Figure 2B, S5).<sup>[25]</sup> These analyses imply that both **6** and **2** will engage in Fe(II) chelation *via* an enolate form of their 1,3-diketone motif (Figure S5). Similarly to the C-5 carboxylate of 2OG/NOG,<sup>[10]</sup> and the carboxylate of FG2216 **7**,<sup>[25]</sup> the GSK1278863 **6** carboxylate is predicted to interact with Tyr169 and Arg230 of OGFOD1. The OGFOD1:**6** model suggests that the inhibitor ring systems bind in two approximately perpendicular planes, one comprising the diketone ring and glycinamide side chain, and the other formed from the cyclohexyl rings (Figure 2C). The model also implies that the cyclohexyl rings engage differently with the OGFOD1 and PHD2 active sites (Figure 2C). By contrast, compounds based on an isoquinoline chemotype (e.g., FG2216 **7**) are predicted to bind in a more co-planar manner<sup>[25]</sup> and were considered less likely to readily lead to OGFOD1-selective inhibitors.

While many reported PHD2 inhibitors possess a glycinamide side chain, the triketone plant growth/HPPD inhibitors do not (Figure 2A). Modelling results suggest that the sulcotrione **2** methyl sulfonyl group and the nitro group of mesotrione **3** and nitisinone **1** may mimic the 2OG/glycinamide side chain binding at their active sites (Figure 2B). It is possible that the enzyme active site may accommodate the side chains of nitisinone **1**, sulcotrione **2**, and mesotrione **3**, which are bulkier than the glycinamide side chain of FG2216 **7**; however, in the absence of structural information, we cannot preclude the possibility that these inhibitors with bulky side chains interact with the enzyme in an alternate orientation.

Both the barbiturate-based PHD inhibitors (e.g., **6**) and the HPPD/plant growth inhibitors (e.g., **1-5**) have a tri-carbonyl motif. However, triketones **1-5** are likely more conformationally flexible than the barbiturates. Notably, the DSF results suggest that the triketones stabilize OGFOD1 more than PHD2 (Figure S3), whereas the glycinamide-containing PHD2 inhibitors (e.g., **6-9**) stabilize PHD2 more than OGFOD1 (Figure S4). We thus explored whether modification of the barbiturate/cyclohexane-1,3-dione-based ring scaffolds could be exploited in the development of selective OGFOD1 inhibitors.




**Figure 3. Fragment-based screening approach for OGFOD1 inhibition.** (A) Structures of diketones, triketones, and structurally related compounds used for fragment-based screening. (B) Inhibitory effect of the fragments on the hydroxylation activity of OGFOD1 (1 μM). The plotted data represent the mean percentage inhibition for the experiment performed in triplicate, while the error bars indicate the standard deviation.

Diketones (e.g., diethyl malonate **10** and 2-acetylcyclohexanone **11**), triketones (e.g., triacetylmethane **12** and substituted 1,3-cyclohexanediones **13-15**), and structurally-related compounds (e.g., 2',6'-dihydroxyacetophenone **16** and barbituric acid **17**) were screened for OGFOD1 inhibition (Figure 3). The preliminary results suggested that the degree of OGFOD1 inhibition may in part relate to the propensity for inhibitor enolization. Thus, while the diketones tested (i.e., **10**, **11**) were poor inhibitors, the triketones (e.g., **12-15**) were more potent

## FULL PAPER


(Figure 3B). However, the phenolic triketone 'mimic' 2',6'-dihydroxyacetophenone **16** was a relatively poor OGFOD1 inhibitor. The most potent template identified for OGFOD1 inhibition was manifested in the C-5 substituted barbiturate derivatives (**18-20**); such compounds are readily enolized, which is likely beneficial for metal chelation. *N*-Methylation of the barbiturate ring (e.g., **20**) improved potency, as did the introduction of an acyl substituent on C-5 of the barbiturate core (e.g., **18** compared to **17**). Replacement of the C-2 oxygen with sulfur (thiobarbituric acid **19**) did not substantially increase the potency relative to the oxygen analogue (i.e., **18**), while replacement of the oxygen at C-4 with a nitrogen (e.g., **21**) abolished any inhibitory activity (Figure 3B). We therefore focused on obtaining selective OGFOD1 inhibitors by modifying the C-5 position of a 1,3-dimethyl barbiturate core. Notably, compounds **22** and **23**, which combine a barbiturate core with a glycnamide ethyl ester side chain, did not manifest selectivity for OGFOD1 over PHD2 (Figure 4; Figure S6).



**Figure 4. Optimization studies of barbiturate inhibitors.** (A) Structures of the barbiturate glycinamide ethyl esters tested against OGFOD1 and PHD2. (B) Panel of phenyl-substituted *N,N*-dimethylbarbiturates, demonstrating the impact of increasing acyl chain length on OGFOD1 inhibition. (C) Panel of barbiturates synthesized with different *N*-alkyl substituents. (D) Panel of C-5 substituted *N,N*-dimethylbarbiturates based on lead compound **CCT3** (**26**).

Therefore, **24** was prepared, in which structural features of **22** are 'merged' with HPPD/plant growth inhibitors **1-3**, by introducing a C-5 aromatic acyl substituent in place of a glycnamide. To investigate the optimal spacing between the barbiturate and the aromatic side chain, the initial screen encompassed a 1,3-dimethyl barbiturate core bearing acetyl **20**, benzoyl **24**, phenylacetyl **25**, and hydrocinnamoyl **26** (or **CCT3**) substituents. The resulting aromatic compounds potently inhibited OGFOD1, with an increase in potency being observed upon extending the carbon chain length from **24** ( $IC_{50} = 30 \mu M$ ) to **25** ( $IC_{50} = 2 \mu M$ ), and from **25** to **26** (**CCT3**) ( $IC_{50} = 0.7 \mu M$ ; assays performed using  $1 \mu M$  OGFOD1; Figure S7). Extending the side chain further (e.g., **27**, **28**) did not provide a substantial increase in potency. Importantly, this series displayed no observable PHD2 inhibition at concentrations up to  $100 \mu M$  (Figure S8). These results suggest that the C-5 substituent is important for obtaining selectivity between OGFOD1 and PHD2.

The importance of the *N*-alkyl substituents on inhibition was examined with barbiturates bearing C-5 cyclopentyl substituents (Figure 4C). Analogues bearing *N,N*-dimethyl (**29**), *N,N*-diethyl (**30**), and *N,N*-dicyclohexyl (**31**) groups were prepared (Figure 4C); these analogues were less potent, suggesting that substituents larger than methyl groups may not be favourable for OGFOD1 inhibition (Figure S9). Note, however, that GSK1278863 **6** potently inhibits OGFOD1 despite the presence of *N,N*-dicyclohexyl groups (Figure 2A).<sup>[33]</sup>



**Figure 5. Selectivity of lead compounds against 2OG oxygenases.** (A) Comparison of inhibition data for compounds **CCT3** and **CCT4** against OGFOD1 and other human 2OG oxygenases. NI: no inhibition observed. (B) Western blot (antibody specific for HIF $\alpha$ ) showing the lack of impact of **CCT3** and **CCT4** on the activity of PHD enzymes in HeLa cells.<sup>[33]</sup> By comparison, the known PHD inhibitor IOX4 inhibits the hydroxylation of HIF $\alpha$ , preventing proteasomal degradation.<sup>[34]</sup> (C) Extent of displacement of 2OG from the active

## FULL PAPER

site of OGFOD1 (blue) and PHD2 (black) by **CCT3** and **CCT4** as observed by CPMG-edited <sup>1</sup>H-NMR. See Figure S15 for additional details.

Modelling suggests that the aryl side chains of **25** and **CCT3** likely do not fit in the OGFOD1 2OG binding site due to steric constraints; instead, the aryl ring may bind in the substrate binding site, potentially contributing to the selectivity of these inhibitors. Varying the C-5 side chain with different mono- and bicyclic aromatic and saturated substituents (**32-45**) did not have a substantial impact on potency (Figure 4D, Figure S10). Similarly, changing the nature of the C-5 link from a ketone to an amide, and extending the carbon chain length beyond two carbons between the carbonyl and the substituent (i.e., **46**) had little effect.

On the basis of these SAR studies, in particular those examining the impact of the C-5 and barbiturate *N*-alkyl substituents, the sub-micromolar potency inhibitors **CCT3** (**26**) and **CCT4** (**42**) were selected for further characterization (Figure 4, S11). These two inhibitors were screened for inhibitory activity against a panel of human 2OG oxygenases (Figure 5A). Of the human enzymes screened, including the HIF prolyl hydroxylase PHD2, the ribosomal oxygenases MINA53 and NO66 (which hydroxylate histidyl residues in ribosomal proteins)<sup>[20]</sup> and the asparaginyl hydroxylase factor-inhibiting HIF (FIH), **CCT3** and **CCT4** only potently inhibited OGFOD1 (with IC<sub>50</sub> values of 0.73 μM and 0.69 μM, respectively). Additionally, these compounds showed poor, or no, inhibition of more distantly related 2OG oxygenases, such as the histone demethylases KDM4A, JARID1B, and JARID1C (Figure 5A). **CCT3** and **CCT4** did potently inhibit the structurally close yeast OGFOD1 homologues Tpa1p and Ofd1 (which hydroxylate RPS23 prolyl residues; Figure S12), consistent with the high levels of similarity between the homologous active sites.<sup>[10]</sup> It is also notable that the sensitivity of the current hydroxylation assay is a limitation for ranking the activities of the potent inhibitors, as the IC<sub>50</sub> values obtained are close to the OGFOD1 concentration used (i.e., the minimum IC<sub>50</sub> value that can be measured is 0.5 μM); thus, these inhibitors may be more potent than represented by the currently reported IC<sub>50</sub> values.

To validate the *in vitro* inhibition results, direct binding interactions between inhibitors **CCT3** and **CCT4** and OGFOD1 and PHD2 were assessed by NMR. While both inhibitors were observed to strongly bind to OGFOD1, as observed using <sup>1</sup>H Carr-Purcell-Meiboom-Gill (CPMG) analyses (Figure S13), only weak binding to PHD2 was observed by water-Ligand Observe Gradient Spectroscopy (wLOGSY) experiments (Figure S14). Competition experiments between the inhibitors and enzyme-bound 2OG were conducted by monitoring the recovery of the enzyme-free 2OG methylene peak at 2.35 ppm using CPMG-edited <sup>1</sup>H-NMR upon addition of the inhibitor.<sup>[35]</sup> The results indicate that **CCT3** and **CCT4** are capable of displacing bound 2OG from the active site of OGFOD1, but not from that of PHD2 (Figure 5C, S15).

We examined the potential inhibition of PHD enzymes by **CCT3** and **CCT4** in the HeLa human cell line.<sup>[36]</sup> As compared to the known PHD inhibitor IOX4,<sup>[34]</sup> **CCT3** and **CCT4** did not stabilize HIFα (Figure 5B). PHD-catalyzed hydroxylation targets HIFα for proteasomal degradation, indicating that these

compounds do not inhibit the PHDs in cells. Based on a MDR1-MDCK assay (performed by Cyprotex, UK; see the Supporting Information), **CCT3** and **CCT4** demonstrate good cell permeability properties, and are predicted to be permeable to the blood brain barrier. Liver microsome stability studies indicate only low levels of clearance of **CCT3** and **CCT4** (Cyprotex, UK).

## Discussion

Our results demonstrate the viability of a template-based approach for the development of selective 2OG oxygenase/prolyl hydroxylase inhibitors capable of differentiating between closely related active sites, such as those of the human prolyl hydroxylases OGFOD1 and PHD2. Furthermore, the optimized inhibitors also did not inhibit the other 2OG oxygenases tested, including other ribosomal oxygenases, as well as histone demethylases. The results suggest that specific inhibitor 'templates' may be preferred for certain oxygenases or oxygenase subfamilies, as supported by work implying differential selectivity between PH and JmjC histone demethylase inhibitors.<sup>[37]</sup> This preference for particular templates may even extend to enzymes with closely related active sites. Appropriately modified barbiturate-based inhibitors may selectively inhibit OGFOD1 because of their ability to support substituents which extend towards active site residues present in OGFOD1 but not in PHD2. By contrast, the glycinamide side chain present in many PHD2 inhibitors (including several compounds in clinical trials, e.g., **6** and **7**) is clearly not required for potent OGFOD1 inhibition. It should also be noted that potent PHD2 inhibitors without a glycinamide side chain are known.<sup>[33,34]</sup> In future work, it will be of interest to further explore the selectivity of the compounds reported here. In this regard, studies with the procollagen C-4 (and C-3) prolyl hydroxylases are of particular interest, especially as the procollagen C-4 PHs are potential therapeutic targets.<sup>[38]</sup>

There is considerable academic and pharmaceutical interest in developing chemical probe compounds to investigate the biological functions of 2OG oxygenases.<sup>[39]</sup> Our results suggest that the development of leads based on known pharmaceutical and agrochemical 'templates' (some of which can penetrate the blood-brain barrier) with well-studied physicochemical properties, such as barbiturates, will be a productive strategy. The combination of the tricarbonyl barbiturate template of the PHD2 inhibitor GSK1278863 **6** with the side chains of agrochemical oxygenase inhibitors (e.g., prohexadione-calcium), followed by subsequent optimization, yielded potent and selective OGFOD1 inhibitors. Future work can now be focused on applying these OGFOD1 inhibitors to investigate the biological roles of OGFOD1, and applying a similar inhibitor development strategy to other ribosomal oxygenases. Based on what is observed in these functional studies, further optimization of these inhibitors may be warranted (e.g., if penetration of the blood-brain barrier is desirable).

It is important to note that many PHD2 inhibitors reported in the literature, including those screened in this work, and those currently in clinical trials, also inhibit OGFOD1.<sup>[33]</sup> Indeed, they may also inhibit other human prolyl hydroxylases and 2OG

## FULL PAPER

oxygenases, including those for which assays are currently not available.<sup>[10,40]</sup> From a clinical perspective, it is also important to note that the barbiturate-related 'triketone' HPPD inhibitor nitisinone, which is used in the treatment of type I tyrosinaemia,<sup>[27]</sup> is an OGFOD1 inhibitor, something that might be taken into consideration if nitisinone successors with improved properties are pursued. In the present work, we have demonstrated that it is possible to attain selectivity between different 2OG oxygenases, with lead compounds that inhibit OGFOD1, but not PHD2. Such 'biochemical selectivity' is not necessarily an issue with clinical applications as the desired pharmacological effect/safety profile may be achieved by controlling metabolism and tissue distribution. However, we propose that, at least for chronic applications, biochemical selectivity could and should be optimized during the development of 2OG oxygenase inhibitors. We also hope that inhibitors selective for particular 2OG oxygenases may enable their biological roles to be deciphered.

## Acknowledgements

We thank the Wellcome Trust, Cancer Research UK, and the Biotechnology and Biological Sciences Research Council (BBSRC) for supporting our research on 2OG-dependent oxygenases. C.L. thanks the Leverhulme Trust for an Early Career Fellowship.

**Keywords:** barbiturate • inhibitors • medicinal chemistry • OGFOD1 • oxygenase

- [1] C. J. Schofield, P. J. Ratcliffe, *Nat. Rev. Mol. Cell Biol.* **2004**, *5*, 343-354.
- [2] S. E. Wilkins, M. I. Abboud, R. L. Hancock, C. J. Schofield, *ChemMedChem* **2016**, *11*, 773-786.
- [3] J. D. Webb, M. L. Coleman, C. W. Pugh, *Cell. Mol. Life Sci.* **2009**, *66*, 3539-3554.
- [4] W. S. Aik, R. Chowdhury, I. J. Clifton, R. J. Hopkinson, T. Leissing, M. A. McDonough, R. Nowak, C. J. Schofield, L. J. Walport, *Introduction to structural studies on 2-oxoglutarate-dependent oxygenases and related enzymes*, Royal Society of Chemistry, **2015**, p. 59.
- [5] C. Loenarz, C. J. Schofield, *Nat. Chem. Biol.* **2008**, *4*, 152-156.
- [6] K. L. Gorres, R. T. Raines, *Crit. Rev. Biochem. Mol. Biol.* **2010**, *45*, 106-124.
- [7] R. S. Singleton, P. Liu-Yi, F. Formenti, W. Ge, R. Sekirnik, R. Fischer, J. Adam, P. J. Pollard, A. Wolf, A. Thalhammer, C. Loenarz, E. Flashman, A. Yamamoto, M. L. Coleman, B. M. Kessler, P. Wappner, C. J. Schofield, P. J. Ratcliffe, M. E. Cockman, *Proc. Natl. Acad. Sci. U. S. A.* **2014**, *111*, 4031-4036.
- [8] C. Loenarz, R. Sekirnik, A. Thalhammer, W. Ge, E. Spivakovsky, M. M. Mackeen, M. A. McDonough, M. E. Cockman, B. M. Kessler, P. J. Ratcliffe, A. Wolf, C. J. Schofield, *Proc. Natl. Acad. Sci. U. S. A.* **2014**, *111*, 4019-4024.
- [9] M. J. Katz, J. M. Acevedo, C. Loenarz, D. Galagovsky, P. Liu-Yi, M. Pérez-Pepe, A. Thalhammer, R. Sekirnik, W. Ge, M. Melani, M. G. Thomas, S. Simonetta, G. L. Boccaccio, C. J. Schofield, M. E. Cockman, P. J. Ratcliffe, P. Wappner, *Proc. Natl. Acad. Sci. U. S. A.* **2014**, *111*, 4025-4030.
- [10] S. Horita, J. S. Scotti, C. Thinner, Y. S. Mottaghi-Taromsari, A. Thalhammer, W. Ge, W. Aik, C. Loenarz, C. J. Schofield, M. A. McDonough, *Structure* **2015**, *23*, 639-652.
- [11] T. M. Schmeing, V. Ramakrishnan, *Nature* **2009**, *461*, 1234-1242.
- [12] W. V. Gilbert, *Trends Biochem. Sci.* **2011**, *36*, 127-132.
- [13] M. V. Nesterchuk, P. V. Sergiev, O. A. Dontsova, *Acta Naturae* **2011**, *3*, 22-33.
- [14] L. E. Alksne, R. A. Anthony, S. W. Liebman, J. R. Warner, *Proc. Natl. Acad. Sci. U. S. A.* **1993**, *90*, 9538-9541.
- [15] R. Chowdhury, I. K. Leung, Y. M. Tian, M. I. Abboud, W. Ge, C. Domene, F. X. Cantrelle, I. Landrieu, A. P. Hardy, C. W. Pugh, P. J. Ratcliffe, T. D. Claridge, C. J. Schofield, *Nat. Commun.* **2016**, *7*, 12673.
- [16] N. R. Rose, M. A. McDonough, O. N. King, A. Kawamura, C. J. Schofield, *Chem. Soc. Rev.* **2011**, *40*, 4364-4397.
- [17] J. D. Vasta, K. A. Andersen, K. M. Deck, C. P. Nizzi, R. S. Eisenstein, R. T. Raines, *ACS Chem. Biol.* **2016**, *11*, 193-199.
- [18] M. C. Chan, J. P. Holt-Martyn, C. J. Schofield, P. J. Ratcliffe, *Mol. Aspects Med.* **2016**, *47-48*, 54-75.
- [19] K. M. Keeling, X. Xue, G. Gunn, D. M. Bedwell, *Annu. Rev. Genomics Hum. Genet.* **2014**, *15*, 371-394.
- [20] W. Ge, A. Wolf, T. Feng, C. H. Ho, R. Sekirnik, A. Zayer, N. Granatino, M. E. Cockman, C. Loenarz, N. D. Loik, A. P. Hardy, T. D. Claridge, R. B. Hamed, R. Chowdhury, L. Gong, C. V. Robinson, D. C. Trudgian, M. Jiang, M. M. Mackeen, J. S. McCullagh, Y. Gordiyenko, A. Thalhammer, A. Yamamoto, M. Yang, P. Liu-Yi, Z. Zhang, M. Schmidt-Zachmann, B. M. Kessler, P. J. Ratcliffe, G. M. Preston, M. L. Coleman, C. J. Schofield, *Nat. Chem. Biol.* **2012**, *8*, 960-962.
- [21] R. Chowdhury, R. Sekirnik, N. C. Brissett, T. Krojer, C. H. Ho, S. S. Ng, I. J. Clifton, W. Ge, N. J. Kershaw, G. C. Fox, J. R. C. Muniz, M. Vollmar, C. Phillips, E. S. Pilka, K. L. Kavanagh, F. von Delft, U. Oppermann, M. A. McDonough, A. J. Doherty, C. J. Schofield, *Nature* **2014**, *510*, 422-426.
- [22] S. Markolovic, Q. Zhuang, S. E. Wilkins, C. D. Eaton, M. I. Abboud, M. J. Katz, H. E. McNeil, R. K. Leśniak, C. Hall, W. B. Struwe, R. Konietzny, S. Davis, M. Yang, W. Ge, J. L. P. Benesch, B. M. Kessler, P. J. Ratcliffe, M. E. Cockman, R. Fischer, P. Wappner, R. Chowdhury, M. L. Coleman, C. J. Schofield, *Nat Chem Biol* **2018**, *14*, 688-695.
- [23] T. Feng, A. Yamamoto, S. E. Wilkins, E. Sokolova, L. A. Yates, M. Münzel, P. Singh, R. J. Hopkinson, R. Fischer, M. E. Cockman, J. Shelley, D. C. Trudgian, J. Schödel, J. S. McCullagh, W. Ge, B. M. Kessler, R. J. Gilbert, L. Y. Frolova, E. Alkalaeva, P. J. Ratcliffe, C. J. Schofield, M. L. Coleman, *Mol Cell* **2014**, *53*, 645-654.
- [24] M. A. McDonough, C. Loenarz, R. Chowdhury, I. J. Clifton, C. J. Schofield, *Curr. Opin. Struct. Biol.* **2010**, *20*, 659-672.
- [25] R. Chowdhury, M. A. McDonough, J. Mecinović, C. Loenarz, E. Flashman, K. S. Hewitson, C. Domene, C. J. Schofield, *Structure* **2009**, *17*, 981-989.
- [26] C. C. Thinner, A. Tumber, C. Yapp, G. Scozzafava, T. Yeh, M. C. Chan, T. A. Tran, K. Hsu, H. Tarhonskaya, L. J. Walport, S. E. Wilkins, E. D. Martinez, S. Müller, C. W. Pugh, P. J. Ratcliffe, P. E. Brennan, A. Kawamura, C. J. Schofield, *Chem. Commun.* **2015**, *51*, 15458-15461.
- [27] P. J. McKiernan, *Drugs* **2006**, *66*, 743-750.
- [28] G. R. Moran, *Arch Biochem Biophys* **2005**, *433*, 117-128.
- [29] M. J. Koury, V. H. Haase, *Nat. Rev. Nephrol.* **2015**, *11*, 394.
- [30] R. Chowdhury, J. I. Candela-Lena, M. C. Chan, D. J. Greenald, K. K. Yeoh, Y. M. Tian, M. A. McDonough, A. Tumber, N. R. Rose, A. Conejo-Garcia, M. Demetriades, S. Mathavan, A. Kawamura, M. K. Lee, F. van Eeden, C. W. Pugh, P. J. Ratcliffe, C. J. Schofield, *ACS Chem Biol* **2013**, *8*, 1488-1496.
- [31] R. Chowdhury, K. K. Yeoh, Y. M. Tian, L. Hillringhaus, E. A. Bagg, N. R. Rose, I. K. Leung, X. S. Li, E. C. Woon, M. Yang, M. A. McDonough, O. N. King, I. J. Clifton, R. J. Klose, T. D. Claridge, P. J. Ratcliffe, C. J. Schofield, A. Kawamura, *EMBO Rep.* **2011**, *12*, 463-469.
- [32] F. H. Niesen, H. Berglund, M. Vedadi, *Nat. Protoc.* **2007**, *2*, 2212-2221.
- [33] T.-L. Yeh, T. M. Leissing, M. I. Abboud, C. C. Thinner, O. Atasoylu, J. P. Holt-Martyn, D. Zhang, A. Tumber, K. Lippl, C. T. Lohans, I. K. H. Leung, H. Morcrette, I. J. Clifton, T. D. W. Claridge, A. Kawamura, E. Flashman, X. Lu, P. J. Ratcliffe, R. Chowdhury, C. W. Pugh, C. J. Schofield, *Chem. Sci.* **2017**, *8*, 7651-7668.


## FULL PAPER

- [34] M. C. Chan, O. Atasoylu, E. Hodson, A. Tumber, I. K. Leung, R. Chowdhury, V. Gómez-Pérez, M. Demetriades, A. M. Rydzik, J. Holt-Martyn, Y. M. Tian, T. Bishop, T. D. Claridge, A. Kawamura, C. W. Pugh, P. J. Ratcliffe, C. J. Schofield, *PLoS One* **2015**, *10*, e0132004.
- [35] M. I. Abboud, T. E. McAllister, I. K. H. Leung, R. Chowdhury, C. Jorgensen, C. Domene, J. Mecinović, K. Lippl, R. L. Hancock, R. J. Hopkinson, A. Kawamura, T. D. W. Claridge, C. J. Schofield, *Chem Commun (Camb)* **2018**, *54*, 3130-3133.
- [36] Y. M. Tian, K. K. Yeoh, M. K. Lee, T. Eriksson, B. M. Kessler, H. B. Kramer, M. J. Edelman, C. Willam, C. W. Pugh, C. J. Schofield, P. J. Ratcliffe, *J. Biol. Chem.* **2011**, *286*, 13041-13051.
- [37] C. C. Thinner, K. S. England, A. Kawamura, R. Chowdhury, C. J. Schofield, R. J. Hopkinson, *Biochim. Biophys. Acta* **2014**, *1839*, 1416-1432.
- [38] J. D. Vasta, R. T. Raines, *J Med Chem* **2018**,
- [39] C. H. Arrowsmith, J. E. Audia, C. Austin, J. Baell, J. Bennett, J. Blagg, C. Bountra, P. E. Brennan, P. J. Brown, M. E. Bunnage, C. Buser-Doepner, R. M. Campbell, A. J. Carter, P. Cohen, R. A. Copeland, B. Cravatt, J. L. Dahlin, D. Dhanak, A. M. Edwards, M. Frederiksen, S. V. Frye, N. Gray, C. E. Grimshaw, D. Hepworth, T. Howe, K. V. Huber, J. Jin, S. Knapp, J. D. Kotz, R. G. Kruger, D. Lowe, M. M. Mader, B. Marsden, A. Mueller-Fahrnow, S. Müller, R. C. O'Hagan, J. P. Overington, D. R. Owen, S. H. Rosenberg, B. Roth, B. Roth, R. Ross, M. Schapira, S. L. Schreiber, B. Shoichet, M. Sundström, G. Superti-Furga, J. Taunton, L. Toledo-Sherman, C. Walpole, M. A. Walters, T. M. Willson, P. Workman, R. N. Young, W. J. Zuercher, *Nat Chem Biol* **2015**, *11*, 536-541.
- [40] F. McMurray, M. Demetriades, W. Aik, M. Merkestein, H. Kramer, D. S. Andrew, C. L. Scudamore, T. A. Hough, S. Wells, F. M. Ashcroft, M. A. McDonough, C. J. Schofield, R. D. Cox, *PLoS One* **2015**, *10*, e0121829.

## FULL PAPER

## Entry for the Table of Contents

## FULL PAPER



Humans have 60-70 2OG-dependent oxygenases, of which some are important therapeutic targets. A priority in the development of oxygenase inhibitors is to identify scaffolds selective for a particular enzyme, made challenging by the highly-conserved oxygenase active sites. Here, we describe the development of potent inhibitors for the oxygenase OGFOD1 which do not inhibit related enzymes, such as the physiologically important oxygenase PHD2.

*Dr. Cyrille C. Thines, Dr. Christopher T. Lohans, Dr. Martine I. Abboud, Dr. Tzu-Lan Yeh, Dr. Anthony Tumber, Dr. Radosław P. Nowak, Dr. Martin Attwood, Dr. Matthew E. Cockman, Prof. Dr. Udo Oppermann, Prof. Dr. Christoph Loenarz, Prof. Dr. Christopher J. Schofield\**

**Page No. – Page No.**

**Selective inhibitors of a human prolyl hydroxylase (OGFOD1) involved in ribosomal decoding**