

Biomimetic synthesis of marine sponge metabolite spiculoic acid **1** and establishment of the absolute configuration of the natural product†

James E. D. Kirkham,^a Victor Lee^{*a} and Jack E. Baldwin^{*b}

Received (in Cambridge, UK) 17th May 2006, Accepted 26th May 2006

First published as an Advance Article on the web 12th June 2006

DOI: 10.1039/b607035c

The synthesis of spiculoic acid **1** using a biomimetic Diels–Alder reaction is described; comparison of the specific rotation of the natural and synthetic material revealed that the enantiomer of the natural product has been synthesized.

Spiculoic acid **1** (Fig. 1) is a polyketide recently isolated from the extracts of the Caribbean marine sponge *Plakortis angulospiculatus* by Andersen *et al.*¹ Spiculoic acid **1** was found to be optically active and its structure and relative stereochemistries were established based on a combination of NMR and mass spectrometry studies. Further activity assay showed that **1** is cytotoxic towards the human breast cancer MCF-7 cell line (IC₅₀ 8 µg/mL).

At the time of its isolation and structure elucidation **1** was considered to be unprecedented due to its spiculane carbon skeleton. Six stereocentres reside within this relatively small bicyclic core, two of which are quaternary and adjacent to each other. A group of similar compounds were subsequently isolated from the marine sponge *Plakortis zyggompha*.²

Andersen *et al.* have proposed that **1** is biosynthesized from **2** through an enzyme catalysed Diels–Alder reaction (Fig. 1).^{1,3} Our continued interest in biomimetic chemistry⁴ prompted us to examine the application of this proposed intramolecular Diels–Alder reaction in the synthesis of **1**.⁵ Herein we report the successful execution of our synthetic plan and the establishment of the absolute configuration of **1**. During the course of our investigation Mehta and Kundu disclosed a synthetic approach to **1** and prepared a highly truncated analogue of the natural product.⁶

Realizing the potential sensitivity of **2** we elected to use **3** as its surrogate. Since the absolute configuration of natural **1** was unknown, we decided to use the chiral structure of **1** depicted in Andersen's publication as our synthetic target. The synthesis of **3** commenced with the conversion of Roche ester **4** into aldehyde **5** in an overall yield of 98% by protection of the primary alcohol using imidazole and *tert*-butyldimethylsilyl chloride,⁷ followed by reduction of the ester⁸ to primary alcohol and oxidation to





Fig. 1 Retrosynthetic analysis of spiculoic acid **1**.

^aChemistry Research Laboratory, University of Oxford, 12 Mansfield Road, Oxford, UK OX1 3TA. E-mail: victor.lee@chem.ox.ac.uk; Fax: +44 (0)1865 285 002; Tel: +44 (0)1865 275 650

^bChemistry Research Laboratory, University of Oxford, 12 Mansfield Road, Oxford, UK OX1 3TA. E-mail: jack.baldwin@chem.ox.ac.uk; Fax: +44 (0)1865 275 632; Tel: +44 (0)1865 275 671

† Electronic supplementary information (ESI) available: ¹H, ¹³C, DEPT, COSY, HSQC, NOESY and TOCSY NMR spectra for (–)-**1**. See DOI: 10.1039/b607035c




Scheme 1 Reagents and conditions: (i) (a) TBSCl, imidazole, THF; (b) LiBH₄, THF, reflux; (c) DMSO, (COCl)₂, ^tPr₂NEt, CH₂Cl₂, –78 °C, 98% over three steps; (ii) **6**, ⁿBu₂BOTf, ^tPr₂NEt, CH₂Cl₂, –78 °C, 89%; (iii) LiBH₄, MeOH, H₂O, 73%; (iv) *para*-methoxybenzaldehyde dimethyl acetal, pyridinium tosylate, CH₂Cl₂, 96%; (v) TBAF, THF, 100%; (vi) IBX, THF, DMSO, 100%.


aldehyde **5** using Swern's conditions⁹ (Scheme 1). Evans' aldol condensation¹⁰ of **5** and oxazolidinone **6**¹¹ delivered secondary alcohol **7** in 89% yield. Reductive removal of the chiral auxiliary was effected with lithium borohydride in wet methanol¹² to afford diol **8** in 73% yield. Diol **8** was converted into *para*-methoxybenzylidene acetal **9** in 96% yield using *para*-methoxybenzaldehyde dimethyl acetal and a catalytic amount of pyridinium tosylate.¹³ This was followed by removal of the TBS group with TBAF to afford primary alcohol **10** in quantitative yield, which was oxidized to aldehyde **11**, again in quantitative yield, using iodoxybenzoic acid (IBX).¹⁴

Wittig olefination of aldehyde **11** with stabilized ylide **12**¹⁵ gave ester **13** in 77% yield (Scheme 2). The *E* stereochemistry of the double bond in **13** was confirmed by NOE studies. Ester **13** was reduced to alcohol **14** with lithium aluminium hydride¹⁶ in 99% yield, which was oxidized to unsaturated aldehyde **15** by iodoxybenzoic acid (IBX) in quantitative yield. Treating aldehyde **15** with ylide **12** delivered the doubly conjugated ester **16** in 65% yield. Again, the geometry of the newly formed double bond was confirmed to be *E* by NOE experiment. Ester **16** was reduced to alcohol **17** in 96% yield, which was followed by oxidation to aldehyde **18** in 100% yield.

Aldehyde **18** was subjected to a Julia/Kocienski olefination¹⁷ with benzylsulfone **19**¹⁸ using lithium bis(trimethylsilyl)amide as base to give *E,E,E* triene **20** in 82% yield (Scheme 3). Reductive cleavage of the *para*-methoxybenzylidene acetal was effected with diisobutylaluminium hydride¹³ to give primary alcohol **21** in 82% yield.




Scheme 2 Reagents and conditions: (i) **12**, toluene, 100 °C; (ii) LiAlH₄, THF, 0 °C; (iii) IBX, DMSO, THF.



Scheme 3 Reagents and conditions: (i) **19**, LiN(TMS)₂, THF, 82%; (ii) DIBAL, CH₂Cl₂, 82%; (iii) IBX, DMSO, THF, 80%; (iv) **23**, toluene, 100 °C, 25%.

yield, which was oxidized to aldehyde **22** in 80% yield with IBX. The final Wittig olefination in our synthesis was conducted by heating a mixture of aldehyde **22** with ylide **23**. Gratifyingly, this delivered bicyclic structure **24** directly in 25% yield. Presumably the desired Wittig reaction took place to afford **3**, which underwent a Diels–Alder reaction *in situ* to give **24**. We also isolated a 5–20% yield of an α,β -unsaturated aldehyde, which was formed by the E1cB elimination of *para*-methoxybenzyl alcohol from **22**.

The final stage of the synthesis of **1** began with removal of the *para*-methoxybenzyl protecting group from **24** with 2,3-dichloro-5,6-dicyanobenzoquinone (DDQ) under buffered conditions¹⁹ to afford secondary alcohol **25**, which was found to co-elute with the *para*-methoxybenzaldehyde formed in the DDQ deprotection during flash chromatography (Scheme 4). Therefore, this partially purified mixture was subjected to a catalytic amount of tetrakis(triphenylphosphine)palladium(0) in the presence of excess morpholine²⁰ to remove the allyl ester. This delivered free acid **26** in a combined yield of 94% over two steps after chromatographic



Scheme 4 Reagents and conditions: (i) DDQ, CH₂Cl₂, pH 7 buffer; (ii) (PPh₃)₄Pd, morpholine, 94% over two steps; (iii) IBX, EtOAc, 70 °C, 45%.




Fig. 2 Selected NOESY correlations of (-)-1.

purification. Finally, oxidation of **26** with IBX in hot ethyl acetate²¹ gave spiculoic acid A (**1**) in 45% yield.

Extensive spectroscopic studies were conducted on compounds **24**, **25**, **26** and **1** to confirm their structures and relative stereochemistries. Analysis of the spectroscopic data for synthetic **1** gave an H-5/H-9 coupling constant of 11.8 Hz, indicating that both protons are axial and that the [4.3.0] bicycle is *trans*. NOESY studies indicated that the correct relative stereochemistry has been achieved *via* the Diels–Alder reaction (Fig. 2). The absolute stereochemistry of **1** was established through knowledge of the absolute configuration of the C-6 and C-8 stereocentres. The ¹H and ¹³C NMR spectra of synthetic **1** were indistinguishable from those of the natural product.

The specific rotation of synthetic **1** ($[\alpha]_D^{22} = -97$, $c = 0.16$, CH₂Cl₂) is comparable in magnitude to that of natural **1** ($[\alpha]_D = +110$, $c = 0.1$, CH₂Cl₂),¹ but of opposite sign. Therefore, we conclude that the absolute configuration of natural spiculoic acid A (**1**) is as depicted in Fig. 3.

In summary, we have demonstrated the application of a biomimetic strategy to the synthesis of spiculoic acid A (**1**) and through our synthesis have established the absolute configuration of the natural product. Analogues iso- (**27**), *nor*- (**28**) and *dinor*- (**29**) spiculoic acids A (Fig. 4), recently isolated from the marine sponge *Plakortis zyggompha*,² all possess specific rotations that are positive and large in value. Therefore, it is tempting to imply that the absolute configurations of these compounds are identical to that of (+)-**1**.





Fig. 3 Absolute configuration of natural spiculoic acid A (**1**).



Isospiculoic acid A (**27**) R₁ = Et, R₂ = Et
Nor-spiculoic acid A (**28**) R₁ = Et, R₂ = Me
Dinor-spiculoic acid A (**29**) R₁ = Me, R₂ = Me

Fig. 4 Speculated absolute configurations of iso- (**27**), *nor*- (**28**) and *dinor*-spiculoic acids A (**29**).

We thank Hoffmann-La Roche for partial support of this work.

Notes and references

- X.-H. Huang, R. van Soest, M. Roberge and R. J. Andersen, *Org. Lett.*, 2004, **6**, 75–78.
- F. Berru , O. P. Thomas, R. Fern ndez and P. Amade, *J. Nat. Prod.*, 2005, **68**, 547–549.
- (a) E. M. Stocking and R. M. Williams, *Angew. Chem., Int. Ed.*, 2003, **42**, 3078–3115; (b) H. Oikawa and T. Tokiwano, *Nat. Prod. Rep.*, 2004, **21**, 321–352; (c) H. Oikawa, *Bull. Chem. Soc. Jpn.*, 2005, **78**, 537–554.
- J. E. Moses and R. M. Adlington, *Chem. Commun.*, 2005, 5945–5952.
- K. Takao, R. Munakata and K. Tadano, *Chem. Rev.*, 2005, **105**, 4779–4807.
- G. Mehta and U. K. Kundu, *Org. Lett.*, 2005, **7**, 5569–5572.
- N. Tsukada, T. Shimada, Y. S. Gyoung, N. Asao and Y. Yamamoto, *J. Org. Chem.*, 1995, **60**, 143–148.
- K. Mori and K. Koseki, *Tetrahedron*, 1988, **44**, 6013–6020.
- J. M. Minguez, S.-Y. Kim, K. A. Giuliano, R. Balachandran, C. Madiraju, B. W. Day and D. P. Curran, *Bioorg. Med. Chem.*, 2003, **11**, 3335–3357.
- D. A. Evans, J. V. Nelson, E. Vogel and T. R. Taber, *J. Am. Chem. Soc.*, 1981, **103**, 3099–3111.
- M. A. Brimble, *Aust. J. Chem.*, 1990, **43**, 1035–1046.
- T. D. Penning, S. W. Djuric, R. A. Haack, V. J. Kalish, J. M. Miyashiro, B. W. Rowell and S. S. Yu, *Synth. Commun.*, 1990, **20**, 307–312.
- Z. H. Peng and K. A. Woerpel, *J. Am. Chem. Soc.*, 2003, **125**, 6018–6019.
- (a) M. Frigerio and M. Santagostino, *Tetrahedron Lett.*, 1994, **35**, 8019–8022; (b) M. Frigerio, M. Santagostino, S. Sputore and G. Palmisano, *J. Org. Chem.*, 1995, **60**, 7272–7276; (c) M. Frigerio, M. Santagostino and S. Sputore, *J. Org. Chem.*, 1999, **64**, 4537–4538.
- S. E. Sen and G. J. Ewing, *J. Org. Chem.*, 1997, **62**, 3529–3536.
- R. St rmer and R. W. Hoffmann, *Chem. Ber.*, 1994, **127**, 2519–2526.
- (a) P. R. Blakemore, W. J. Cole, P. J. Kocienski and A. Morley, *Synlett*, 1998, 26–29; (b) P. R. Blakemore, *J. Chem. Soc., Perkin Trans. 1*, 2002, 2563–2585.
- H. Hilpert and B. Wirz, *Tetrahedron*, 2001, **57**, 681–694.
- P. Wipf and T. H. Graham, *J. Am. Chem. Soc.*, 2004, **126**, 15346–15347.
- S. Friedrich-Bochnitschek, H. Waldmann and H. Kunz, *J. Org. Chem.*, 1989, **54**, 751–756.
- J. D. More and N. S. Finney, *Org. Lett.*, 2002, **4**, 3001–3003.