

## The Raman Spectra of Boron Trifluoride, Trichloride, and Tribromide. The Effect of the Boron Isotopes

Thomas F. Anderson, Edwin N. Lassettre, and Don M. Yost

Citation: [The Journal of Chemical Physics](#) **4**, 703 (1936); doi: 10.1063/1.1749772

View online: <http://dx.doi.org/10.1063/1.1749772>

View Table of Contents: <http://scitation.aip.org/content/aip/journal/jcp/4/11?ver=pdfcov>

Published by the [AIP Publishing](#)

---

### Articles you may be interested in

[Studies of Electron Impact Excitation, Negative Ion Formation, and Negative IonMolecule Reactions in Boron Trifluoride and Boron Trichloride](#)

J. Chem. Phys. **56**, 3336 (1972); 10.1063/1.1677702

[Infrared and Raman Spectra of Nitrogen Trifluoride](#)

J. Chem. Phys. **23**, 1248 (1955); 10.1063/1.1742250

[Thermodynamic Properties of Gaseous Boron Trifluoride, Boron Trichloride, and Boron Tribromide](#)

J. Chem. Phys. **14**, 729 (1946); 10.1063/1.1724094

[The Raman Spectrum of Boron Trifluoride Gas](#)

J. Chem. Phys. **6**, 424 (1938); 10.1063/1.1750286

[Raman Spectra of Benzene and Isotope Effect](#)

J. Chem. Phys. **6**, 8 (1938); 10.1063/1.1750124

---



## The Raman Spectra of Boron Trifluoride, Trichloride, and Tribromide. The Effect of the Boron Isotopes\*

THOMAS F. ANDERSON, EDWIN N. LASSETTRE AND DON M. YOST, *California Institute of Technology, Pasadena, California*

(Received August 12, 1936)

The Raman frequencies  $\nu_1\{1\}$ ,  $2\nu_2\{2\}$ ,  $\nu_3\{2\}$ , and  $\nu_4\{2\}$  for  $\text{BF}_3(\text{g})$ ,  $\text{BCl}_3(\text{l})$ , and  $\text{BBr}_3(\text{l})$  were found to be 886 and 783, 1394, 1038 and 1105, 440; 471, 924, 958 and 996, 243; 279, 743, 806 and 846, and 151  $\text{cm}^{-1}$ , respectively. The assignment of frequencies was determined by polarization experiments on  $\text{BCl}_3$  and is confirmed for the three compounds by the results of a normal coordinate treatment. The doubling of  $\nu_1$  in  $\text{BF}_3$  is ascribed to vibrational resonance between  $\nu_1$  and  $2\nu_4$ . The doubling of  $\nu_3$  in all three compounds is shown to be due to the presence of the two isotopes of boron  $\text{B}^{10}$  and  $\text{B}^{11}$ . A satisfactory assignment of the infrared frequencies of  $\text{BCl}_3$  is made.

### INTRODUCTION

THE Raman spectrum of boron trichloride has been reported by Bhagavantam and by Venkateswaran.<sup>1</sup> Cassie<sup>2</sup> has proposed an assignment for the infrared frequencies he observed for boron trichloride, but he assumed frequencies differing rather widely from the Raman frequencies as well as the presence of forbidden lines in the infrared spectrum.

We have reinvestigated the Raman spectrum of boron trichloride and report as well the Raman spectra of boron tribromide and boron trifluoride. In all three compounds a doubling of certain frequencies by the presence of the boron isotopes  $\text{B}^{10}$  and  $\text{B}^{11}$  is observed. The infrared spectrum of boron trichloride is satisfactorily explained by an assignment of frequencies which is confirmed by the results of a normal coordinate treatment.

The work on boron trifluoride was started at this laboratory in 1934 by Mr. S. T. Gross.

This and other similar investigations were assisted by a grant-in-aid of research to one of us (D. M. Y.) from the National Research Council. The boron tribromide used was prepared as an N.Y.A. project by Mr. David Pressman.

### GENERAL THEORETICAL CONSIDERATIONS

The shape of the boron trichloride molecule is rather definitely shown to be that of a plane equilateral triangle with the symmetry  $D_{3h}$ . This

follows from its zero dipole moment<sup>3</sup> and the results of electron diffraction experiments.<sup>4</sup> The results of the present investigation are in agreement with the assumption that the other boron trihalides also have the symmetry  $D_{3h}$ .

The  $\text{BX}_3$  molecule then has the four symmetry modes of vibration indicated in Fig. 1.  $\nu_1\{1\}$ ,  $\nu_3\{2\}$ , and  $\nu_4\{2\}$  are allowed in the Raman effect with the depolarizations 0, 6/7, and 6/7, respectively.<sup>5</sup> Numbers in braces adjoining  $\nu$ 's are degeneracies.  $\nu_2\{1\}$ ,  $\nu_3\{2\}$  and  $\nu_4\{2\}$  are active in the infrared. The selection rules for simple combinations and overtones obtained by the methods of group theory<sup>6</sup> are indicated in Table I.

The fact must be borne in mind that the presence of the isotopes of boron  $\text{B}^{11}$  and  $\text{B}^{10}$  in the approximate ratio of 4 to 1 will produce a doubling of those frequencies in which the boron atom moves appreciably. Thus in the Raman effect we should expect  $\nu_3$  and  $2\nu_2$  each to appear as a doublet separated in frequencies by about 5 percent, and with the lower frequency about four times as intense as the higher. Analogous splits in frequencies should appear in the infrared spectrum with a ratio of extinction coefficients for the two components of 4 to 1. A normal coordi-

\* Contribution No. 562 from the Gates and Crellin Chemical Laboratory of the California Institute of Technology.

<sup>1</sup> Bhagavantam, *Ind. J. Phys.* **5**, 73 (1930); Venkateswaran, *ibid.* **6**, 284 (1931).

<sup>2</sup> A. B. D. Cassie, *Proc. Roy. Soc. A* **148**, 87 (1935).

<sup>3</sup> E. Bergmann and L. Engel, *Physik. Zeits.* **32**, 425 (1931); H. Ulich and W. Nespital, *Zeits. f. Elektrochem.* **37**, 559 (1931); W. Nespital, *Zeits. f. physik. Chemie* **B16**, 153 (1932).

<sup>4</sup> L. O. Brockway, *Rev. Mod. Phys.* **8**, 231 (1936); private communication.

<sup>5</sup> E. B. Wilson, Jr., *J. Chem. Phys.* **2**, 432 (1934).

<sup>6</sup> E. Wigner, *Gött. Nachr.*, 133 (1930); G. Placzek, *The Structure of Molecules* (Blackie and Son, London, 1932); L. Tisza, *Zeits. f. Physik* **82**, 48 (1933); E. B. Wilson, Jr., *Phys. Rev.* **45**, 706 (1934).




FIG. 1. Symmetry modes of the plane symmetrical  $BX_3$  molecule.

nate treatment should enable one to predict the isotopic doubling of frequencies with greater accuracy.

### EXPERIMENTAL

Boron trifluoride was prepared by treating ammonium fluoborate with concentrated sulfuric acid and heating. The reaction chamber was of glass, so silicon tetrafluoride was expected as an impurity. To prepare ammonium fluoborate, boric acid was treated with hydrofluoric acid in a copper beaker; the solution was neutralized with  $NH_4OH$  and concentrated by heating. Upon cooling the solution,  $NH_4BF_4$  crystallized out. The  $BF_3$  so prepared was fractionated repeatedly; the final product had a melting point of  $-127 \pm 1^\circ$ , in good agreement with the value given in the literature. The Raman spectrum of the somewhat cloudy liquid under its own vapor pressure of 6 atmospheres at the temperature of a solid carbon dioxide alcohol bath was taken using a technique similar to that previously employed for phosphorus trifluoride.<sup>7</sup> The Raman spectrum of the gas at a pressure of six atmospheres was taken using the 2537A line of mercury as a source of radiation. The spectrum from the gas was much more satisfactory than that from the liquid.

Kahlbaum's C.P. boron trichloride ( $b_{760} 12.5^\circ$ ) was distilled into a Raman tube and sealed in without further purification. The Raman spectrum was obtained at room temperature from the liquid under its own vapor pressure. The polarizations of the two most intense lines were roughly determined. For this purpose<sup>8</sup> the mercury lamp was placed below the horizontal Raman tube with a series of parallel vanes of black cardboard between lamp and tube to insure that the light entering the tube was

TABLE I. An addition table showing the selection rules for simple combinations and overtones of modes of vibration of  $BX_3$  molecules.

	$\nu_1\{1\}R^*$	$\nu_2\{1\}I$	$\nu_3\{2\}RI$	$\nu_4\{2\}RI$
$\nu_1\{1\}R$	$R$	$I$	$RI$	$RI$
$\nu_2\{1\}I$		$R^*$	$R$	$R$
$\nu_3\{2\}RI$			$RI$	$RI$
$\nu_4\{2\}RI$				$RI$

$R$  indicates a Raman active combination;  $I$  indicates an infrared active combination.

\*  $3\nu_2$  is infrared active.

parallel. The scattered light from the Raman tube then passed successively through a half-wave plate for 4400A, a large Nicol prism with vertical axis, and a lens which focused the light on the 0.4 mm slit of the spectrograph. With the axis of the half-wave plate at an angle of  $45^\circ$  to that of the Nicol, only horizontally polarized light from the Raman tube passed through to the slit. When the axis of half-wave plate and Nicol were parallel, only the perpendicular component reached the slit. The advantage of this arrangement is that all light entering the spectrograph is vertically polarized so that there is no effect of selective transmission or reflection of the light from the slit or prism faces, in contrast to previously used arrangements. Two one-hour exposures were made on the same photographic plate with the two settings of the half-wave plate. Relative intensities of the lines in the two spectra were estimated visually and compared with the relative intensities of the Rayleigh lines which are known to be polarized.

Boron tribromide ( $b_{740} 90.1^\circ$ ) was prepared by passing bromine vapor over a mixture of boric oxide and carbon at  $250^\circ$ . It was carefully fractionated, the fraction used having a boiling point of  $91.0 \pm 0.2^\circ$ . Difficulty was had in sealing the material in the Raman tube without the liberation of small amounts of bromine. The bromine was therefore removed by sealing a small amount of mercury in the Raman tube with the boron tribromide. The mercurous bromide, which was formed on shaking, settled out nicely and Raman spectra of the clear colorless liquid remaining were obtained.

### RESULTS AND ASSIGNMENT OF FREQUENCIES

The observed Raman frequencies and assignments are presented in Table II. We shall discuss

<sup>7</sup> D. M. Yost and T. F. Anderson, J. Chem. Phys. 2, 624 (1934).

<sup>8</sup> The arrangement here described, which has been in use in this laboratory for some time, was developed by Dr. Fred B. Stitt of this laboratory. He will publish a more detailed account shortly.

first the more complete results for  $\text{BCl}_3$  and proceed then to a discussion of  $\text{BBr}_3$  and  $\text{BF}_3$ .

### Boron trichloride

The results for boron trichloride agree with those obtained by Venkateswaran<sup>1</sup> with the exception that the broad band he observed at  $947\text{ cm}^{-1}$  is resolved into a triplet. This group, consisting of a sharp line and two broad lines, is reproduced in Fig. 2. The depolarization of the  $471\text{ cm}^{-1}$  line was found to be less than 0.25; it was therefore assigned to the mode  $\nu_1$ . The depolarization of the line at  $253\text{ cm}^{-1}$  was close to unity and it was therefore taken to be  $\nu_4$ . The frequency  $\nu_3$  was taken to be the expected isotopic doublet  $946, 989\text{ cm}^{-1}$ , the broadness of this pair distinguishing it from the sharp line at  $902\text{ cm}^{-1}$ . The latter frequency was then assumed to be  $2\nu_2$  whose weaker isotopic component is obscured by the  $946\text{ cm}^{-1}$  frequency.

When we make the assignment of infrared absorption frequencies<sup>2</sup> indicated in Table III, we arrive at the values for the vibrational frequencies for  $\text{BCl}_3(g)$  indicated in Table II. These frequencies differ only slightly from the Raman frequencies for  $\text{BCl}_3(l)$ . The agreement between observed and calculated frequencies is quite satisfactory with the exception of the assignment of the frequency  $845\text{ cm}^{-1}$ . Its nearness to the frequency  $834\text{ cm}^{-1}$  observed in the Raman spectrum of phosgene<sup>9</sup> and the fact that phosgene is often a by-product in the preparation of  $\text{BCl}_3$  and difficult to separate from it, suggest that the presence of this band may be due to a trace of phosgene in the  $\text{BCl}_3$

TABLE II. The Raman spectra of boron trifluoride, boron trichloride, and boron tribromide.

SUBSTANCE	$\nu_1\{1\}$ ( $\text{cm}^{-1}$ )	$2\nu_2\{1\}$ ( $\text{cm}^{-1}$ )	$\nu_3\{2\}$ ( $\text{cm}^{-1}$ )	$\nu_4\{2\}$ ( $\text{cm}^{-1}$ )
$\text{B}^{11}\text{F}_3(g)$	886(7), 783(3)	1394( $\frac{1}{2}$ ?)*	1038(1)	440(4b) $\pm 25^*$
$\text{B}^{10}\text{F}_3(g)$	886(7), 783(3)	(1452)	1105( $\frac{1}{2}$ ?)*	440(4b) $\pm 25^*$
$\text{B}^{11}\text{F}_3(l)$	883(1), —	—	—	—
$\text{B}^{11}\text{Cl}_3(g)^\dagger$	471	924	958	243
$\text{B}^{10}\text{Cl}_3(g)^\dagger$	471	(966)	996	243
$\text{B}^{11}\text{Cl}_3(l)$	471(10)	902(3)	946(3b)	253(8)
$\text{B}^{10}\text{Cl}_3(l)$	471(10)	(943)	989(1b)	253(8)
$\text{B}^{11}\text{Br}_3(l)$	279(10)	743( $\frac{1}{2}$ )	806(3/4b)	151(8)
$\text{B}^{10}\text{Br}_3(l)$	279(10)	(780)	846( $\frac{1}{2}$ b)	151(8)

Numbers in parentheses adjoining frequency values indicate relative intensities. *b* = broad; all other lines are sharp. Numbers in braces adjoining  $\nu$ s are multiplicities. Frequencies in parentheses are estimated.

\* The assignment of these frequencies is provisional for  $\text{BF}_3(g)$ .

† Frequencies obtained from infrared data for  $\text{BCl}_3(g)$ .

<sup>9</sup> Kohlrausch, *Der Smekal-Raman-Effekt* (Julius Springer, Berlin, 1931).




FIG. 2. The effect of the boron isotopes in the Raman spectrum of  $\text{BCl}_3$ .

used. The infrared spectrum of phosgene has not been reported. The weaker isotopic components of  $\nu_3 + \nu_4$  and of  $\nu_1 + \nu_3$  are evidently too weak to be observed, while that of  $3\nu_2$  is obscured by  $\nu_1 + \nu_3$  from  $\text{B}^{11}\text{Cl}_3$ . In other cases the estimated relative intensities of the bands are in accord with the relative abundances of the boron isotopes. Unfortunately Cassie has not published data from which extinction coefficients could be calculated. The assignment of frequencies is confirmed by the results of the normal coordinate treatment which follows this section.

### Boron tribromide

The Raman spectrum of boron tribromide is very similar in appearance to that of boron

TABLE III. The infrared spectrum of boron trichloride.

$\nu_{\text{obs.}}$ ( $\text{cm}^{-1}$ )	$\nu_{\text{calc.}}$ ( $\text{cm}^{-1}$ )	Combination
714(4)	714	$\nu_1 + \nu_4$
845(3)	834(R)	$\text{COCl}_2$ (?)
958(100)	958	$\nu_3, \text{B}^{11}\text{Cl}_3$
996(50)	996	$\nu_3, \text{B}^{10}\text{Cl}_3$
1206(2)	1201	$\nu_3, \text{B}^{11}\text{Cl}_3 + \nu_4$
1387(6)	1386	$3\nu_2, \text{B}^{11}\text{Cl}_3$
1429(6)	1429; 1452	$\nu_1 + \nu_3, \text{B}^{11}\text{Cl}_3; 3\nu_2, \text{B}^{10}\text{Cl}_3$
1912(2.5)	1916	$2\nu_3, \text{B}^{11}\text{Cl}_3$
1995(2)	1992	$2\nu_3, \text{B}^{10}\text{Cl}_3$

R = observed in Raman spectrum of  $\text{COCl}_2(l)$ .

TABLE IV. Calculated force constants  $K$ ,  $H$ ,  $G$ , and  $K'$ , calculated frequencies for trihalides containing  $B^{10}$ , and calculated internuclear distances  $B-X$ , for boron trihalides.

SUBSTANCE	$K \times 10^{-5}$ (dynes/cm)	$H \times 10^{-5}$ (dynes/cm)	$G \times 10^{-5}$ (dynes/cm)	$K' \times 10^{-5}$ (dynes/cm)	$\nu_3 B^{10} X_3$ calc ( $cm^{-1}$ )	$\nu_3 B^{10} X_3$ obs. ( $cm^{-1}$ )	$\nu_4 B^{10} X_3$ calc. ( $cm^{-1}$ )	$\nu_4 B^{10} X_3$ obs. ( $cm^{-1}$ )	B-X calc. (Å)	B-X obs. (Å)
$BF_3(g)$	4.24	0.60	0.88	1.73	1080	1105(?)	416	416*	1.49	1.42†
$BCl_3(g)$	3.30	0.276	0.434	0.653	1000	996	243	243	1.72	1.73
$BBr_3(l)$	2.50	0.252	0.270	0.61	843	846	151	151	1.85	1.87

\* This value was assumed for  $\nu_1$ ,  $B^{11}F_3$  in order to obtain real values for the constants.

† The sum of Pauling's double bond radii plus 0.04 Å was taken as an estimate for the B-F distance. See L. Pauling, L. O. Brockway and J. Y. Beach, J. Am. Chem. Soc. **57**, 2705 (1935).

trichloride with  $2\nu_2$  and the isotopic shift of  $\nu_3$  both appearing. By analogy with the assignment of the  $BCl_3$  frequencies the assignment of frequencies indicated in Table II was made; confirmation is again obtained from the normal coordinate treatment.

### Boron trifluoride

The Raman spectrum obtained from boron trifluoride is much more difficult to interpret. In addition to the frequencies listed in Table II, a line, doubtless due to the presence of  $SiF_4$  ( $\nu_1=800(10) \text{ cm}^{-1}$ ),<sup>10</sup> was observed at  $801(4) \text{ cm}^{-1}$ . The line assigned to  $\nu_4$  whose center is at  $440 \text{ cm}^{-1}$ , is  $50 \text{ cm}^{-1}$  in width. Its integrated intensity is therefore quite great. We then attribute the strong doublet, which appears as  $\nu_1$  in our assignment, to vibrational resonance between  $\nu_1$  and  $2\nu_4$ . Aside from the presence of impurities this appears to be the only reasonable possibility of accounting for the existence of both lines. If we introduce the perturbation term

$$H' = \gamma Q_1(Q_{4a}^2 + Q_{4b}^2)$$

(where the  $Q$ 's represent the normal coordinates and  $\gamma$  is a constant) in the potential energy expression one obtains the following energy levels from the solution of the secular equation:<sup>11</sup>

$$\begin{aligned}\lambda_1\{2\} &= 2\nu_4, \\ \lambda_2\{1\} &= \frac{1}{2}[2\nu_4 + \nu_1 + \{(2\nu_4 - \nu_1)^2 + 8P^2\}^{\frac{1}{2}}], \\ \lambda_3\{1\} &= \frac{1}{2}[2\nu_4 + \nu_1 - \{(2\nu_4 - \nu_1)^2 + 8P^2\}^{\frac{1}{2}}].\end{aligned}$$

We should expect  $\lambda_2$  and  $\lambda_3$  to appear, while the simple overtone  $\lambda_1$  should be very weak in the Raman effect.  $P$ , the perturbation energy defined in the usual manner, has the value  $35 \text{ cm}^{-1}$  when

it is assumed that  $\nu_1 = 2\nu_4 = 835 \text{ cm}^{-1}$ . This value is of the usual order of magnitude for  $P$ . The assignment of other observed frequencies is confirmed by the rather satisfying results of the following normal coordinate treatment. An investigation of the infrared absorption spectrum of  $BF_3$  would doubtless yield a unique assignment.

### NORMAL COORDINATES; COMPUTATION OF ISOTOPIC SHIFTS IN FREQUENCIES

Normal coordinate treatments of the plane symmetrical  $AB_3$  type molecule have been made by a number of authors.<sup>12</sup> We shall use the valence force potential function  $V$  with force constants chosen so that

$$\begin{aligned}2V &= K(\Delta R_1^2 + \Delta R_2^2 + \Delta R_3^2) \\ &\quad + HR_0^2(\Delta \alpha_{12}^2 + \Delta \alpha_{23}^2 + \Delta \alpha_{13}^2) \\ &\quad + GR_0^2(\Delta \beta_1^2 + \Delta \beta_2^2 + \Delta \beta_3^2) \\ &\quad + 2K'(\Delta R_1 \Delta R_2 + \Delta R_2 \Delta R_3 + \Delta R_1 \Delta R_3).\end{aligned}$$

$R_1$  is here the distance  $A-B_i$  and  $R_0$  is this distance when the atoms are at equilibrium positions.  $\alpha_i$  is the angle  $B_i-A-B_j$  while  $\beta_i$  is the angle made by the line  $A-B_i$  with the  $B_3$  plane. The frequencies are then given by the equations:

$$\begin{aligned}m\lambda_1 - K - 2K' &= 0, \\ mM\lambda_2 - (3m + M)G &= 0, \\ m^2\lambda_3 - s^2 - \left\{\frac{1}{2}(K - K')(3m/M + 2) \right. \\ &\quad \left. + H(9m/2M + 3)\right\}m\lambda_3, \\ &\quad + H(K - K')(9m/M + 3) = 0.\end{aligned}$$

Using the vibrational data for the boron trihalides containing the isotope  $B^{11}$  we have determined the various force constants with the

<sup>10</sup> D. M. Yost, E. N. Lassettre, and S. T. Gross, J. Chem. Phys. **4**, 325 (1936).

<sup>11</sup> T. F. Anderson, J. Chem. Phys. **4**, 161 (1936).

<sup>12</sup> D. M. Dennison, Phil. Mag. **1**, 195 (1926); F. Lechner, Wien. Bericht **141**, 633 (1932); J. B. Howard and E. B. Wilson, Jr., J. Chem. Phys. **2**, 630 (1934); J. Rosenthal, Phys. Rev. **47**, 235 (1935). We have applied in part the results of Howard and Wilson in our treatment.

results indicated in Table IV. It is interesting to note that the constant  $G$ , usually assumed equal to  $H$  in the treatment of pyramidal type molecules, is nearly equal to  $H$  for heavy halogen atoms, but is considerably greater than  $H$  for light halogen atoms bound to the central boron.

Using these force constants we have calculated the frequencies  $\nu_3$  and  $\nu_4$  to be expected for the compounds containing the  $B^{10}$  isotope. The results agree well with the observed values as indicated in the table. The fact that the frequency  $\nu_4$  has little isotopic shift indicates that

the symmetry modes are very nearly the normal modes in these cases.

Finally in Table IV, columns 10 and 11 are presented values of the internuclear distances calculated directly from the values of  $K$  by Badger's rule.<sup>13</sup> The agreement with the internuclear distances determined by electron diffraction experiments<sup>4</sup> is much better than would have been expected, judging from the experience of previous investigators.

<sup>13</sup> R. M. Badger, *J. Chem. Phys.* **2**, 128 (1934); **3**, 710 (1935).

NOVEMBER, 1936

JOURNAL OF CHEMICAL PHYSICS

VOLUME 4

## The Entropy of Carbon Tetrachloride

R. C. LORD, JR. AND E. R. BLANCHARD, *Department of Chemistry, Johns Hopkins University, Baltimore, Maryland*  
(Received July 27, 1936)

A new calculation of the entropy of carbon tetrachloride has been made from Raman spectra and electron diffraction data. The effect of the isotopes of chlorine has been investigated. A new value of the calorimetric entropy has been computed from recent (and in part unpublished) thermal data. The agreement of the entropies is sufficiently good as to leave little reason to doubt the validity of the spectroscopic calculations

SOME time ago Yost and Blair<sup>1</sup> noted a difference of about three entropy units between the entropy of carbon tetrachloride calculated from the existing thermal data<sup>2</sup> and a value of the entropy which they calculated from spectroscopic data by statistical methods. Assuming that the thermal data were substantially correct, they suggested that the discrepancy arose in the statistical calculation from uncertainty about the vibrational states of the molecule. The purpose of their note was to point out the dubious value, in their opinion, of spectroscopic entropy calculations made on the basis of the customary assumption that molecular vibrations are quasi-harmonic. In view of the good spectroscopic data available and the straightforward interpretation of the vibrational structure of carbon tetrachloride, it seemed to us unlikely that uncertainties of the sort suggested by Yost and Blair could account for the discrepancy between the thermal and spectroscopic

entropies. Rather it appeared that the difficulty probably lay in the experimental thermal data. We have had put at our disposal some new unpublished heat capacity data for carbon tetrachloride. Furthermore we have made a new calculation of the statistical entropy, thoroughly investigating several possible sources of the disagreement. The theoretical and calorimetric data have been found to agree well within the limits of the probable experimental error in the latter.

### STATISTICAL ENTROPY

It appeared worth while to refine the statistical computations by taking into account the isotopes  $Cl^{35}$  and  $Cl^{37}$ . The presence of these isotopes creates five different isotopic configurations for carbon tetrachloride, and the final value of the entropy has been obtained from the entropies of the configurations. It is to be remarked that the symmetry numbers of the various configurations are not uniformly twelve. At first sight, this variance in symmetry number might be thought

<sup>1</sup> Yost and Blair, *J. Am. Chem. Soc.* **55**, 2610 (1933).

<sup>2</sup> Latimer, *J. Am. Chem. Soc.* **44**, 90 (1922).