

β -Nitro-5,10,15-tritolylicorroles

Manuela Stefanelli,[†] Giuseppe Pomarico,[†] Luca Tortora,[†] Sara Nardis,[†] Frank R. Fronczek,[‡] Gregory T. McCandless,[‡] Kevin M. Smith,^{*,‡} Machima Manowong,[§] Yuanyuan Fang,[§] Ping Chen,[§] Karl M. Kadish,^{*,§} Angela Rosa,^{*,[⊥]} Giampaolo Ricciardi,[⊥] and Roberto Paolesse^{*,†}

[†]Dipartimento di Scienze e Tecnologie Chimiche, Università di Roma Tor Vergata, 00133 Roma, Italy


[‡]Department of Chemistry, Louisiana State University, Baton Rouge, Louisiana 70803, United States

[§]Department of Chemistry, University of Houston, Houston, Texas 77204-5003, United States

[⊥]Dipartimento di Chimica, Università della Basilicata, 85100 Potenza, Italy

Supporting Information

ABSTRACT: Functionalization of the β -pyrrolic positions of the corrole macrocycle with $-\text{NO}_2$ groups is limited at present to metallocorrolates due to the instability exhibited by corrole free bases under oxidizing conditions. A careful choice of the oxidant can limit the transformation of corroles into decomposition products or isocorrole species, preserving the corrole aromaticity, and thus allowing the insertion of nitro groups onto the corrole framework. Here we report results obtained by reacting 5,10,15-tritolylicorrole (TTCorrH_3) with the $\text{AgNO}_2/\text{NaNO}_2$ system, to give mono- and dinitrocorrole derivatives when stoichiometry is carefully controlled. Reactions were found to be regioselective, affording the 3- $\text{NO}_2\text{TTCorrH}_3$ and 3,17- $(\text{NO}_2)_2\text{TTCorrH}_3$ isomers as the main products in the case of mono- and disubstitution, in 53 and 20% yields, respectively. In both cases, traces of other mono- and disubstituted isomers were detected, which were structurally characterized by X-ray crystallography. The influence of the β -nitro substituents on the corrole properties is studied in detail by UV–visible, electrochemical, and spectroelectrochemical characterization of these functionalized corroles. Density functional theory (DFT) and time-dependent DFT (TDDFT) calculations of the ground and excited state properties of these β -nitrocorrole derivatives also afforded significant information, closely matching the experimental observations. It is found that the β - NO_2 substituents conjugate with the π -aromatic system of the macrocycle, which initiates significant changes in both the spectroscopic and redox properties of the so functionalized corroles. This effect is more pronounced when the nitro group is introduced at the 2-position, because in this case the conjugation is, for steric reasons, more efficient than in the 3-nitro isomer.



■ INTRODUCTION


The establishment of efficient synthetic methodologies for the preparation of corrole^{1–4} has allowed a more detailed investigation of its chelating properties as a ligand, and of its unique behavior in several reactions.⁵ The investigation of corrole functionalization reactions is fundamental for the potential exploitation of this macrocycle in practical uses, since the manipulation of the peripheral positions can deeply affect the corrole properties, as shown by the first seminal examples reported in the literature.⁶

Compared with porphyrins, the peripheral functionalization of the corrole macrocycle is possible only in a limited number of cases, but recently some successful substitutions have been reported in the literature.^{2,7} In particular we have been interested in the nitration reaction, because the insertion of one or more $-\text{NO}_2$ groups onto the β -pyrrole positions represents one of the most valuable modifications for researchers interested in functionalization of the corrole skeleton; for example, the potential of the nitro group in

activating nucleophilic substitution on the adjacent β -carbons of the pyrrole rings is well-known. Furthermore, it is also interesting to evaluate the influence of the strong electron withdrawing character of the nitro group on the behavior of the corrole ring, which is characterized by high electron density and its noninnocent character as ligand.⁸ In this connection, we recently reported the first successful example of a vicarious nucleophilic substitution performed on corrole nitro-derivatives by reacting copper and germanium β -(NO_2) corrolates with 4-amino-4H-1,2,4-triazole.⁹ The reaction afforded β -amino- β -nitro derivatives, which can have great synthetic value for the elaboration of new chromophores, as well as extended aromatic β -fused architectures. To date β -nitrocorrole derivatives of copper,⁹ germanium,¹⁰ gallium,¹¹ silver,¹² cobalt,¹³ and iron¹⁴ have been prepared by using a variety of nitrating systems and reaction conditions, while all attempts to nitrate the corrole

Received: April 17, 2012



Chart 1. Molecular Structures of β -Nitrocorroles Presented in This Work

free-base failed, mainly because of the lability of corroles under these substitution conditions. This was evidenced under a variety of reaction conditions with the formation of isocorrole species. The recent development of demetalation protocols for the effective removal of silver¹⁵ and copper¹⁶ from the corresponding corrole complexes has allowed the preparation of nitrocorrole free bases from the application of the reductive DBU/THF system to a silver 3-NO₂ corrolate (3-NO₂TtBuCorrAg), obtained by using a large excess of AgNO₂.¹⁵ Although this approach allowed circumvention of the instability of corrole under the nitrating systems tested to date, it opened the way to preparation of otherwise unavailable 3-NO₂ corroles. Such preparations of functionalized free bases in a single step is highly desirable, particularly in the preparation of polynitrated free bases. Prompted by the recent results obtained in the case of nitration of copper corrole complexes⁹ carried out with an optimized ratio of silver salt and sodium nitrite mixture as the nitrating system, we decided to apply the same reaction conditions to corrole free bases, surmising that minimizing the oxidant stoichiometry could preserve corrole aromaticity during the reaction, but still allow nucleophilic attack by the NO₂[−] ion upon the macrocycle.

We report here the first successful example of nitration of the corrole free base, obtained by reacting 5,10,15-tritolylcorrole (TTCorrH₃) with the AgNO₂/NaNO₂ system. Variation of the molar ratio of reagents allowed us to obtain 3-nitro- and 3,17-dinitro-derivatives of corrole in satisfying yield, together with traces of other isomers, namely 2-NO₂-, 2,3-(NO₂)₂-, and 3,12-(NO₂)₂ corroles (Chart 1). While the substitution pattern of 2-NO₂TTCorrH₃ was established structurally by X-ray crystallography, in the case of 2,3-(NO₂)₂TTCorrH₃ and 3,12-(NO₂)₂TTCorrH₃ isomers, the substitution pattern was unambiguously confirmed by preparation of the corresponding Co triphenylphosphine complexes, which were also structurally characterized.

The definition of the synthetic protocols for the preparation of free base nitrocorroles prompted us to study in detail the influence of the introduction of the nitro group at the corrole β -positions. This has been accomplished through a detailed UV–visible, electrochemical, and spectroelectrochemical characterization of these functionalized corroles. The changes in the UV–visible spectra and in the first reduction potential following β -nitration have been rationalized by density functional theory (DFT) and time-dependent DFT (TDDFT) calculations of the ground and excited states of the TTCorrH₃, 2-NO₂TTCorrH₃, 3-NO₂TTCorrH₃, and 3,17-(NO₂)₂TTCorrH₃ series of free base corroles. Taken together, the experimental and theoretical

results provide informative details on the behavior of these macrocycles.

EXPERIMENTAL SECTION

General. Silica gel 60 (70–230 mesh, Sigma Aldrich) was used for column chromatography. Reagents and solvents (Aldrich, Merck or Fluka) were of the highest grade available and were used without further purification, except for benzonitrile (PhCN), which was distilled over P₂O₅ under vacuum prior to use, and tetra-*n*-butylammonium perchlorate (TBAP), which was dried under vacuum at 30 °C for at least one week prior to use. ¹H NMR spectra were recorded on a Bruker AV300 (300 MHz) spectrometer. Chemical shifts are given in parts per million relative to residual CHCl₃ (7.25 ppm). UV–visible spectra were measured on a Cary 50 spectrophotometer. Mass spectra (FAB mode) were recorded on a VGQuattro spectrometer in the positive-ion mode using *m*-nitrobenzyl alcohol (Aldrich) as a matrix.

Diffraction data for chloroform solvates of 2-NO₂TTCorrH₃ and 2,3-(NO₂)₂TTCorrCoPPh₃ and for a dichloromethane solvate of 3,12-(NO₂)₂TTCorrCoPPh₃ were collected at low temperature on a Bruker Kappa Apex-II CCD diffractometer with either Cu K α (λ = 1.54184 Å) or MoK α (λ = 0.71073 Å) radiation and an Oxford Cryosystems Cryostream chiller. Refinement was by full-matrix least-squares using SHELXL,¹⁷ with H atoms in idealized positions, except for those on N, for which coordinates were refined. Absorption corrections were applied with a multiscan method (Bruker SADABS¹⁸).

Synthesis of β -Nitrocorroles. Preparation of 3-(NO₂)TTCorrH₃. NaNO₂ (75 mg, 1.1 mmol) and AgNO₂ (19 mg, 0.12 mmol) were added to a refluxing solution of TTCorrH₃ (70 mg, 0.12 mmol) in DMF (12 mL), and the progress of the reaction was followed by TLC analysis and UV–visible spectroscopy. The reaction was complete in 10 min, as indicated by the UV–visible spectrum of the reaction mixture that showed the spectral features of the desired product. The reaction mixture was cooled, quenched by addition of few drops of aqueous hydrazine solution, precipitated by adding distilled water, and then filtered. The crude product was taken up in CH₂Cl₂ and applied to a silica gel column for chromatographic purification. Elution with CH₂Cl₂/hexane (7:3) afforded traces of the silver complex, TTCorrAg, as a first reddish band followed by a second fraction containing a complex mixture of green and brownish compounds that were not easy to separate using a column. A subsequent main green fraction was isolated and crystallized from CH₂Cl₂/MeOH giving the 3-nitroderivative as an emerald green powder (38 mg, 52% yield). The spectroscopic characterization of 3-NO₂TTCorrH₃ was fully in agreement with data in the literature.¹⁶

The second fraction was subjected to preparative TLC on silica gel plates (CH₂Cl₂/hexane 1:1 as eluant). A green fraction having 0.52 R_f was isolated and identified as 2-NO₂TTCorrH₃.

2-NO₂TTCorrH₃. Mp > 300 °C. UV–visible (CHCl₃): λ_{max} nm (log ϵ) 408 (4.52), 460 (4.53), 588 (4.33), 715 (4.31). ¹H NMR (300 MHz, CDCl₃): 9.13 (d, 1H, *J* = 4.20 Hz β -pyrrole), 9.08 (s, 1H, β -pyrrole), 8.74 (d, 1H, *J* = 4.83 Hz β -pyrrole), 8.64 (d, 1H, *J* = 4.89 Hz

β -pyrrole), 8.34 (d, 1H, $J = 4.95$ Hz β -pyrrole), 8.28 (m, 3H, β -pyrrole + phenyls), 8.03 (d, 2H, $J = 7.92$ Hz phenyls), 7.95 (d, 2H, $J = 7.83$ Hz phenyls), 7.67 (d, 2H, $J = 7.86$ Hz phenyls), 7.59 (m, 6H phenyls), 2.70 (s, 3H, p -CH₃), 2.67 (s, 6H, p -CH₃). Anal. calcd for C₄₀H₃₁N₅O₂: C, 78.2; H, 5.1; N, 11.3%. Found: C, 78.3; H, 5.1; N, 11.4%. MS (FAB): m/z 613(M⁺). Crystal data: C₄₀H₃₁N₅O₂ · 1.47 CHCl₃, triclinic space group $P\bar{1}$, $a = 9.6282(10)$, $b = 14.5927(14)$, $c = 15.2094(15)$ Å, $\alpha = 64.348(5)$, $\beta = 83.229(5)$, $\gamma = 87.496(5)^\circ$, $V = 1912.8(3)$ Å³, $T = 90.0(5)$ K, $Z = 2$, $\rho_{\text{calcd}} = 1.370$ g cm⁻³, $\mu(\text{Cu K}\alpha) = 3.42$ mm⁻¹. A total of 19 520 data was collected at to $\theta = 68.7^\circ$. $R = 0.059$ for 6350 data with $I > 2\sigma(I)$ of 6674 unique data and 510 refined parameters, CCDC 847685.

Preparation of 3,17-(NO₂)₂TTCorrH₃. TTCorrH₃ (100 mg, 0.18 mmol) was dissolved in DMF (15 mL), and the solution was heated at reflux. Then, NaNO₂ (99 mg, 1.44 mmol) and AgNO₂ (55 mg, 0.36 mmol) were added and the progress of the reaction was followed by TLC analysis and UV–visible spectroscopy. The reaction was complete in 30 min. The reaction mixture was cooled, quenched by addition of a few drops of aqueous hydrazine solution, precipitated by adding distilled water, and then filtered. The crude product was taken up in CH₂Cl₂ and applied to a silica gel column for chromatographic purification. Elution with CH₂Cl₂ afforded the isocorrolo 3-(NO₂)-5-(OH)TTisoCorH₂ (10 mg, 9% yield) as the first fraction, followed by two green bands corresponding to the dinitroisomers 2,3-(NO₂)₂TTCorrH₃ (2 mg, 1.7% yield) and 3,12-(NO₂)₂TTCorrH₃ (1 mg, 0.8% yield), respectively. The last brilliant green fraction was collected and crystallized from CH₂Cl₂/*n*-hexane giving the 3,17-(NO₂)₂TTCorrH₃ as a brilliant green powder (24 mg, 20% yield).

3,17-(NO₂)₂-TTCorrH₃. Mp > 300 °C. UV–visible (CHCl₃): λ_{max} nm (log ϵ) 439 (4.54), 483 (4.63), 707 (4.52). ¹H NMR (300 MHz, CDCl₃): 8.94 (s, 2H, β -pyrrole), 8.52 (d, 2H, $J = 4.80$ Hz β -pyrrole), 8.23 (d, 2H, $J = 4.86$ Hz β -pyrrole), 7.95 (m, 6H, phenyls), 7.55 (m, 6H, phenyls), 2.68 (s, 3H, p -CH₃), 2.63 (s, 6H, p -CH₃). Anal. calcd for C₄₀H₃₀N₆O₄: C, 72.9; H, 4.6; N, 12.8%. Found: C, 72.8; H, 4.6; N, 12.7%. MS (FAB): m/z 658 (M⁺).

Preparation of 2,3-(NO₂)₂-TTCorrCoPPh₃. 2,3-(NO₂)₂-TTCorrH₃ (5 mg, 0.08 mmol) in CH₂Cl₂ (10 mL) and cobalt(II) acetate (60 mg, 0.24 mmol) and triphenylphosphine (63 mg, 0.24 mmol) in CH₃OH (15 mL) were mixed, and the resulting solution was heated at reflux for 1 h. The solvent was evaporated and the residue dissolved in CH₂Cl₂ and filtered through a silica plug to afford the complex as a green-bluish fraction. Crystallization from CH₂Cl₂/CH₃OH afforded the title compound in an almost quantitative yield.

2,3-(NO₂)₂-TTCorrCoPPh₃. Mp > 300 °C. UV–visible (CHCl₃): λ_{max} nm (log ϵ) 370 (3.92), 552 (3.47), ¹H NMR (300 MHz, CDCl₃): 9.06 (d, 1H, $J = 4.50$ Hz β -pyrrole), 8.35 (d, 1H, $J = 4.89$ Hz β -pyrrole), 8.15 (d, 1H, $J = 4.89$ Hz β -pyrrole), 8.10 (d, 1H, $J = 5.07$ Hz β -pyrrole), 8.02 (d, 1H, $J = 5.07$ Hz β -pyrrole), 7.98 (d, 1H, $J = 4.47$ Hz β -pyrrole), 7.58–7.36 (m, 15 H, phenyls), 7.22 (m, 3 H, *p*-phosphine), 6.94 (m, 6 H, *m*-phosphine), 5.43 (m, 6 H, *o*-phosphine), 2.60 (s, 3H, p -CH₃), 2.59 (s, 6H, p -CH₃). Anal. calcd for C₅₈H₄₂CoN₆O₄P: C, 71.31; H, 4.33; N, 8.60%. Found: C, 71.28; H, 4.36; N, 8.63%. MS (FAB): m/z 715 (M⁺ – PPh₃). Crystal data: C₅₈H₄₂CoN₆O₄P · 0.81 CHCl₃, orthorhombic space group $Fdd2$, $a = 37.190(3)$, $b = 37.996(6)$, $c = 13.8041(10)$ Å, $V = 19,506(4)$ Å³, $T = 90.0(5)$ K, $Z = 16$, $D_x = 1.463$ g cm⁻³, $\mu(\text{Mo K}\alpha) = 0.58$ mm⁻¹. A total of 36 267 data was collected to $\theta_{\text{max}} = 30.0^\circ$. $R = 0.059$ for 10 933 data with $I > 2\sigma(I)$ of 12 276 unique data and 634 refined parameters, Flack¹⁹ parameter 0.022(12). Disordered solvent contribution was removed using the SQUEEZE procedure,²⁰ CCDC 869740

Preparation of 3,12-(NO₂)₂-TTCorrCoPPh₃. 3,12-(NO₂)₂-TTCorrCoPPh₃ was prepared as reported above for the 2,3-dinitro isomer.

3,12-(NO₂)₂-TTCorrCoPPh₃. Mp > 300 °C. UV–visible (CHCl₃): λ_{max} nm (log ϵ) 450 (4.31), 594 (4.08), ¹H NMR (300 MHz, CDCl₃): 8.98 (s, 1H, β -pyrrole), 8.67 (s, 1H, β -pyrrole), 8.53 (d, 2H, $J = 4.65$ Hz β -pyrrole), 8.30 (d, 2H, $J = 5.04$ Hz β -pyrrole), 8.21 (d, 2H, $J = 5.13$ Hz β -pyrrole), 8.07 (d, 2H, $J = 4.53$ Hz β -pyrrole), 7.66–7.32 (m, 15 H, phenyls), 7.28 (m, 3 H, *p*-phosphine), 7.18 (m, 6 H, *m*-phosphine), 6.53 (m, 6 H, *o*-phosphine), 2.62 (s, 3H, p -CH₃), 2.57 (s, 3H, p -CH₃), 2.54 (s, 3H, p -CH₃). Anal. calcd for C₅₈H₄₂CoN₆O₄P: C,

71.31; H, 4.33; N, 8.60%. Found: C, 71.27; H, 4.35; N, 8.64%. MS (FAB): m/z 715 (M⁺ – PPh₃). Crystal data: C₅₈H₄₂CoN₆O₄P · 0.5 CH₂Cl₂, monoclinic space group $P2_1/c$, $a = 18.225(3)$, $b = 9.2487(14)$, $c = 29.294(4)$ Å, $\beta = 105.212(10)^\circ$, $V = 4764.8(13)$ Å³, $T = 120.0(5)$ K, $Z = 4$, $D_x = 1.421$ g cm⁻³, $\mu(\text{Cu K}\alpha) = 4.12$ mm⁻¹. A total of 19 491 data was collected to $\theta_{\text{max}} = 64.9^\circ$, $R = 0.059$ for 6277 reflections with $I > 2\sigma(I)$ of 7795 unique data and 644 refined parameters. Disordered CH₂Cl₂ was modeled as a half-populated site near a 2-fold axis, CCDC 869741.


Electrochemical and Spectroelectrochemical Measurements. Cyclic voltammetry was carried out with an EG&G model 173 potentiostat/galvanostat. A homemade three-electrode electrochemistry cell was used and consisted of a glassy carbon working electrode, a platinum wire counter electrode, and a saturated calomel reference electrode (SCE). The SCE was separated from the bulk of the solution by a fritted-glass bridge of low porosity which contained the solvent/supporting electrolyte mixture. All potentials are referenced to the SCE.

UV–visible spectroelectrochemical experiments were performed with an optically transparent platinum thin-layer electrode of the type described in the literature.²¹ Potentials were applied with an EG&G Model 173 potentiostat/galvanostat. Time-resolved UV–visible spectra were recorded with a Hewlett-Packard Model 8453 diode array rapid-scanning spectrophotometer.

Determination of Equilibrium Constants of Free Base Corroles. In order to study the protonation or deprotonation process, microliter quantities of TFA/CH₂Cl₂ and piperidine/CH₂Cl₂ were added gradually to 6.0 mL CH₂Cl₂ solution of TTCorrH₃ in a 1.0 cm homemade glass cell. Solutions of the corroles were freshly degassed by N₂ flux before each measurement and the spectra recorded after each addition of titration reagents. The concentration of the corrole solutions were in the 10⁻⁶ M range, in order to avoid aggregation of the macrocycle.

The free base is protonated in acid and deprotonated in basic solutions. The spectroscopic data were analyzed by the Hill equation, which gives the equilibrium constant (K) for the reactions.

Quantum Chemical Calculations. All calculations were performed with Turbomole V6.3 2011²² using both pure (BP86)²³ and hybrid (B3LYP)²⁴ functionals in combination with the extensively polarized basis sets of triple- ζ quality including high angular momentum polarization functions (def2-TZVPP).²⁵ In the BP86 calculations, the resolution of the identity (density fitting) approach²⁶ was used to save computer time in combination with the def2-TZVPP/J auxiliary basis set.²⁷ The ground state molecular structures of TTCorrH₃, 2-NO₂TTCorrH₃, 3-NO₂TTCorrH₃, 3,17-(NO₂)₂-TTCorrH₃, and related monoanionic species were fully optimized (without constraints) in the gas-phase and in dichloromethane solution. For the optimization of the monoanions, the unrestricted formalism was used. Solvation effects on the optimized structures were modeled through a dielectric continuum model, which was chosen to be the COSMO model.²⁸ The optimized structures were confirmed to be local minima by calculating the harmonic vibrational frequencies. The thermodynamic parameters for the one-electron reduction reaction of the investigated metal-free corroles were computed at the BP86 level, at the geometries optimized in dichloromethane solution. The zero-point energies (ZPE), thermal corrections, and entropy terms for the optimized geometries of the neutral and monoanionic species were obtained from frequency calculations in dichloromethane. Vertical absorption energies and oscillator strengths of the lowest singlet excited states of TTCorrH₃ and its nitro derivatives were computed in dichloromethane solution at TDDFT²⁹ level using the hybrid B3LYP²⁴ functional and the DFT/B3LYP ground state geometries optimized in dichloromethane. The calculated excitation energies contain, apart from the altered “solvated” orbitals (slow term), also the contributions from the “fast” solvent response term.³⁰

Scheme 1. Synthesis of $(\text{NO}_2)_x\text{TTCorrH}_3$ 

RESULTS AND DISCUSSION

Synthesis. In the past few years, we have been particularly interested in the elaboration of synthetic protocols for the synthesis of β -nitrocorrole derivatives. By using different nitrating systems, we have been able to establish diverse methodologies for the convenient preparation of mono- and dinitro-metallocorrolates, starting in most cases from metallocorroles, such as the Cu, Ge, and Fe derivatives. Our attempts to introduce nitro groups directly onto the free base, to obtain the corresponding functionalized corrole failed, since the required β -substitution took place in combination with other structural modifications, thereby precluding the isolation of the pure nitrocorrole free base. For example, using a large excess of AgNO_2 , both functionalization and metalation of the macrocycle occurred simultaneously, affording the corresponding Ag(III) 3-nitrocorrole as the reaction product.¹² In the case of TFA/NaNO_2 or HCl/NaNO_2 systems, the formation of β -nitroisocorroles was observed, but these were converted in a second step into the corresponding aromatic corrole complexes by cobalt insertion.¹³ Our hypothesis for the nitration reaction mechanism is that the first step requires an oxidant species for the generation of the corrole π -cation radical, which is then attacked by nitrite to afford the final product. However, corrole free bases demonstrate extreme sensitivity under oxidizing reaction conditions, leading to a quite facile conversion of the macrocycle into the corresponding isocorrole species.^{9,13,15,31–35} Therefore, the nitration reaction conditions made the preparation of aromatic nitro-compounds difficult, instead favoring the formation of isocorrole derivatives through nucleophile insertion onto a *meso*-carbon position, which was observed for the 3- NO_2 -5-OH-isocorroles.^{9,31} Since the oxidation step is crucial for the reaction outcome, we surmised

that a careful choice of the oxidant species could potentially drive the reaction pathway toward nitrocorroles, inhibiting the further transformation to isocorroles.

We recently reported that when the nitration reaction is performed on corrole using an equimolar amount of an oxidizing agent, such as chloranil in combination with an excess of NaNO_2 , the 3- NO_2 -5-OH-isocorrole is rapidly obtained as the main reaction product, confirming the regioselectivity of both mononitration and the formation of the single 5-substituted isocorrole isomer.³¹ Our attention turned toward other oxidants, and prompted by the results reported recently for nitration of copper corrolates, we tested the $\text{AgNO}_2/\text{NaNO}_2$ system, which allows the preparation of mono- and dinitro-derivatives in good yields just by varying the reagent stoichiometry.⁹

Thus, TTCorrH_3 was reacted in refluxing DMF with a mixture of sodium and silver nitrites using a corrole/ $\text{AgNO}_2/\text{NaNO}_2$ molar ratio of 1:1:9 (Scheme 1) with the intention of achieving mononitration of the corrole. The reaction proceeded very fast and in 10 min complete disappearance of the starting compound was observed by TLC analysis, together with the formation of a green major product. The UV–visible spectrum of the mixture changed rapidly even from the beginning of the reaction and showed at the end the typical spectral features of 3- $\text{NO}_2\text{TTCorrH}_3$,¹⁵ indicating it as the main product. Chromatographic purification afforded traces of the silver complex of the corrole as the first rose-red band together with a second fraction containing a mixture of brownish and greenish products, inseparable on the column. A third green fraction was the major reaction product and corresponded to 3- $\text{NO}_2\text{TTCorrH}_3$, obtained in 52% yield. It is worth mentioning that the formation of the 3-nitro-5-hydroxy-isocorrole was not

observed, showing that the amount of silver nitrite was correctly chosen to allow the β -functionalization, and to preserve at the same time the integrity of the nitrocorrole free base formed in the reaction medium. The second band was further purified on a preparative silica gel TLC plate, using a mixture of CH_2Cl_2 /hexane as eluant. A green band ($R_f = 0.52$) was isolated in sufficient amount to be characterized using the usual spectroscopic techniques. This fraction showed an unusual UV–visible spectrum, characterized by a split Soret band (of similar intensity) at 408 and 460 nm, a less intense narrow band centered at 588 nm, and a broad band at about 715 nm. The FAB mass spectrum of this compound afforded a molecular peak at m/z 613, which is consistent with a mononitrocorrole. The ^1H NMR spectrum and integral calculations confirmed it as a monosubstituted corrole showing a proton singlet at 9.08 ppm. It is noteworthy that the pattern of substitution is clearly different from that of the known 3- $\text{NO}_2\text{TTCorrH}_3$, showing a pyrrolic doublet more deshielded compared with the singlet, just as observed for the 2-bromocorrole isomer obtained by reacting the TTCorrH_3 with EtMgBr .³³ The definitive identification of position C2 as the site of substitution for this compound was afforded by X-ray crystallographic analysis, carried out on a single crystal obtained by slow diffusion of *n*-hexane into a dilute dichloromethane solution. This unambiguously confirmed the new monosubstituted corrole as the 2- $\text{NO}_2\text{TTCorrH}_3$ isomer (Figure 1).

The three protonated N atoms in the interior of the molecule force marked deviations of the corrole system from coplanarity. The mean deviation of its 23 atoms is 0.159 Å, with a maximum of 0.415(4) Å. The nitrated “A” pyrrole and the adjacent “B” pyrrole not directly bonded to it come nearest to being




Figure 1. Molecular structure of 2- $\text{NO}_2\text{TTCorrH}_3$, with 50% ellipsoids.

coplanar, forming a dihedral angle of $10.5(6)^\circ$. The main distortion of the corrole is a twist, in which the other two pyrroles “C” and “D” tip alternately above and below this plane, forming dihedral angles of $18.1(3)^\circ$ and $20.6(2)^\circ$ with it and $23.8(3)^\circ$ with each other. The N atoms of the two alternately tipped pyrroles are somewhat pyramidal, with their H atoms lying $0.14(4)$ and $-0.30(4)$ Å out of the best planes of their respective pyrroles. The third N–H hydrogen atom is not tipped out of plane, but forms an N–H \cdots N hydrogen bond (N \cdots N 2.623(3) Å; N–H \cdots N $134(3)^\circ$). The nitro group lies essentially in the plane of the pyrrole to which it is bonded, with a C1–C2–N–O torsion angle $1.2(4)^\circ$. The two directly bonded pyrroles exhibit a N–C1–C19–N twist of $7.6(3)^\circ$.

With the aim of driving the reaction toward dinitrocorrole derivatives, we modified the amounts of nitrating agents by altering the corrole/ AgNO_2 molar ratio to 1:2 (Scheme 1). Of course, in this case, we were aware that the increase of the oxidant reagent could induce the formation of the isocorrole, instead of forming dinitrocorroles. The reaction of TTCorrH_3 with silver and sodium nitrites in a 1:2:8 molar ratio, afforded in 30 min a mixture of compounds observable from the TLC analysis as green and bluish-green bands together with a number of decomposition products.

Chromatographic purification on silica gel using dichloromethane as eluant afforded, after elution of small traces of the silver tritolylcorrolate, a first major fraction consisting of the 3- NO_2 -5-OH-TTisoCorrH₂, obtained in 9% yield. Afterward, a small quantity of two green fractions was isolated and subjected to spectroscopic identification. The band having the greater R_f value showed a UV–visible spectral profile quite similar to 3- $\text{NO}_2\text{TTCorrH}_3$, but generally red-shifted, while the second fraction displayed a strongly red-shifted Soret band at 472 nm together with two satellite bands at 613 and 666 nm. The mass spectrometry analysis afforded a molecular peak at m/z 658 for both compounds, tentatively identifying them as two different dinitrocorrole isomers. The proton NMR spectra of both compounds were of poor resolution, hampering definitive identification of the sites of substitution by the nitro groups.

The last green fraction obtained by column chromatographic purification was the main reaction product, obtained in 20% yield, and fully characterized as 3,17- $(\text{NO}_2)_2\text{TTCorrH}_3$. The UV–visible spectrum was unusual, having a split Soret band at 439 and 483 nm and an intense broad absorption at ca. 700 nm.

To allow a more definitive identification of the other dinitrocorrole isomers, we decided to prepare the corresponding Co complexes, by reaction of these products with $\text{Co}(\text{OAc})_2$ and PPh_3 ; the reaction provided cobalt complexes having well resolved ^1H NMR spectra. In the case of the first compound, the lack of proton singlets together with the presence of a downfield shifted doublet above 9 ppm, clearly separated from the other five pyrrolic resonances at ca. 8–8.5 ppm, led to the conclusion that for this compound the substitution occurred on the C2 and C3 positions of the corrole macrocycle. We were able to obtain crystals of this isomer suitable for X-ray crystallographic analysis, which helped to elucidate the structure of the corresponding free base as 2,3- $(\text{NO}_2)_2\text{TTCorrH}_3$.

The structure of 2,3- $(\text{NO}_2)_2\text{TTCorrCoPPh}_3$ is shown in Figure 2. The Co atom has square pyramidal coordination with Co–N distances in the range 1.869(3)–1.886(3) Å (mean 1.880 Å), Co–P distance 2.2321(8) Å, and the Co is 0.2634(13) Å out of the best plane of the four pyrrole N atoms. The 23-atom C_{19}N_4 corrole core deviates only slightly




Figure 2. Molecular structure of 2,3-(NO₂)₂TTCorrCoPPh₃, with 50% ellipsoids and H atoms not shown.

from planarity, with an rms deviation 0.059 Å and maximum deviation 0.119(3) Å for C7. The small distortion from planarity is a slight saddle. One nitro group lies nearly in the corrole best plane, with C1–C2–N–O torsion angle $-6.9(5)^\circ$, while the other is more nearly orthogonal to the corrole, with C2–C3–N–O torsion angle $-73.1(4)^\circ$.

The second Co complex showed two proton singlets at 8.98 and 8.67 ppm, respectively, and three different proton signals of equal intensity at 2.62, 2.57, and 2.54 ppm, indicating it to be an asymmetric product. Also in this case, the X-ray crystallographic characterization (Figure 3) led to identification of 3,12-




Figure 3. Molecular structure of 3,12-(NO₂)₂TTCorrCoPPh₃, with 30% ellipsoids and H atoms not shown.

(NO₂)₂TTCorrCoPPh₃, confirming the corresponding corrole free base as the 3,12-(NO₂)₂TTCorrH₃ isomer. It is interesting to note that this isomer is, to the best of our knowledge, the first example of a disubstituted corrole where pyrroles A and C are functionalized, instead the usual A and D pattern.

The structure of 3,12-(NO₂)₂TTCorrCoPPh₃ is illustrated in Figure 3 and is quite similar to that of 2,3-(NO₂)₂TTCorrCoPPh₃, with a somewhat more nonplanar corrole core. The Co atom has square pyramidal coordination with Co–N distances in the range 1.873(3)–1.892(3) Å (mean 1.878 Å), Co–P distance 2.2113(11) Å, and the Co is 0.2734(15) Å out of the best plane of the four pyrrole N atoms. The 23-atom C₁₉N₄ corrole core is approximately three times

more distorted from planarity than that of 2,3-(NO₂)₂TTCorrCoPPh₃, having an rms deviation of 0.159 Å and maximum deviation 0.339(4) Å for C8. The distortion is best described as a saddle, although all four N atoms lie on the same side of the 23-atom best plane by a mean distance of 0.118 Å. The two nitro groups are twisted out of the corrole best plane by similar amounts, forming torsion angles: C4–C3–N–O, $-40.0(6)^\circ$ and C11–C12–N–O, $29.2(5)^\circ$.

It is worth mentioning that in all cases the use of the AgNO₂/NaNO₂ system allowed the β -functionalization of the macrocycle, without inducing the concomitant metalation, which occurs when the reaction is carried out with an excess of silver nitrite. In this case the AgNO₂ acts only as oxidant, and the proper amount chosen prevented the metalation of the macrocycle. The transient formation of the silver corrole complex, then demetalated by the final addition of hydrazine can be safely excluded, because hydrazine is unable to promote silver corrole demetalation and because the same products were again obtained when the nitration reaction was carried out avoiding the addition of this reagent.

Electrochemistry and Spectroelectrochemistry. The redox behavior of free base corroles is complicated in nonaqueous media by the facile gain or loss of protons, which can give in solution a mixture of the neutral corrole along with either its protonated [(Corr)H₄]⁺ or deprotonated [(Corr)H₂][–] form. To investigate the effect of nitration and the solvent on both the overall oxidation/reduction mechanism and the electrochemically initiated protonation/deprotonation process, a series of three corroles, TTCorrH₃, 3-NO₂TTCorrH₃, and 3,17-(NO₂)₂TTCorrH₃, were selected for electrochemical and spectroelectrochemical characterization in pyridine, PhCN, and CH₂Cl₂.

We earlier reported the electrochemistry and spectroelectrochemistry of several *meso*-substituted corroles in PhCN and pyridine, including TTCorrH₃.³⁶ It was demonstrated that the electroactive species in PhCN was the neutral corrole CorrH₃, while in pyridine, the monoanion [(Corr)H₂][–] is the prevalent species,³⁶ the deprotonation process being almost complete. A similar solvent effect on deprotonation is seen for the currently investigated corroles, as seen by the cyclic voltammograms in Figure 4 for TTCorrH₃ in CH₂Cl₂, PhCN and pyridine.

The first reduction of TTCorrH₃ is irreversible in all three solvents and located at $E_p = -1.33$ to -1.40 V for a scan rate of 0.1 V/s. Only a residual amount of neutral CorrH₃ is present in the basic pyridine solvent and a small peak current is seen for reduction of the fully protonated corrole. This contrasts with CH₂Cl₂, where CorrH₃ is the main electroactive species in solution and the peak current is much higher for the first reduction. Sweeping the potential from -0.40 to values beyond the first reduction in PhCN or pyridine drives to completion the deprotonation of TTCorrH₃, after which the deprotonated [(TTCorr)H₂][–] product undergoes a reversible one-electron ring-reduction, which is located at $E_{1/2} = -1.90$ V in PhCN and -1.88 V in pyridine.

As indicated above, TTCorrH₃ undergoes deprotonation in both CH₂Cl₂ and PhCN after the first one-electron reduction and this then generates [(Corr)H₂][–]. As a result, when reversing the scan at potential negative of the first reduction and then scanning in a positive direction, a new redox couple assigned to [(Corr)H₂][–]/[(•Corr)H₂] can be seen in CH₂Cl₂ and PhCN. This process is located at $E_{1/2} = -0.09$ to -0.08 V and it is not observed in CH₂Cl₂ or PhCN, when sweeping the potential toward more positive values from the initial potential




Figure 4. Cyclic voltammograms of TTCorrH₃ in (a) CH₂Cl₂, (b) PhCN, and (c) pyridine containing 0.1 M TBAP.

of -0.4 V. This oxidation of $[(\text{Corr})\text{H}_2]^-$ is however seen in pyridine on all scans, in a positive or negative direction ($E_{1/2} = 0.09$ V).

To investigate the effect of nitration on both the overall oxidation/reduction mechanism and the protonation/deprotonation processes, 3-NO₂TTCorrH₃ and 3,17-(NO₂)₂TTCorrH₃ were also characterized electrochemically and spectroelectrochemically in pyridine, PhCN, and CH₂Cl₂.

The addition of one or two nitro substituents to the β -pyrrole positions of the macrocycle leads to a positive shift in $E_{1/2}$ for each oxidation or reduction of the nitro-substituted free base corroles, as compared to measured half wave potentials for the oxidation or reduction of TTCorrH₃. This is as expected, due to the strong electron-withdrawing character of the NO₂ substituents, with the magnitude of the positive shift in $E_{1/2}$ varying as a function of both the number of the nitro groups on the macrocycle (0, 1, or 2), as well as the specific electrode reaction.

A summary of redox potentials for of TTCorrH₃, 3-NO₂TTCorrH₃, and 3,17-(NO₂)₂TTCorrH₃ in the different

solvents is given in Table 1, while typical cyclic voltammograms are shown in Figure 5 for the three corroles in pyridine,




Figure 5. Cyclic voltammograms of TTCorrH₃, 3-NO₂TTCorrH₃, and 3,17-(NO₂)₂TTCorrH₃ in pyridine containing 0.1 M TBAP.

containing 0.1 M TBAP. As shown in this figure, the addition of one or two nitro substituents to the β -pyrrole positions of the macrocycle leads to similar positive shifts in $E_{1/2}$ for the first two oxidations. A positive shift in $E_{1/2}$ of 130–160 mV for each additional nitro group is at the 3 and 17 positions of the macrocycle. For example, the $[(\text{Corr})\text{H}_2]^-/[(\text{Corr})\text{H}_2]$ reaction in pyridine is located at $E_{1/2} = 0.09$ V for TTCorrH₃ (with no NO₂ group), while 3-NO₂TTCorrH₃ (with one NO₂ group) is oxidized at $E_{1/2} = 0.25$ V and 3,17-(NO₂)₂TTCorrH₃ (with two NO₂ groups) is oxidized at $E_{1/2} = 0.41$ V. It is also worth mentioning that a much larger substituent effect of the nitro group is observed for the reductions than for the oxidations. For example, a 540 mV shift in $E_{1/2}$ for the second reduction is seen upon going from TTCorrH₃ to 3-NO₂TTCorrH₃. Although the magnitude of the shift in $E_{1/2}$ for the second and third reductions is reduced to 290 and 310 mV respectively, upon going from 3-NO₂TTCorrH₃ to 3,17-

Table 1. Half-Wave or Peak Potentials (V vs SCE) for the Different Forms of (NO₂)_xTTCorrH₃ in Solvents Containing 0.1 M TBAP

solvent	cpd	ox		red		
		second	first	first	second	third
pyridine	TTCorrH ₃	0.56 ^a	0.09	−1.33 ^a	−1.88	
	3-NO ₂ TTCorrH ₃	0.70 ^a	0.25	−0.87 ^a	−1.34	−1.71
	3,17-(NO ₂) ₂ TTCorrH ₃	0.83 ^a	0.41	−0.75 ^a	−1.05	−1.40
CH ₂ Cl ₂	TTCorrH ₃		0.33	−1.40 ^a		
	3-NO ₂ TTCorrH ₃		0.50 ^a	−0.84 ^a	−1.42 ^a	−1.60 ^a
	3,17-(NO ₂) ₂ TTCorrH ₃			−0.62 ^a	−1.06	−1.39
PhCN	TTCorrH ₃		0.34	−1.40 ^a	−1.90	
	3-NO ₂ TTCorrH ₃			−0.84 ^a	−1.39	−1.76 ^a
	3,17-(NO ₂) ₂ TTCorrH ₃		0.94	−0.62 ^a	−1.08	−1.38

^aPeak potential at scan rate of 0.1 V/s. ^bThis redox couple can only be seen on the second positive potential sweep after scanning through the first reduction.




Figure 6. UV–visible spectral changes of 3-NO₂TTCorrH₃ in CH₂Cl₂ during (a) the controlled reduction at −0.95 V, (b) successive addition of piperidine (inset shows the Hill plot), (c) the controlled oxidation at 0.70 V, and (d) successive addition of TFA (inset shows the Hill plot).

(NO₂)₂TTCorrH₃, the effect of NO₂ on these processes is still much larger than that observed for the first oxidation. This trend in substituent effects was also observed for β-NO₂ substituted Fe(III), Cu(III), and Ge(IV) corroles having similar structures^{9,10,37} and fits with the major effect of the nitro group being to lower the energy of the lowest unoccupied molecular orbital (LUMO), vide infra.

As mentioned above, the first reduction of the investigated corroles is irreversible in pyridine, as well as in CH₂Cl₂ and PhCN (see Table 1). In all three solvents, the reduction leads to formation of the deprotonated [(Corr)H₂][−] species as the final product. An example of the thin-layer UV–visible spectral changes observed during the first one-electron reduction of 3-NO₂TTCorrH₃ in CH₂Cl₂ are shown in Figure 6a. As seen in this figure, the initial spectrum of 3-NO₂TTCorrH₃ is characterized by three separated Soret bands at 396, 438, and 464 nm and two visible Q-like bands at 589 and 667 nm. Controlled potential reduction at −0.95 V in the thin-layer cell leads to a decreased intensity of the 396 and 464 nm bands, as well as to a complete disappearance of the 438 nm band, while the two initially separated visible bands become merged into a single peak at 630–667 nm (Figure 6a). These spectral changes match perfectly spectral changes observed in the same solvent during a titration of 3-NO₂TTCorrH₃ with piperidine (Figure 6b) and gives further evidence for formation of the

deprotonated species in the electrochemical reaction, where [(Corr)H₂][−] is characterized by absorption bands at 396, 463, and 666 nm. The diagnostic log–log plot of the titration data is shown in the inset of Figure 6b and has a slope of 1.0 with a log *K* = 3.5 for dissociation of one proton under the given solution condition.

As shown by the cyclic voltammogram insert in Figure 6c, 3-NO₂TTCorrH₃ undergoes an irreversible to quasi-reversible one-electron oxidation at *E*_p = 0.50 V in CH₂Cl₂ at scan rate of 0.1 V/s. Similar irreversible oxidations have been observed for some free-base porphyrins, where the product of the first electron abstraction involves a protonation process,³⁸ although in the case of free-base corroles the oxidation mechanism seems to be more complicated. Our previous studies^{36b} suggest that the one-electron oxidation of triarylcorroles initially gives the corrole radical cation, [(Corr)H₃]⁺, which then rapidly loses a proton after reaction with another molecule of the neutral corrole in solution to generate a mixture of the free-base neutral corrole radical, (Corr)H₂, and the corrole monocation, [(Corr)H₄]⁺. In Figure 6c and d, it is possible to compare the spectral changes observed for 3-NO₂TTCorrH₃, when a controlled potential oxidation at 0.70 V was carried out (see Figure 6c), and when the same macrocycle was titrated with successive additions of TFA in CH₂Cl₂ solution. The final UV–

visible spectra under these two conditions are quite similar to each other.

As expected, the introduction of one or two nitro groups at the β -positions of the corrole macrocycle influences the corresponding acid–base properties of the molecule. Spectral changes of the type reported in Figure 6b and d for 3-NO₂TTCorrH₃ were exploited to obtain equilibrium constants for deprotonation ($\log K_a$) and protonation ($\log K_b$) processes of the three investigated free-base corroles in CH₂Cl₂, and the corresponding protonation–deprotonation constants are reported in Table 2. The equilibrium constant for proton

Table 2. Equilibrium Constants for Deprotonation ($\log K_a$) and Protonation ($\log K_b$) of (NO₂)_xTTCorrH₃ in CH₂Cl₂

cpd	$\log K_a$	$\log K_b$
	reacting with piperidine	reacting with trifluoroacetic acid
TTCorrH ₃	1.1	4.5
3-NO ₂ TTCorrH ₃	3.5	3.9
3,17-(NO ₂) ₂ TTCorrH ₃	4.9	3.3

dissociation increased upon going from TTCorrH₃ ($\log K_a = 1.1$) to 3-NO₂TTCorrH₃ ($\log K_a = 3.5$) and then to 3,17-(NO₂)₂TTCorrH₃ ($\log K_a = 4.9$), while for $\log K_b$, a reverse trend is obtained (see Table 2), indicating that the acidity of free-base corroles increases, while the basicity decreases upon increasing the number of peripheral nitro substituents. It is interesting to note that the effect of the nitro groups is higher for the increase of the acidity constants than that observed for the corresponding K_b .

Theoretical Studies. To rationalize the effects of introducing nitro groups at the periphery of TTCorrH₃, the structural, electronic, optical, and electrochemical properties of TTCorrH₃ and its nitro derivatives were theoretically investigated using DFT and TDDFT methods.

(a) **Molecular Structures.** In metal-free corroles the inner hydrogen atoms can be distributed in different ways among the four nitrogen atoms,³⁹ which results in the formation of tautomers. In this study only one tautomeric form of TTCorrH₃, 2-NO₂TTCorrH₃, 3-NO₂TTCorrH₃, and 3,17-(NO₂)₂TTCorrH₃ was examined theoretically. In the choice of the tautomers of TTCorrH₃ and its nitro derivatives, we have been guided by the X-ray structural data available for TPCorrH₃ (TPCorrH₃ = 5,10,15-triphenylcorrole)^{39c} and 2-NO₂TTCorrH₃, respectively. In both compounds the inner hydrogen atoms are located on B, C, and D pyrrole rings. The

structural analysis was performed in the gas-phase and in dichloromethane solution, using both BP86 and B3LYP functionals. The optimized structures showed very little sensitivity to the used functional and solvation effects. Table 3 collects the relevant structural parameters computed at the B3LYP level, in the gas-phase, for TTCorrH₃, 2-NO₂TTCorrH₃, 3-NO₂TTCorrH₃, and 3,17-(NO₂)₂TTCorrH₃. For 2-NO₂TTCorrH₃, comparison is made between the theoretical and experimental data.

As observed in other metal-free triarylcorroles,^{39c,40} in TTCorrH₃ the macrocycle exhibits a significant distortion from planarity to minimize the steric hindrance between the inner protons. The macrocycle adopts a puckered conformation in which the pyrrole rings turn either slightly up or down. According to the computed twist angle between the pyrrole mean planes, θ_1 , the pyrrolenine ring A and the adjacent pyrrole not directly bonded to it, B, are close to be coplanar, mostly due to the presence of an intramolecular hydrogen bond between the N₂–H group and the iminic N₁ atom (N_1 –N₂ = 2.621 Å, $\angle N_2$ –H₂...N₁ = 132.2°). The D and C rings are tipped up and down with respect to the 23 atom mean plane forming a dihedral angle (θ_3) of 22.2° with each other. The two directly bonded pyrroles, A and D, show a N–C–C–N twist angle (φ_1) of 10.3°. Comparison of the structural parameters of 2-NO₂TTCorrH₃ with those of TTCorrH₃ shows that the introduction of a nitro group at the C₂ position has very little impact on the geometry of the corrole macrocycle. As indicated by the C₁–C₂–N₅–O₁ angle (φ_2), the NO₂ group is substantially coplanar with the pyrrolenine ring A. This, together with a relatively short C₂–N₅ bond length (1.430 Å), suggests conjugation between the corrole and the nitro substituent. The geometrical data listed in Table 3 for 2-NO₂TTCorrH₃ show agreement between theory and experiment, thus confirming that the puckering of the corrole macrocycle is dictated by intrinsic electronic factors. Because of steric hindrance with the adjacent phenyl ring, the nitro group in 3-NO₂TTCorrH₃ is no longer coplanar with the pyrrolenine ring A, the C₄–C₃–N₅–O₁ angle amounting to 31.4°. This results in decreased conjugation between the nitro group and the corrole macrocycle, as confirmed by the elongation of the C–N bond. In turn, the phenyl ring on the 5 position tilts toward the bipyrrolic block by 50.2°. This is reflected in a slight increase of macrocycle puckering. In 3,17-(NO₂)₂TTCorrH₃, the puckering of the corrole macrocycle is further enhanced relative to the mononitrated corroles, due to the cooperative effects of the nitro substituents. It is worth noting, however,

Table 3. Selected Bond Lengths (Å), Bond Angles (deg), and Metrical Parameters Calculated for TTCorrH₃ and its Nitro Derivatives, 2-NO₂TTCorrH₃, 3-NO₂TTCorrH₃, and 3,17-(NO₂)₂TTCorrH₃

	TTCorrH ₃	2-NO ₂ TTCorrH ₃ ^a	3-NO ₂ TTCorrH ₃	3,17-(NO ₂) ₂ TTCorrH ₃
C–N _(nitro) ^b		1.430/1.421(3)	1.445	1.447/1.441
N ₁ –N ₂	2.621	2.639/2.623(3)	2.623	2.627
N ₂ –H ₂ ...N ₁	132.2	131.5/134(3)	130.4	129.8
θ_1 ^c	10.2	9.4/10.5(6)	12.4	13.7
θ_3 ^d	22.2	22.4/23.8(3)	23.6	27.9
φ_1 ^e	10.3	11.0/7.3(3)	12.1	14.4
φ_2 ^f		2.8/1.2(4)	31.4	32.9/20.0

^aExperimental values (this work) in italics. ^bC₂–N₅, C₃–N₅, and C₃–N₅/C₁₇–N₆ bond lengths for 2-NO₂TTCorrH₃, 3-NO₂TTCorrH₃, and 3,17-(NO₂)₂TTCorrH₃, respectively. ^cDihedral angle between the B and A pyrrole mean planes. ^dDihedral angle between the C and D pyrrole mean planes. ^eN₁–C₅–C₁₉–N₄ torsional angle. ^fC₁–C₂–N₅–O₁, C₄–C₃–N₅–O₁, and C₄–C₃–N₅–O₁/C₁₆–C₁₇–N₆–O₃ torsional angles for 2-NO₂TTCorrH₃, 3-NO₂TTCorrH₃, and 3,17-(NO₂)₂TTCorrH₃, respectively.

that while the twisting of the nitro group on the 3 position is very much the same as in 3-NO₂TTCorrH₃ (32.9° vs 31.4°), that of the nitro group on the 17 position is only 20.0°, leading to a more effective conjugation with the pertinent pyrrole.

(b) *Ground-State Electronic Structure and Optical Spectra.* To provide an interpretation of the UV/vis spectral changes accompanying the nitration of TTCorrH₃, TDDFT calculations of the electronic absorption spectra in dichloromethane solution were performed for 2-NO₂TTCorrH₃, 3-NO₂TTCorrH₃, 3,17-(NO₂)₂TTCorrH₃, and the parent TTCorrH₃. Before dealing with the excited states, we briefly discuss the ground-state electronic structure of the investigated metal-free corroles. To highlight the electronic effects of the nitro groups, the electronic structure of the unsubstituted free-base is taken as a reference. An energy level scheme of the highest occupied and lowest unoccupied Kohn–Sham orbitals of TTCorrH₃ and its nitro derivatives is shown in Figure 7.




Figure 7. Energy level scheme for TTCorrH₃ and its nitro derivatives. The Gouterman-derived MOs are indicated with red lines.

The plots of the two highest occupied and of the two lowest unoccupied MOs of these compounds are displayed in Figure 8. The HOMO–1 and HOMO of TTCorrH₃ resemble the HOMO–1(a_u) and HOMO(b_{1u}) of TPPH₂ (D_{2h} symmetry).⁴¹ Less pronounced is the similarity between the LUMO and LUMO+1 of TTCorrH₃ and the LUMO(b_{2g}) and LUMO+1(b_{3g}) of TPPH₂ (D_{2h} symmetry).⁴¹ In the nitro derivatives and in the parent TTCorrH₃, a quite large energy gap separates the HOMO–1 from the lower lying MOs, which are largely localized on the aryl groups and on the pyrrolenine ring. In the nitro derivatives, one or two MOs having some amplitude on the nitro groups interpose between the LUMO+1 and the high-

lying set of π -phenyl orbitals. According to the level scheme of Figure 7, the outstanding effect of introducing nitro groups into TTCorrH₃ is the stabilization of the LUMO and, to a lesser extent, of the HOMO–1. As a result, the LUMO/LUMO+1 splitting increases dramatically on going from TTCorrH₃ to the nitro derivatives. The large stabilization of the LUMO moving from TTCorrH₃ to the nitro derivatives is due to the C–N_(nitro) π -bonding character of this orbital well visible in the plots displayed in Figure 8. In 3-NO₂TTCorrH₃ the LUMO is slightly less stabilized than in 2-NO₂TTCorrH₃ because the nitro group is rotated by 31.4° with respect to the pertinent pyrrole plane and hence the overlap between the C3 and N5 2p_z orbitals is less effective than in 2-NO₂TTCorrH₃ where the nitro group is almost coplanar with the pertinent pyrrole. In the dinitro derivative both the LUMO and LUMO+1 experience a significant downward shift as compared to the parent TTCorrH₃. This is because both these MOs have C–N_(nitro) π -bonding character. Due to the presence of two C–N_(nitro) π -bonding interactions, the stabilization of the LUMO and LUMO+1 is quite large in 3,17-(NO₂)₂-TTCorrH₃. However, it should be noted that, for overlap reasons, the C–N_(nitro) π -bonding interaction involving the nitro group on the 3 position is somewhat less effective than that involving the nitro group on the 17 position. As pointed out earlier, the nitro groups on the 3 and 17 positions are rotated with respect to the plane of the pertinent pyrrole by 32.9° and 20.0°, respectively.

In summary, comparison between the one-electron levels of the nitro corroles and those of the parent free-base reveals that the most striking differences are the increased LUMO/LUMO+1 splitting and the diminished HOMO/LUMO energy gap, which are mainly due to the downshift of the LUMO in the nitro derivatives.

We see now how the previously discussed electronic structure changes accompanying the nitration of the corrole macrocycle reflect on the UV–visible spectroscopic properties of the investigated nitrocorroles. The excitation energies and oscillator strengths calculated for the lowest singlet excited states of TTCorrH₃ and its nitro derivatives are reported in Tables 4–7 together with the major one-electron transitions contributing to the excited-state solution vectors.

In Figure 9, the computed and experimental absorption spectra of these compounds in CH₂Cl₂ are compared. Considering first TTCorrH₃ as point of reference for the nitro derivatives, only two excited states are computed in the energy regime of the four Q bands, the 1¹A and 2¹A. As found in TPPH₂,⁴² these states originate from out-of-phase combination of the Gouterman transitions. However, due to the lifting of the LUMO/LUMO+1 degeneracy induced by the appreciable distortion of the macrocycle in TTCorrH₃, the mixing coefficients of the Gouterman transitions change significantly relative to TPPH₂. As a matter of fact, the 1¹A excited state in TTCorrH₃ is mostly described by the HOMO → LUMO transition, the transition out of the HOMO–1 into the LUMO+1 entering in this state with only a minor (15%) weight. The 2¹A is instead a nearly 50:50 mixture of the HOMO → LUMO+1 and HOMO–1 → LUMO transitions. Due to a less effective mixing of the contributing transitions and, hence, to a less effective cancellation of their large dipole moments, the oscillator strength computed for the 1¹A state is significantly larger than that of the 2¹A. By analogy with the assignment previously proposed for TPPH₂,⁴² the Q bands observed in the CH₂Cl₂ absorption spectrum of TTCorrH₃ at 652 and 574 nm are assigned to the 1¹A and 2¹A states,




Figure 8. Plots of the frontier orbitals of TTCorrH₃ and its nitro derivatives.

Table 4. Composition, Vertical Excitation Energies, E (eV/nm), and Oscillator Strengths, f , for the Lowest Optically Allowed Excited States of TTCorrH₃ in CH₂Cl₂

state	composition (%)	E	f
1 ¹ A	150a → 151a (78)	2.12/585	0.2905
	149a → 152a (15)		
2 ¹ A	150a → 152a (47)	2.30/539	0.03580
	149a → 151a (46)		
3 ¹ A	149a → 151a (48)	2.96/418	1.122
	150a → 152a (47)		
4 ¹ A	149a → 152a (80)	3.05/406	1.439
	150a → 151a (14)		

respectively, although the computed absorption wavelengths are somewhat blue-shifted with respect to the experimental values. According to the TDDFT calculations, the Q bands at 616 and 465 nm are identified as vibrational in origin. The B band profile of TTCorrH₃ is nicely accounted for by the nearly degenerate 3¹A and 4¹A excited states. These originate from in-phase combinations of the Gouterman type transitions and, hence, have large oscillator strength (see Table 4).

Coming to the nitrocorroles, in the energy regime of the Q bands, TDDFT calculations predict, just as for the parent TTCorrH₃, only two excited states, the 1¹A and 2¹A. As can be inferred from the composition of these states in Tables 4–7 for all three nitro derivatives, the 1¹A is a nearly pure (>90%) HOMO → LUMO state and the 2¹A is a mixture of the HOMO → LUMO+1 and HOMO–1 → LUMO transitions. The lowest excited state nicely accounts for the energy and intensity of the longest wavelength Q-band. The 2¹A excited state is responsible for the higher energy Q-band appearing as a

distinct peak at 588 nm in the UV–visible spectrum of 2-NO₂TTCorrH₃ and as a broad feature at around 600 nm in the spectra of the other nitro derivatives. The calculations also account for the observed red shift of this band on going from TTCorrH₃ to the nitro derivatives. Distinct from TTCorrH₃, the nitro corroles exhibit a complex B band system characterized by several overlapping features. The increased complexity of the B band system in nitrocorroles is related to the involvement of non-Gouterman transitions in the excited states accounting for this spectral region. The calculations indicate that the Gouterman-type transitions, which account for the nearly degenerate B bands of TTCorrH₃, mix to a variable extent with transitions out of the HOMO–2 (a MO largely localized on the aryl groups and on the pyrrolene ring) into the LUMO, or with transitions out of the HOMO into the LUMO+2 (a MO with sizable amplitude on the nitro groups). As for 2-NO₂TTCorrH₃, the two prominent B bands are well accounted for by the 4¹A and 5¹A excited states computed at 449 and 392 nm and with oscillator strength of 0.8068 and 0.7931, respectively. According to their composition (see Table 5), these states are largely composed of the Gouterman-type transitions with the above-mentioned non-Gouterman transitions entering with a minor, although not negligible, weight. Four intense excited states are computed in the energy regime of the three B band system of 3-NO₂TTCorrH₃. On the basis of their energy, the 3¹A and 4¹A are assigned to the lowest energy features at 465 and 438 nm, respectively, and the nearly degenerate 5¹A and 6¹A states to the most prominent feature at 397 nm.

The B band system of the dinitrocorrole is characterized by an intense, sharp feature at 483 nm followed to the blue by a broad, less intense feature. The energy and oscillator strength




Figure 9. Comparison between computed (TDDFT/B3LYP) and experimental absorption spectra of TTCorrH₃ and its nitro derivatives in CH₂Cl₂.

computed for the 3^1A excited state leaves no doubt about the assignment of the longer wavelength B band to this state. It is noteworthy that the 3^1A state consists of the same Gouterman transitions as the 2^1A , HOMO-1 \rightarrow LUMO and HOMO \rightarrow LUMO+1, with approximately reversed weights. The large transition dipoles of these transitions reinforce in the 3^1A and cancel in the 2^1A state with the result that the former is by far more intense than the latter.

Table 5. Composition, Vertical Excitation Energies, E (eV/nm), and Oscillator Strengths, f , for the Lowest Optically Allowed Excited States of 2-NO₂TTCorrH₃ in CH₂Cl₂

state	composition (%)	E	f
1^1A	161a \rightarrow 162a (94)	1.74/711	0.2655
2^1A	160a \rightarrow 162a (72)	2.21/560	0.1042
	161a \rightarrow 163a (25)		
3^1A	161a \rightarrow 164a (82)	2.61/474	0.07460
	160a \rightarrow 163a (16)		
4^1A	161a \rightarrow 163a (59)	2.76/449	0.8068
	160a \rightarrow 162a (20)		
	160a \rightarrow 164a (16)		
5^1A	160a \rightarrow 163a (49)	3.16/392	0.7931
	159a \rightarrow 162a (31)		

Table 6. Composition, Vertical Excitation Energies, E (eV/nm), and Oscillator Strengths, f , for the Lowest Optically Allowed Excited States of 3-NO₂TTCorrH₃ in CH₂Cl₂

state	composition (%)	E	f
1^1A	161a \rightarrow 162a (91)	1.86/667	0.3338
2^1A	160a \rightarrow 162a (69)	2.19/565	0.1304
	161a \rightarrow 163a (27)		
3^1A	161a \rightarrow 164a (38)	2.65/468	0.3584
	161a \rightarrow 163a (37)		
	160a \rightarrow 162a (14)		
4^1A	160a \rightarrow 163a (32)	2.73/455	0.3085
	161a \rightarrow 164a (27)		
	161a \rightarrow 163a (23)		
5^1A	159a \rightarrow 162a (54)	3.07/403	0.4981
	161a \rightarrow 164a (17)		
	160a \rightarrow 163a (14)		
6^1A	159a \rightarrow 162a (36)	3.13/396	0.5669
	160a \rightarrow 163a (33)		
	161a \rightarrow 164a (14)		

In the energy regime of the broad B feature, the calculations predict four excited states, two of which, the nearly degenerate 6^1A and 7^1A , account for most of the intensity of this band.

(c) *Reduction Potentials.* The redox behavior of metal-free corroles is rather complex owing to the proton transfer

Table 7. Composition, Vertical Excitation Energies, E (eV/nm), and Oscillator Strengths, f , for the Lowest Optically Allowed Excited States of 3,17-(NO₂)₂TTCorrH₃ in CH₂Cl₂

state	composition (%)	E	f
1^1A	172a \rightarrow 173a (96)	1.72/720	0.4537
2^1A	171a \rightarrow 173a (59)	2.04/608	0.01098
	172a \rightarrow 174a (40)		
3^1A	172a \rightarrow 174a (54)	2.46/503	0.6455
	171a \rightarrow 173a (35)		
4^1A	171a \rightarrow 174a (71)	2.69/462	0.01524
	172a \rightarrow 176a (20)		
5^1A	170a \rightarrow 173a (68)	2.81/442	0.03678
	172a \rightarrow 175a (24)		
6^1A	172a \rightarrow 176a (47)	2.90/427	0.1994
	170a \rightarrow 173a (16)		
	171a \rightarrow 175a (14)		
7^1A	172a \rightarrow 175a (37)	2.93/422	0.1740
	170a \rightarrow 173a (20)		
	171a \rightarrow 174a (16)		


Table 8. Thermodynamic Parameters (eV) for the One-Electron Reduction Reaction 1 and One-Electron Reduction Potentials (V) for TTCorrH₃, 2-NO₂TTCorrH₃, 3-NO₂TTCorrH₃, and 3,17-(NO₂)₂TTCorrH₃

	$\Delta H_{\text{ox/red}}^{\circ}$	$-T\Delta S_{\text{ox/red}}^{\circ}$	ΔE_{solv}^a	$\Delta G_{\text{ox/red}}^{\circ}$	$E_{\text{ox/red}}^{\circ}$ vs SCE	$E_{1/2}^b$ vs SCE
TTCorrH ₃	−2.93	0.02	−1.41	−2.91	−1.46	−1.40
2-(NO ₂)TTCorrH ₃	−3.49	0.03	−1.50	−3.52	−0.85	
3-(NO ₂)TTCorrH ₃	−3.44	0.03	−1.47	−3.47	−0.90	−0.84
3,17-(NO ₂) ₂ TTCorrH ₃	−3.78	0.04	−1.31	−3.82	−0.55	−0.62

^a ΔE_{solv} is the difference between the dielectric energy of the reduced species and the dielectric energy of the oxidized species. ^bExperimental values in dichloromethane solution, this work.

equilibrium reaction $\text{CorrH}_3 \rightleftharpoons [\text{CorrH}_2]^-$. As seen in the previous section, the reduction potentials of the three-protonated species, CorrH₃, positively shifts when one or two nitro groups are introduced at the β -position of the macrocycle. Also the addition of a second nitro group to the macrocycle decreases the effect of nitration on the reduction potentials under the same experimental conditions. To rationalize the observed trend, we have computed the first reduction potential of 2-NO₂TTCorrH₃, 3-NO₂TTCorrH₃, 3,17-(NO₂)₂TTCorrH₃, and the parent TTCorrH₃ in dichloromethane solution, using a DFT/COSMO model.


The standard reduction potential, $E_{\text{ox/red}}^{\circ}$, for the reaction



is related to the standard free energy change relative to the standard hydrogen electrode (SHE) through the equation

$$E_{\text{ox/red}}^{\circ} = -\frac{\Delta G_{\text{ox/red}}^{\circ} - \Delta G_{\text{SHE}}^{\circ}}{nF} \quad (2)$$

In eq 2, n is the number of electrons consumed in the reduction reaction, 1 in the present case, and F is a constant. If the free energies are expressed in molar units, F is the Faraday constant (the negative of the charge on one mole of electrons), but if energies are expressed in electron volts per molecule, as in the present paper, then F is equal to the unit charge e^- . $\Delta G_{\text{ox/red}}^{\circ}$ is the free energy change associated with reaction 1, whereas $\Delta G_{\text{SHE}}^{\circ}$ is the free energy change of the following reaction



$\Delta G_{\text{SHE}}^{\circ}$ has been established in the literature to be -4.28 eV .⁴³

As the experimental half-wave potentials of the investigated metal-free corroles were measured using SCE as the reference electrode, the theoretical reduction potentials obtained from eq 2 were converted to SCE referenced potentials by subtracting 0.09 V, which was obtained by taking the literature value of the reduction potential of aqueous SCE relative to SHE at 298 K (0.24 V)⁴⁴ and correcting it to its corresponding value in dichloromethane solution using the literature value of the liquid junction potential (l_{jp}) for dichloromethane–water (0.15 V).⁴⁵ The corrected one-electron reduction potentials of TTCorrH₃, 2-NO₂TTCorrH₃, 3-NO₂TTCorrH₃, and 3,17-(NO₂)₂TTCorrH₃ are compared with the experimental half-wave potentials in Table 8 where the thermodynamic parameters computed for the one-electron reduction process in eq 1 are also reported. The data in Table 8 indicate that there is an excellent agreement between the theoretically predicted one-electron reduction potentials and the available experimental values. The thermodynamic data reveal that the one-electron reduction process is energetically controlled by enthalpic factors, the entropic factors playing a negligible role. The

ΔE_{solv} values indicate that the energetic contribution of solvation to the enthalpic term, though relevant, is substantially constant along the series. Thus, the $\Delta H_{\text{ox/red}}^{\circ}$ trend is primarily determined by the electronic enthalpy that favors energetically the reduced form over the oxidized one in the order $\text{TTCorrH}_3 \ll 2\text{-NO}_2\text{TTCorrH}_3 \cong 3\text{-NO}_2\text{TTCorrH}_3 < 3,17\text{-(NO}_2)_2\text{TTCorrH}_3$. In nitrocorroles, the one-electron reduced form is much more stable than the oxidized form as compared to TTCorrH₃, and this is consistent with the added electron entering a MO with C–N_(nitro) π -bonding character in the former, as witnessed by the significant shortening of the C–N_(nitro) bond. In 2-NO₂TTCorrH₃ and 3-NO₂TTCorrH₃, the C–N_(nitro) bond shortens by 0.027 and 0.033 Å, respectively. In the dinitro derivative, the two C–N_(nitro) bonds shorten, on the average, by 0.025 Å.

CONCLUSIONS

We have reported the first example of efficient insertion of nitro groups into the β -pyrrole positions of corrole free bases, effectively overcoming the formation of isocorrole by controlling the amount and the nature of the oxidant used for the reaction. The regioselectivity usually observed in the functionalization of corrole ring is also demonstrated for the nitration reaction, producing the 3-nitro- and the 3,17-dinitro-corroles as the main reaction products of mono- and disubstitution; we have also been able to characterize some other minor regioisomers, such as the 2-nitro, 2,3-, and the 3,12-dinitro derivatives. It is interesting to note that this last regioisomer is the first example of antipodal functionalization of corrole ring.

The introduction of nitro substituents at the β -pyrrole positions of the corrole ring strongly influence the chemical and spectroscopic behavior of the macrocycle. The strong electron-withdrawing character of the nitro group leads to a positive shift of the $E_{1/2}$ of the redox processes of corrole and to an increase of the macrocycle acidity. It is noteworthy that the introduction of the first nitro group is more effective than the second substituent in changing the properties of the functionalized macrocycle. Optical absorption spectra of β -nitrocorroles are strongly influenced by the peripheral nitro groups, which increase the number of bands, characterized also by a significant red shift.

The theoretical results on these β -nitrocorrole derivatives also afforded significant information, closely matching the experimental observations. It was found that the β -NO₂ substituents conjugate with the π -aromatic system of the macrocycle, which initiates significant changes in both the spectroscopic and redox properties of the so functionalized corroles. This effect is more pronounced when the nitro group is introduced at the 2-position, because in this case the conjugation is, for steric reasons, more efficient than in the 3-nitro isomer.

In closing, these nitro-functionalized corroles are also useful starting materials for further modifications of the corrole ring and their availability can open new opportunities for the preparation of functionalized corroles; these studies are now in progress in our laboratories.

■ ASSOCIATED CONTENT

● Supporting Information

CIF files of 2-NO₂TTCorrH₃, 2,3-(NO₂)₂TTCorrCoPPh₃, and 3,12-(NO₂)₂TTCorrCoPPh₃. This material is available free of charge via the Internet at <http://pubs.acs.org>

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: roberto.paolesse@uniroma2.it (R.P.); kkadish@uh.edu (K.M.K.); kmsmith@lsu.edu (K.M.S.); angela.rosa@unibas.it (A.R.).

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We gratefully acknowledge the support of MIUR, Italy (PRIN project 2009Z9ASCA), the US National Institutes of Health (K.M.S., grant CA132861), and the Robert A. Welch Foundation (K.M.K., grant E-680).

■ REFERENCES

- (1) Nardis, S.; Monti, D.; Paolesse, R. *Mini-Rev. Org. Chem.* **2005**, *2*, 355.
- (2) Paolesse, R. *Synlett* **2008**, 2215.
- (3) Paolesse, R. *The Porphyrin Handbook*; Kadish, K. M., Smith, K. M., Guillard, R., Eds.; Academic Press: New York, 2000; Vol. 2, pp 201–232.
- (4) Gryko, D. T. *J. Porphyrins Phthalocyanines* **2008**, *12*, 906.
- (5) Gross, Z.; Aviv-Harel, I. *Coord. Chem. Rev.* **2011**, *255*, 717–736.
- (6) (a) Gross, Z.; Aviv-Harel, I. *Chem.—Eur. J.* **2009**, *15*, 8382–8394. (b) Kupersmidt, L.; Okun, Z.; Amit, T.; Mandel, S.; Saltsman, I.; Mohammed, A.; Bar-Am, O.; Gross, Z.; Youdim, M. B. H. *J. Neurochem.* **2010**, *113*, 363–373. (c) Kanamori, A.; Catrinescu, M. M.; Mohammed, A.; Gross, Z.; Levin, L. A. *J. Neurochem.* **2010**, *114*, 488–498.
- (7) Lemon, C. M.; Brothers, P. J. *J. Porphyrins Phthalocyanines* **2011**, *15*, 809–834.
- (8) Walker, F. A.; Licoccia, S.; Paolesse, R. *J. Inorg. Biochem.* **2006**, *100*, 810 and references therein.
- (9) Stefanelli, M.; Mandoj, F.; Mastroianni, M.; Nardis, S.; Mohite, P.; Fronczek, F. R.; Smith, K. M.; Kadish, K. M.; Xiao, X.; Ou, Z.; Chen, P.; Paolesse, R. *Inorg. Chem.* **2011**, *50*, 8281–8292.
- (10) Mastroianni, M.; Zhu, W.; Stefanelli, M.; Nardis, S.; Fronczek, F. R.; Smith, K. M.; Ou, Z.; Kadish, K. M.; Paolesse, R. *Inorg. Chem.* **2008**, *47*, 11680–11687.
- (11) Saltsman, I.; Mohammed, A.; Goldberg, I.; Tkachenko, E.; Botoshansky, M.; Gross, Z. *J. Am. Chem. Soc.* **2002**, *124*, 7411–7420.
- (12) Stefanelli, M.; Mastroianni, M.; Nardis, S.; Licoccia, S.; Fronczek, F. R.; Smith, K. M.; Zhu, W.; Ou, Z.; Kadish, K. M.; Paolesse, R. *Inorg. Chem.* **2007**, *46*, 10791–10799.
- (13) Pomarico, G.; Fronczek, F. R.; Nardis, S.; Smith, K. M.; Paolesse, R. *J. Porphyrins Phthalocyanines* **2011**, *15*, 1086–1092.
- (14) (a) Stefanelli, M.; Nardis, S.; Tortora, L.; Fronczek, F. R.; Smith, K. M.; Licoccia, S.; Paolesse, R. *Chem. Commun.* **2011**, *47*, 4255–4257. (15) Stefanelli, M.; Shen, J.; Zhu, W.; Mastroianni, M.; Mandoj, F.; Nardis, S.; Ou, Z.; Kadish, K. M.; Fronczek, F. R.; Smith, K. M.; Paolesse, R. *Inorg. Chem.* **2009**, *48*, 6879–6887.
- (16) (a) Mandoj, F.; Nardis, S.; Pomarico, G.; Paolesse, R. *J. Porphyrins Phthalocyanines* **2008**, *12*, 19–26. (b) Ngo, T. H.; Van Rossom, W.; Dehaen, W.; Maes, W. *Org. Biomol. Chem.* **2009**, *7*, 439–443. (c) Capar, C.; Thomas, K. E.; Ghosh, A. *J. Porphyrins Phthalocyanines* **2008**, *12*, 964–967.
- (17) Sheldrick, G. M. *Acta Crystallogr., Sect. A* **2008**, *64*, 112.
- (18) APEX2, SAINT and SADABS; Bruker AXS Inc. Madison, Wisconsin, USA, 2007.
- (19) Flack, H. D. *Acta Crystallogr.* **1983**, *A39*, 876–881.
- (20) Spek, A. L. *Acta Crystallogr. Sect. D-Biol. Crystallogr.* **2009**, *65*, 148.
- (21) Lin, X. Q.; Kadish, K. M. *Anal. Chem.* **1985**, *57*, 1498–1501.
- (22) (a) TURBOMOLE V6.3 2011, a development of University of Karlsruhe and Forschungszentrum Karlsruhe GmbH, 1989–2007, TURBOMOLE GmbH; available from <http://www.turbomole.com>. (b) Ahlrichs, R.; Bär, M.; Häser, M.; Horn, H.; Kölmel, C. *Chem. Phys. Lett.* **1989**, *162*, 165.
- (23) (a) Becke, A. *Phys. Rev. A* **1988**, *38*, 3098. (b) Perdew, J. P. *Phys. Rev. B* **1986**, *33*, 8822.
- (24) (a) Becke, A. D. *J. Chem. Phys.* **1993**, *98*, 5648. (b) Lee, C.; Yang, W.; Parr, R. G. *Phys. Rev. B* **1988**, *37*, 785.
- (25) Weigend, F.; Ahlrichs, R. *Phys. Chem. Chem. Phys.* **2005**, *7*, 3297.
- (26) Eichkorn, K.; Treutler, C. *Phys. Chem. Lett.* **1995**, *240*, 283.
- (27) Weigend, F. *Phys. Chem. Chem. Phys.* **2006**, *8*, 1057–1065.
- (28) (a) Klamt, A.; Schürmann, G. *J. Chem. Soc., Perkin Trans.* **1993**, *2*, 799. (b) Klamt, A.; Jonas, V. *J. Chem. Phys.* **1996**, *105*, 9972.
- (29) (a) Gross, E. U. K.; Dobson, J. F.; Petersilka, M. In *Density Functional Theory; Topics in Current Chemistry*; Nalewajski, R. F., Ed.; Springer: Heidelberg, 1996. (b) Casida, M. E. In *Recent Advances in Density Functional Methods*; Chong, D. P., Ed.; World Scientific: Singapore, 1995; Vol. 1, p 155. (c) Parr, R. G.; Yang, W. *Density Functional Theory of Atoms and Molecules*; Oxford University Press: New York, 1989. (d) Drew, A.; Head-Gordon, M. *Chem. Rev.* **2005**, *105*, 4009.
- (30) (a) Klamt, A. *J. Phys. Chem.* **1996**, *100*, 3349–3353. (b) Scalmani, G.; Frisch, M. J.; Mennucci, B.; Tomasi, J.; Cammi, R.; Barone, V. *J. Chem. Phys.* **2006**, *124*, 094107–094121.
- (31) Tortora, L.; Nardis, S.; Fronczek, F. R.; Smith, K. M.; Paolesse, R. *Chem. Commun.* **2011**, *47*, 4243–4245.
- (32) Nardis, S.; Pomarico, G.; Fronczek, F. R.; Vicente, M. G. H.; Paolesse, R. *Tetrahedron Lett.* **2007**, *48*, 8643–8646.
- (33) Nardis, S.; Pomarico, G.; Mandoj, F.; Fronczek, F. R.; Smith, K. M.; Paolesse, R. *J. Porphyrins Phthalocyanines* **2010**, *14*, 753–757.
- (34) Setsune, J.; Tsukajima, A.; Watanabe, J. *Tetrahedron Lett.* **2006**, *47*, 1817–1820.
- (35) Flint, D. L.; Fowler, R. L.; LeSaulnier, T. D.; Long, A. C.; O'Brien, A. Y.; Geier, G. R., III. *J. Org. Chem.* **2010**, *75*, 553–563.
- (36) (a) Ou, Z.; Shen, J.; Shao, J.; E, W.; Galewski, M.; Gryko, D. T.; Kadish, K. M. *Inorg. Chem.* **2007**, *46*, 2775–2786. (b) Shen, J.; Shao, J.; Ou, Z.; E, W.; Koszarna, B.; Gryko, D. T.; Kadish, K. M. *Inorg. Chem.* **2006**, *45*, 2251–2265.
- (37) Nardis, S.; Stefanelli, M.; Mohite, P.; Pomarico, G.; Tortora, L.; Manowong, M.; Chen, P.; Kadish, K. M.; Fronczek, F. R.; McCandless, G. T.; Smith, K. M.; Paolesse, R. *Inorg. Chem.* **2012**, *51*, 3910–3920.
- (38) (a) Kadish, K. M.; Chen, P.; Enakieva, Y. Y.; Nefedov, S. E.; Gorbunova, Y. G.; Tsivadze, A. Y.; Lemeune, A.; Stern, C.; Guillard, R. *J. Electroanal. Chem.* **2011**, *656*, 61–71. (b) Inisan, C.; Saillard, J.-Y.; Guillard, R.; Tabard, A.; Mest, Y. L. *New J. Chem.* **1998**, 823–830.
- (39) (a) *Porphyrins: excited states and dynamics*; Gouterman, M., Rentzepis, P. M., Straub, K. D., Eds.; American Chemical Society: New York, 1987; Vol. 321. (b) Storm, C. B.; Teklu, Y. *J. Am. Chem. Soc.* **1972**, *94*, 1745–1747. (c) Ding, T.; Harvey, J. D.; Ziegler, C. J. *J. Porphyrins Phthalocyanines* **2005**, *9*, 22–27.
- (40) (a) Gross, Z.; Galili, N.; Simkhovich, L.; Saltsman, I.; Botoshansky, M.; Bläser, D.; Boese, R.; Goldberg, I. *Org. Lett.* **1999**, *1*, 599. (b) Simkhovich, L.; Goldberg, I.; Gross, Z. *J. Inorg. Biochem.* **2000**, *80*, 235–238. (c) Paolesse, R.; Nardis, S.; Venanzi, M.; Mastroianni, M.; Russo, M.; Fronczek, F. R.; Vicente, M. G. H. *Chem.—Eur. J.* **2003**, *9*, 1192–1197.
- (41) Palummo, M.; Hogan, C.; Sottile, F.; Bagala, P.; Rubio, A. *J. Chem. Phys.* **2009**, *131*, 084102–084107.

- (42) De Luca, G.; Romeo, A.; Monsù Scolaro, L.; Ricciardi, G.; Rosa, A. *Inorg. Chem.* **2009**, *48*, 8493.
- (43) Kelly, C. P.; Cramer, C. J.; Truhlar, D. G. *J. Phys. Chem. B* **2006**, *110*, 16066–16081.
- (44) Bard, A. J.; Faulkner, L. R. *Electrochemical Methods: Fundamentals and Applications*; Wiley: New York, 2001.
- (45) (a) Krishtalik, L. I.; Alpatova, N. M.; Ovsyannikova, E. V. *J. Electroanal. Chem.* **1992**, *329*, 1–8. (b) Tsierkezos, N. *J. Sol. Chem.* **2008**, *37*, 1437–1448.