Organic & Biomolecular Chemistry

Cite this: Org. Biomol. Chem., 2011, 9, 4425

www.rsc.org/obc

Asymmetric alkynylation of aldehydes with propiolates without high reagent loading and any additives[†]

Naoto Kojima,* Shogo Nishijima, Kaoru Tsuge and Tetsuaki Tanaka*

Received 29th March 2011, Accepted 20th April 2011 DOI: 10.1039/c1ob05489a

The asymmetric alkynylation of aliphatic and aromatic aldehydes with propiolates was mediated by dialkylzinc and a novel prolinol catalyst without high reagent loading and any additives, such as Ti(O*i*-Pr)₄, to give the corresponding γ -hydroxy- α , β -acetylenic esters with high enantiomeric excess of up to 95%.

Chiral propargyl alcohols are versatile building blocks for fine chemicals, pharmaceuticals, and natural products.¹ A typical example is chiral γ -hydroxy- α , β -acetylenic esters, considered attractive molecules because they contain three functionalities: a hydroxyl group, an alkyne, and an ester, in a small structure. Many efficient syntheses that employed them as useful and versatile intermediates have been reported.^{2,3} The two most common methods to prepare optically active γ -hydroxy- α , β -acetylenic esters are i) the asymmetric reduction of γ -oxo- α , β -acetylenic esters with a chiral reagent, and ii) the asymmetric alkynylation of aldehydes with propiolates. The latter has attracted widespread interest in recent years because it is able to achieve both new C-C bond formation and construction of a stereogenic centre. However, previous studies of the asymmetric alkynylation of aldehydes have concentrated on the addition of simple terminal alkynes, for example, phenylacetylene.⁴ Difficulty arises in the asymmetric alkynylation using propiolate as the nucleophile because the reactivity of propiolate is greatly different from that of simple terminal alkynes.5

Recently, Pu and co-workers reported that BINOL and H₈-BINOL derivatives catalysed the Et₂Zn-mediated asymmetric alkynylation of aldehydes with methyl propiolate.⁶ Wang's group developed the enantioselective addition of methyl propiolate to aromatic aldehydes with β -sulfonamide alcohol as the ligand.⁷ Although previous studies, including those methods, gave products in good yields with high enantioselectivities, most of them required excess amounts of reagents and additives, such as Ti(O*i*-Pr)₄. Trost *et al.*⁸ and Wang and co-workers⁹ independently demonstrated the Me₂Zn-mediated enantioselective addition of methyl propiolate to aldehydes without any additives. However, their reactions still required excess amounts of Me_2Zn and propiolate. Herein, we disclose a dialkylzinc-mediated asymmetric alkynylation of aliphatic and aromatic aldehydes using propiolates, which is catalysed by a novel prolinol system and does not require any additives and high reagent loading.

We focused on a prolinol derivative for its potential use as a chiral ligand in the asymmetric alkynylation with propiolates, because there are many reports of the asymmetric alkylation and alkenylation of aldehydes with prolinol as the catalyst.¹⁰ However, to our knowledge, there are few studies of the asymmetric alkynylation of aldehydes using prolinol as the catalyst.⁸

First, we attempted to perform the enantioselective addition of ethyl propiolate to cyclohexanecarbaldehyde using (S)-(+)diphenyl(1-methylpyrrolidin-2-yl)methanol (DPMPM, **2c**) as the ligand, and obtained chiral γ -hydroxy- α , β -acetylenic ester **1a** in good yield with moderate enantioselectivity (88%, 69% ee) (Table 1, Entry 3).¹¹ At the same time, Zhang and co-workers reported the enantioselective addition of alkynes to aldehydes using *N*-benzyl prolinol as the catalyst.¹² As the reaction using propiolates was not

 Table 1
 Ligand screening in the model reaction



" Determined by 1H NMR. Ee was determined by HPLC.

Graduate School of Pharmaceutical Sciences, Osaka University, 1-6 Yamadaoka, Suita, Osaka 565-0879, Japan. E-mail: kojima@phs.osakau.ac.jp, t-tanaka@phs.osaka-u.ac.jp; Fax: +81-6-6879-8214; Tel: +81-6-6879-8210

[†] Electronic supplementary information (ESI) available: Synthesis of ligand **2d**, experimental procedures of asymmetric alkynylation, determination of stereochemistry of products, characterization data, and ¹H/¹³C NMR spectra of products. See DOI: 10.1039/c1ob05489a



reported in their letter, we examined the asymmetric alkynylation with their chiral ligands (**2a** and **b**).

The reactions catalysed by their chiral ligands (**2a** and **b**) gave γ -hydroxy- α , β -acetylenic esters in good yields, but the enantioselectivities were low (~50% ee) (Entries 1 and 2). As a result of several investigations, we found that a novel *N*-methyl prolinol derivative with two 3,5-dimethylphenyl moieties (**2d**)¹³ catalysed the asymmetric alkynylation in high yield with good enantioselectivity (Entry 4).

Table 2 shows the optimization of the reaction conditions. To a solution of 30 mol% of ligand 2d in toluene, Et₂Zn (2.0 equiv.) and ethyl propiolate (2.0 equiv.) were added at room temperature. After 12 h, cyclohexanecarbaldehyde (1.0 equiv.) was added and the reaction mixture was stirred for 3 h, giving chiral propargyl alcohol 1a in good yield with high enantioselectivity (Entry 1). The previously reported dialkylzinc-mediated alkynylation of aldehydes required a mixing step prior to the addition of aldehydes to prepare an alkynylzinc species. Without this step, the alkyl adduct was obtained as a by-product¹⁴ because chiral ligands, such as BINOL¹⁵ and β -sulfonamide alcohol,¹⁶ would catalyse the addition of dialkylzinc to aldehydes. Surprisingly, the reaction that did not include a mixing step prior to the addition of aldehyde in the presence of 2d gave propargyl alcohol 1a in high yield, and no ethyl adduct was detected (Entry 2). Furthermore, this improvement not only simplified the reaction but also increased the enantioselectivity. We surmise that ligand 2d can catalyse the formation of an alkynylzinc species more efficiently than previous ligands and therefore, the long mixing time afforded dialkynylzinc species, thereby decreasing the enantioselectivity.¹⁴ Fortunately, the efficiency of our catalyst 2d enabled the reduction of the amounts of Et₂Zn and propiolate (Entries 2-5). The reaction

using 1.0 equiv. of propiolate and 1.2 equiv. of Et_2Zn gave almost the same result as that of Entry 2 (Entry 4). When the reaction temperature was lowered to 0 °C, the yield was decreased (Entry 6). On the other hand, there was no significant change in both yield and selectivity when the reaction was conducted at 40 °C compared with the case at room temperature (Entry 7). Increasing the amount of **2d** did not improve yield or selectivity (Entry 8), and the yield in the reaction using 0.1 equiv. of **2d** was reduced (Entry 9). The use of CH₂Cl₂, *n*-hexane, and Et₂O instead of toluene decreased both yield and enantioselectivity, whereas the reaction in THF hardly occurred (Entries 10–13).

Under the above optimal conditions (Table 2, Entry 4), we examined the alkynylation of aliphatic aldehydes with ethyl propiolate (Table 3). The reaction with propionaldehyde gave alcohol **1b** in moderate yield but with high enantioselectivity (Entry 1). Isobutylaldehyde gave the corresponding alcohol **1c** in good yield with high enantioselectivity (Entry 3). The use of a bulky aldehyde, pivalaldehyde, resulted in only a small decrease in enantioselectivity (Entry 4).

Next, we examined the alkynylation of aromatic aldehydes (Table 4). The asymmetric alkynylation of benzaldehyde with ethyl propiolate under the optimal conditions afforded the desired alkynylated compound **1e** with high enantioselectivity, but the yield was low (32%) because the conditions gave an ethylation product (44%) as the main product (Entry 1). The use of Me₂Zn in place of Et₂Zn dramatically decreased the alkyl adduct (3%) and increased the yield of **1e** to 52% (Entry 2).¹⁷ By increasing the reagent loading (Me₂Zn and ethyl propiolate) to 2.0 equiv., the yield of **1e** was improved to 78% with high enantioselectivity (95% ee) (Entry 3). Having established the optimal reaction conditions for aromatic aldehydes, we investigated the alkynylation of various

2

3e

4





^a Aldehyde (1.0 mmol), ethyl propiolate (1.0 mmol), Et₂Zn (1 M solution in n-hexane, 1.2 mmol), and 2d (0.3 mmol) in toluene (6 mL) were stirred at rt under Ar. ^b Isolated yield. ^c Ee was determined by HPLC. The stereochemistry of the products was confirmed by the modified Mosher method. d Determined by 1H NMR. The reaction was carried out with 3.0 mmol of aldehyde.

 Table 4
 Asymmetric alkynylation of aromatic aldehydes^a



^a Aldehyde (1.0 mmol), ethyl propiolate (2.0 mmol), Me₂Zn (1 M solution in n-hexane, 2.0 mmol), and 2d (0.3 mmol) in toluene (6 mL) were stirred at rt under Ar. ^b Isolated yield. ^c Ee was determined by HPLC. The stereochemistry of the products was confirmed by the modified Mosher method after transformation to the corresponding allyl alcohols. For details, see Electronic Supplementary Information.^d Ethyl propiolate (1.0 eq.) and Et_2Zn (1.2 eq.) were used. ^e Ethyl propiolate (1.0 eq.) and Me_2Zn (1.2 eq.) were used. ^{*f*} Absolute configuration was determined by comparing the optical rotation with the literature value after conversion into the known methyl ester. For details, see Electronic Supplementary Information

aromatic aldehydes. The reactions of benzaldehydes containing a bromine atom at the ortho-, meta-, and para-positions gave good yields and high enantioselectivities, respectively (Entries 46). Both *p*-anisaldehyde and *p*-formylphenyl acetate, respectively, were converted into the corresponding alcohols 1i and 1j with high enantioselectivities, but the yield of the former was moderate (54%) because of less reactivity due to the electron-donating substituent (Entries 7 and 8). The desired product was also obtained when heterocyclic aldehyde was employed as the substrate, although the yield and the selectivity were decreased (Entry 9). The results showed that this catalytic system has a broad generality for aliphatic and aromatic aldehvdes.

Asymmetric alkynylation in the presence of chiral prolinol 2d afforded (R)-propargyl alcohol. The absolute configuration of the newly created chiral centre is predictable based on the model (Fig. 1). The model is analogous to that proposed by Novori and co-workers for the Me₂Zn addition to aldehydes.¹⁸ The carbonyl oxygen atom of the aldehyde coordinates to the zinc of the bimetallic complex generated from the chiral ligand, zinc acetylide, and Et₂Zn, and then an alkynyl group attacks the carbonyl carbon of the aldehyde as shown, giving an (R)-isomer.



Fig. 1 Model predicting the stereochemistry.

In conclusion, we have demonstrated the enantioselective addition of propiolates to aliphatic and aromatic aldehydes in the presence of a novel prolinol catalyst without high reagent loading and any additives. Our chiral ligand systems make it possible to prepare easily chiral γ -hydroxy- α , β -acetylenic esters. It is interesting to catalyse the asymmetric addition of propiolates to aldehydes by structurally simple ligand 2d with good enantioselectivity. The scope, mechanism, and synthetic application are under investigation.

Acknowledgements

This work was partially supported by a Grant-in-Aid for Young Scientists (B) [No. 21790113] from The Ministry of Education, Culture, Sports, Science and Technology, Japan (MEXT) and a Grant-in-Aid for Scientific Research on Innovative Areas [No. 22136006] from Japan Society for the Promotion of Science (JSPS).

Notes and references

- 1 Acetylene Chemistry, ed. F. Diederich, P. J. Stang and R. R. Tykwinski, WILEY-VCH, Weinheim, Germany, 2005.
- 2 Review of synthesis and applications of chiral γ -hydroxy- α , β -acetylenic esters: G. Gao and L. Pu, Sci. China Chem., 2010, 53, 21.
- 3 Recent reports of applications of γ -hydroxy- α , β -acetylenic esters in organic synthesis: (a) R. Csuk, A. Barthel, S. Schwarz, H. Kommera and R. Paschke, Bioorg. Med. Chem., 2010, 18, 2549; (b) V. Shekhar, D. K. Reddy, V. Suresh, D. C. Babu and Y. Venkateswarlu, Tetrahedron Lett., 2010, 51, 946; (c) Y. Zhang, H.-D. Hao and Y. Wu, Synlett, 2010, 905; (d) B. M. Trost, B. M. O'Boyle and D. Hund, J. Am. Chem. Soc., 2009, 128, 15061; (e) N. Kojima, S. Nishijima and T. Tanaka, Synlett, 2009, 3171; (f) Y. Yue, M. Turlington, X.-Q. Yu and L. Pu, J. Org. Chem., 2009, 74, 8681; (g) L. Lin, Q. Zhao, A.-N. Li, F. Ren, F. Yang and R. Wang, Org. Biomol. Chem., 2009, 7, 3663; (h) L. Liu,

- 4 Recent review of enantioselective addition of alkynes to carbonyl groups: B. M. Trost and A. H. Weiss, *Adv. Synth. Catal.*, 2009, **351**, 963.
- 5 For example: S. P. Shahi and K. Koide, *Angew. Chem., Int. Ed.*, 2004, **43**, 2525.
- 6 (a) G. Gao, Q. Wang, X.-Q. Yu, R.-G. Xie and L. Pu, Angew. Chem., Int. Ed., 2006, 45, 122; (b) M. Turlington, A. M. DeBerardinis and L. Pu, Org. Lett., 2009, 11, 2441; (c) X. Chen, W. Chen, L. Wang, X.-Q. Yu, D.-S. Huang and L. Pu, Tetrahedron, 2010, 66, 1990.
- 7 L. Lin, X. Jiang, W. Liu, L. Qiu, Z. Xu, J. Xu, A. S. C. Chan and R. Wang, *Org. Lett.*, 2007, **9**, 2329.
- 8 B. M. Trost, A. H. Weiss and A. J. von Wangelin, J. Am. Chem. Soc., 2006, **128**, 8.

- 9 J.-C. Zhong, S.-C. Hou, Q.-H. Bian, M.-M. Yin, R.-S. Na, B. Zheng, Z.-Y. Li, S.-Z. Liu and M. Wang, *Chem.-Eur. J.*, 2009, **15**, 3069.
- 10 L. Pu and H.-B. Yu, Chem. Rev., 2001, 101, 757.
- 11 Niwa and Soai reported the enantioselective addition of phenylacetylene to benzaldehyde in the presence of DPMPM (2c). However, there was only one example, and the enantioselectivity was only 4% ee. S. Niwa and K. Soai, J. Chem. Soc., Perkin Trans. 1, 1990, 937.
- 12 Z. Xu, N. Wu, Z. Ding, T. Wang, J. Mao and Y. Zhang, *Tetrahedron Lett.*, 2009, **50**, 926.
- 13 Ligand **2d** was easily prepared from L-proline in 4 steps. For details, see Electronic Supplementary Information.
- 14 S. Dahmen, Org. Lett., 2004, 6, 2113.
- 15 X.-W. Yang, J.-H. Sheng, C.-S. Da, H.-S. Wang, W. Su, R. Wang and A. S. C. Chan, J. Org. Chem., 2000, 65, 295.
- 16 (a) M. Yus, D. J. Ramón and O. Preito, *Tetrahedron: Asymmetry*, 2002, 13, 1573; (b) T. Bauer, J. Tarasiuk and K. Pasniczek, *Tetrahedron: Asymmetry*, 2002, 13, 77.
- 17 Noyori and co-workers reported that Me₂Zn was less reactive than Et₂Zn in the asymmetric alkylation of aldehydes in the presence of a chiral amino alcohol: M. Kitamura, S. Okada, S. Suga and R. Noyori, J. Am. Chem. Soc., 1989, 111, 4028.
- 18 (a) T. Rasmussen and P.-O. Norrby, J. Am. Chem. Soc., 2003, **125**, 5130; (b) M. Yamakawa and R. Noyori, Organometallics, 1999, **18**, 128; (c) M. Yamakawa and R. Noyori, J. Am. Chem. Soc., 1995, **117**, 6327.