Cycloisomerization versus Hydration Reactions in Aqueous Media: A Au(III)-NHC Catalyst That Makes the Difference

2012 Vol. 14, No. 10 2520–2523

ORGANIC LETTERS

Eder Tomás-Mendivil,[†] Patrick Y. Toullec,[‡] Josefina Díez,[†] Salvador Conejero,^{*,§} Véronique Michelet,^{*,‡} and Victorio Cadierno^{*,†}

Laboratorio de Compuestos Organometálicos y Catálisis (Unidad Asociada al CSIC), Departamento de Química Orgánica e Inorgánica, Universidad de Oviedo, Julián Clavería 8, E-33006 Oviedo, Spain, ChimieParisTech, Laboratoire Charles Friedel, ENSCP, UMR 7223, 11 rue P. et M. Curie, F-75231, Paris Cedex 05, France, and Departamento de Química Inorgánica, Instituto de Investigaciones Químicas (IIQ), CSIC/Universidad de Sevilla, Avda. Américo Vespucio 49, E-41092, Sevilla, Spain

sconejero@iiq.csic.es; veronique-michelet@chimie-paristech.fr; vcm@uniovi.es

Received March 30, 2012



A novel water-soluble Au(III)-NHC complex has been synthesized and successfully applied in the intramolecular cyclization of γ -alkynoic acids into enol-lactones under biphasic toluene/water conditions, thus representing a rare example of an active and selective catalyst for this transformation in aqueous media. Remarkably, competing alkyne hydration processes were not observed, even during the desymmetrization reaction of challenging 1,6-diyne substrates. In addition, after phase separation, the water-soluble Au(III) catalyst could be recycled 10 times without loss of activity or selectivity.

After a long induction period, gold complexes have emerged in recent years as versatile and powerful catalysts

10.1021/ol300811e © 2012 American Chemical Society **Published on Web 04/30/2012** for synthetic organic chemistry.¹ The catalytic hydration of alkynes by Na[AuCl₄] reported by Fukuda and Utimoto in 1991 initiated rapid development of the field.² Since then, gold complexes have been widely employed in many organic transformations, with special focus on the electrophilic π -activation of unsaturated carbon–carbon bonds (alkynes, enynes, alkenes, allenes, etc.) toward nucleophiles.^{1,3} Relativistic effects have been evoked to explain the superior effectiveness of gold as a π -activator compared to other transition metals.⁴ In addition, it has also been largely demonstrated that the stability and Lewis acidity of Au(I) and Au(III) species can be easily modulated by the surrounding ligands, thus enabling the fine tuning of the activity and the chemo-, regio-, and/or stereoselectivity of

[†]Universidad de Oviedo.

[‡]ChimieParisTech.

[§] CSIC/Universidad de Sevilla.

Hashmi, A. S. K.; Hutchings, G. J. Angew. Chem., Int. Ed. 2006, 45, 7896–7936. (b) Hashmi, A. S. K. Chem. Rev. 2007, 107, 3180–3211.
 (c) Li, Z.; Brouwer, C.; He, C. Chem. Rev. 2008, 108, 3239–3265. (d) Arcadi, A. Chem. Rev. 2008, 108, 3266–3325. (e) Fürstner, A. Chem. Soc. Rev. 2009, 38, 3208–3221. (f) Hashmi, A. S. K. Angew. Chem., Int. Ed. 2010, 49, 5232–5241. (g) Corma, A.; Leyva-Pérez, A.; Sabater, M. J. Chem. Rev. 2011, 111, 1657–1712.

⁽²⁾ Fukuda, Y.; Utimoto, K. J. Org. Chem. 1991, 56, 3729-3731.

⁽³⁾ See, for example: (a) Fürstner, A.; Davies, P. W. Angew. Chem., Int. Ed. 2007, 46, 3410–3449. (b) Shen, H. C. Tetrahedron 2008, 64, 3885– 3903. (c) Muzart, J. Tetrahedron 2008, 64, 5815–5849. (d) Shen, H. C. Tetrahedron 2008, 64, 7847–7870. (e) Jiménez-Núñez, E.; Echavarren, A. M. Chem. Rev. 2008, 108, 3326–3350. (f) Patil, N.-T.; Yamamoto, Y. Chem. Rev. 2008, 108, 3395–3442. (g) Michelet, V.; Toullec, P. Y.; Genêt, J.-P. Angew. Chem., Int. Ed. 2008, 47, 4268–4315. (h) Belmont, P.; Parker, E. Eur. J. Org. Chem. 2009, 6075–6089. (i) Wang, S.; Zhang, G.; Zhang, L. Synlett 2010, 692–706. (j) Krause, N.; Winter, C. Chem. Rev. 2011, 111, 1994–2009. (k) Pradal, A.; Toullec, P. Y.; Michelet, V. Synthesis 2011, 1501–1514.

^{(4) (}a) Gorin, D. J.; Toste, F. D. *Nature* **2007**, *446*, 395–403. (b) Leyva-Pérez, A.; Corma, A. *Angew. Chem., Int. Ed.* **2012**, *51*, 614–635.

^{(5) (}a) For a specific review on ligand effects in homogeneous Au catalysis, see: Gorin, D. J.; Sherry, B. D.; Toste, F. D. *Chem. Rev.* **2008**, *108*, 3351–3378. (b) For a relevant recent example, see: Wang, W.; Hammond, G. B.; Xu, B. J. Am. Chem. Soc. **2012**, *134*, 5697–5705.

the catalytic processes in which gold is involved.⁵ In this context, gold complexes with *N*-heterocyclic carbene (NHC) ligands are gaining a great significance due to the stability provided by the strong σ -donation and weaker π -back-bonding ability of such ligands in comparison to the more commonly used tertiary phosphines.⁶

On the other hand, since the discoveries made by Breslow and Grieco in the early 1980s on the positive effect of water on the rate and endo/exo selectivity of Diels-Alder reactions,⁷ the development of organic transformations in aqueous media has become one of the major cornerstones in modern chemistry.⁸ In addition, the use of water as an alternative, available, safe, and cost-effective solvent fullfills the principles of "Green Chemistry",⁹ creating an answer to the growing concerns associated with the environmental impact of chemical processes.¹⁰ Following this general trend, several Au-catalyzed reactions conducted in aqueous media have been successfully developed in recent years.¹ However, concerning the use of gold-NHC systems, studies in water have been scarce and restricted to catalytic hydrations of alkynes¹¹ and nitriles¹² and Meyer-Schuster-type rearrangements.13

We wish therefore to describe herein that, despite the high tendency of gold to promote the hydration of C=C bonds,^{1-3,11,14} cycloisomerization reaction of γ -alkynoic acids can be selectively performed in aqueous media in the presence of a new water-soluble Au(III)-NHC complex.¹⁵

(9) (a) Anastas, P. T.; Warner, J. C. In *Green Chemistry: Theory and Practice*; Oxford University Press: Oxford, 1998. (b) Sheldon, R. A.; Arends, I.; Hanefeld, U. *Green Chemistry and Catalysis*; Wiley-VCH: Weinheim, 2007.

(10) Kerton, F. M. In *Alternative Solvents for Green Chemistry*; RSC Publishing: Cambridge, 2009.

(12) Ramón, R. S.; Marion, N.; Nolan, S. P. Chem.—Eur. J. 2009, 15, 8695–8697.

(14) For leading references using Au(III) catalysts, see: (a) Casado, R.; Contel, M.; Laguna, M.; Romero, P.; Sanz, S. J. Am. Chem. Soc. **2003**, *125*, 11925–11935. (b) Wang, W.; Xu, B.; Hammond, G. B. J. Org. Chem. **2009**, *74*, 1640–1643.

(15) Competitive hydration of the substrates is probably the major limitation when catalytic transformations of alkynes are performed in water: (a) Chen, L.; Li, C.-J. *Adv. Synth. Catal.* **2006**, *348*, 1459–1484. (b) Li, C.-J. *Acc. Chem. Res.* **2010**, *43*, 581–590.

Such a catalytic reaction constitutes nowadays the major route for the construction of five-membered exocyclic enol-lactones with complete atom economy. It is also worthy of note that, although a plethora of transition-metal catalysts including gold have been developed for this synthetically useful transformation,^{16–18} efficient systems able to operate in aqueous media are still rare.¹⁹

The route designed to synthesize the water-soluble Au-(III)-NHC catalyst **3** is depicted in Scheme 1. First, the novel zwitterionic imidazolium salt **2** was prepared by reacting the known 2-(1*H*-imidazol-1-ylmethyl)pyridine 1^{20} with 1,3-propane sultone.²¹ Then, sequential treatment of **2** with Ag₂O, [AuCl(SMe₂)], and excess of PhICl₂ led to the clean formation of **3**, *via* the corresponding Ag(I)- and Au(I)-NHC intermediates.²² Complex **3**, isolated as an airstable yellow solid in 63% yield, was characterized by means of standard spectroscopic techniques as well as elemental analysis. In addition, X-ray-quality crystals were obtained by slow diffusion of toluene into a saturated DMSO solution, allowing us to confirm unambiguously the molecular structure of this new Au(III)-NHC derivative (see Supporting Information).

As expected, complex **3** readily dissolves in water at room temperature. However, in this medium an equilibrium is established between **3** and the zwitterionic derivative **4**, the latter resulting from the release of HCl and concomitant coordination of the pyridyl unit to the gold

(17) For other cycloisomerization reactions based on Au catalysts, see: (a) Harkat, H.; Weibel, J.-M.; Pale, P. *Tetrahedron Lett.* **2006**, *47*, 6273–6276. (b) Marchal, E.; Uriac, P.; Legouin, B.; Toupet, L.; van de Weghe, P. *Tetrahedron* **2007**, *63*, 9979–9990. (c) Harkat, H.; Dembelé, A. Y.; Weibel, J.-M.; Blanc, A.; Pale, P. *Tetrahedron* **2009**, *65*, 1871–1879. (d) Sperger, C. A.; Fiksdahl, A. J. Org. Chem. **2010**, *75*, 4542–4553.

(18) For reviews on catalytic additions of carboxylic acids to alkynes, see: (a) Alonso, F.; Beletskaya, I. P.; Yus, M. Chem. Rev. 2004, 104, 3079–3160. (b) Nakamura, I.; Yamamoto, Y. Chem. Rev. 2004, 104, 2127–2198. (c) Beller, M.; Seayad, J.; Tillack, A.; Jiao, H. Angew. Chem., Int. Ed. 2004, 43, 3368–3398. (d) Jiménez-Nuñez, E.; Echavarren, A. M. Chem. Commun. 2007, 333–346. (e) Bruneau, C.; Dixneuf, P. H. Acc. Chem. Res. 1999, 32, 311–323.

(19) Only two catalytic systems, involving Pt and Cu complexes, active in aqueous media have been previously described: (a) Mindt, T. L.; Schibli, R. *J. Org. Chem.* **2007**, *72*, 10247–10250. (b) Alemán, J.; del Solar, V.; Navarro-Ranninger, C. *Chem. Commun.* **2010**, *46*, 454–456. (c) The presence of water was tolerated in ref 16c.

(20) Chiu, P. L.; Lai, C.-L.; Chang, C.-F.; Hu, C.-H.; Lee, H. M. Organometallics 2005, 24, 6169–6178.

(21) Related zwitterionic imidazolium salts have been previously synthesized following this route: Moore, L. R.; Cooks, S. M.; Anderson, M. S.; Schanz, H.-J.; Griffin, S. T.; Rogers, R. D.; Kirk, M. C.; Shaughnessy, K. H. *Organometallics* **2006**, *25*, 5151–5158.

(22) Trace amounts of HCl in the dichloromethane employed is probably responsible for the protonation of the sulfonate group. In addition, HCl may also be present in the iodobenzene dichloride employed as oxidant, since it was synthesized by chlorination of iodobenzene with NaClO₂ in hydrochloric acid solution as described in Zhao, X.-F.; Zhang, C. *Synthesis* **2007**, *4*, 551–557.

⁽⁶⁾ For reviews on the catalytic uses of NHC-containing Au(I) and Au(III) complexes, see: (a) Marion, N.; Nolan, S. P. *Chem. Soc. Rev.* 2008, *37*, 1776–1782. (b) Nolan, S. P. *Acc. Chem. Res.* 2011, *44*, 91–100. (c) Gaillard, S.; Cazin, C. S. J.; Nolan, S. P. *Acc. Chem. Res.* 2012, published ASAP ahead of print; DOI: 10.1021/ar200188f).

^{(7) (}a) Rideout, D. C.; Breslow, R. J. Am. Chem. Soc. **1980**, 102, 7816–7817. (b) Grieco, P. A.; Yoshida, K.; Garner, P. J. Org. Chem. **1983**, 48, 3137–3139.

⁽⁸⁾ See, for example: (a) Organic Reactions in Water; Lindström, U. M., Ed.; Blackwell Publishing: Oxford, 2007. (b) Li, C.-J.; Chan, T. H. In Comprehensive Organic Reactions in Aqueous Media; Wiley-VCH: Weinheim, 2007. (c) Genin, E.; Leseurre, L.; Michelet, V. In Génie des Procédés Verts et Durables: Outils et Méthodes; Dunod: Paris, 2010. (d) Li, C.-J. Chem. Rev. 2005, 105, 3095–3166. (e) Shaughnessy, K. H. Chem. Rev. 2009, 109, 643–710 and references therein.

^{(11) (}a) Schneider, S. K.; Herrmann, W. A.; Herdtweck, E. Z. Anorg. All. Chem. 2003, 629, 2363–2370. (b) de Frémont, P.; Singh, R.; Stevens, E. D.; Petersen, J. L.; Nolan, S. P. Organometallics 2007, 26, 1376–1385.
(c) Marion, N.; Ramón, R. S.; Nolan, S. P. J. Am. Chem. Soc. 2009, 131, 448–449. (d) Almássy, A.; Nagy, C. E.; Bényei, A. C.; Joó, F. Organometallics 2010, 29, 2484–2490. (e) Czégéni, C. E.; Papp, G.; Kathó, Á.; Joó, F. J. Mol. Catal. A: Chem. 2011, 340, 1–8.

^{(13) (}a) Ramón, R. S.; Marion, N.; Nolan, S. P. *Tetrahedron* **2009**, *65*, 1767–1773. (b) Ramón, R. S.; Gaillard, S.; Slawin, A. M. Z.; Porta, A.; D'Alfonso, A.; Zanoni, G.; Nolan, S. P. *Organometallics* **2010**, *29*, 3665–3668. (c) Merlini, V.; Gaillard, S.; Porta, A.; Zanoni, G.; Vidari, G.; Nolan, S. P. *Tetrahedron Lett.* **2011**, *52*, 1124–1127.

⁽¹⁶⁾ Representative examples published by some of us: (a) Genin, E.; Toullec, P. Y.; Antoniotti, S.; Brancour, C.; Genêt, J.-P.; Michelet, V. J. Am. Chem. Soc. 2006, 128, 3112–3113. (b) Genin, E.; Toullec, P. Y.; Antoniotti, S.; Brancour, C.; Genêt, J.-P.; Michelet, V. ARKIVOC 2007, 67–78. (c) Toullec, P. Y.; Genin, E.; Antoniotti, S.; Genêt, J.-P.; Michelet, V. Synlett 2008, 707–711. (d) Neaţu, F.; Li, Z.; Richards, R.; Toullec, P. Y.; Genêt, J.-P.; Dumbuya, K.; Gottfried, J. M.; Steinrück, H.-P.; Pârvulescu, V. I.; Michelet, V. Chem.—Eur. J. 2008, 14, 9412– 9418. (e) Neaţu, F.; Pârvulescu, V. I.; Michelet, V.; Genêt, J.-P.; Goguet, A.; Hardacre, C. New J. Chem. 2009, 33, 102–106. (f) Neaţu, F.; Proteşescu, L.; Florea, M.; Pârvulescu, V. I.; Teodorescu, C. M.; Apostol, N.; Toullec, P. Y.; Michelet, V. Green Chem. 2010, 12, 2145– 2149.

center (Scheme 1). Complex **4** is the major species present in solution at rt, and its structure was also determined by X-ray crystallography (see Supporting Information). Chelation of the NHC ligand in water is readily evidenced by ¹H NMR spectroscopy, where the singlet signal corresponding to the methylenic protons of the picolyl moiety in **3** (δ 5.63 ppm) undergoes an AB splitting (δ 5.65 and 5.81 ppm; J = 16.0 Hz) as a consequence of the conformational rigidity of the six-membered metallacycle in **4**.²³





As shown in Table 1, complex 3 turned out to be an efficient and selective catalyst for the cycloisomerization of model 4-pentynoic acid 5a in different mono- and biphasic water/organic solvent mixtures (entries 1-4). Reactions, performed at rt under aerobic conditions with a metal loading of 2.5 mol %, led to the quantitative transformation of 5a into enol-lactone 6a, with no byproduct resulting from the hydration of the C \equiv C bond of **5a** being detected by ¹H NMR spectroscopy in the crude reaction mixtures. Best results in terms of activity were obtained using a biphasic toluene/water solvent system,²⁴ where total consumption of 5a was observed after only 1 h, thus allowing the isolation of **6a** in high yield (81%; entry 4). The process is also operative in pure water (entry 5). However, both the activity and selectivity of 3 in this medium were lower, the latter due to the partial hydrolysis of 6a to form 3acetylpropanoic acid.²⁵ Brønsted acid catalysis mediated by the HCl released after dissolving complex 3 in water can

be ruled out, as demonstrated by the result shown in entry $6.^{26}$ The use of a base (KOH, entry 7) associated with the catalyst allowed the formation of the desired lactone.^{17a} Conversely, no catalytic activity was observed in the presence of the free ligand **2** alone (entry 8). These results fully support that the intramolecular nucleophilic attack of the carboxylic unit is taking place upon coordination of the alkyne to the metal center. The required vacant site is most probably generated by decoordination of the picolyl moiety in the *in situ* formed complex **4**, owing to the well-known hemilabile character of this group.²⁷ As shown in entries 9 and 10, simple Au(III) and Au(I) sources were comparatively much less effective and selective, thus highlighting the synthetic potential of complex **3**.

Table 1. Catalytic Cycloisomerization of 5a in Aqueous Media^a



entry	conditions	time (h)	yield $(\%)^e$
1^b	3 (2.5 mol %), MeCN/H ₂ O	6	>99
2^b	3 (2.5 mol %), Et ₂ O/H ₂ O	5	>99
3^b	3 (2.5 mol %), CH ₂ Cl ₂ /H ₂ O	2	>99
4^b	3 (2.5 mol %), toluene/H ₂ O	1	>99 (81)
5^c	3 (2.5 mol %), H ₂ O	1	50^{f}
6^b	$HCl (2.5 \text{ mol } \%)$, toluene/ H_2O	1	1
$7^{b,d}$	3 + KOH (2.5 mol %), toluene/H ₂ O	1	>99
8^b	2 (2.5 mol %), toluene/H ₂ O	1	0
9^b	AuCl ₃ (2.5 mol %), toluene/H ₂ O	4	90^g
10^b	AuCl (2.5 mol %), toluene/ H_2O	6	70^{f}

^{*a*}Reactions were performed at rt under aerobic conditions starting from 0.3 mmol of **5a**. ^{*b*} 1 mL of the corresponding organic solvent and 1 mL of distilled water were employed. ^{*c*} 2 mL of distilled water were employed. ^{*d*} 2.5 mol % of each reagent was employed. ^{*e*} Determined by ¹H NMR. Isolated yields are given in brackets. ^{*f*} A mixture of **6a** and 3-acetylpropanoic acid in ca. 3:1 ratio is formed. ^{*g*} A mixture of **6a** and 3-acetylpropanoic acid in ca. 5:1 ratio is formed.

The scope of the reaction was then studied for various γ -alkynoic acids in a biphasic toluene/water mixture (Table 2).²⁸ Complete conversions were observed for substrates **5b**-**h**, all bearing a terminal alkyne unit and alkyl or alkenyl side chains, after 2–4 h of stirring at rt in the presence of 2.5 mol % of **3** (entries 1–7). The reactions led to the selective formation of lactones **6b**-**h**, which could be isolated in good yields (66–97%), without detection of any byproduct by ¹H NMR in the crude reaction mixtures. As shown in entries 8 and 9, the process is also operative with the internal alkynes **5i**-**j**. Nevertheless, longer reaction times or higher temperatures were required. In addition, traces of the corresponding six-membered ring lactones, resulting from an *endo* instead of an *exo*

⁽²³⁾ Related Au(III) complexes containing chelating picolyl-NHC ligands have been described. See, for example: (a) Pažický, M.; Loos, A.; Ferreira, M. J.; Serra, D.; Vinokurov, N.; Rominger, F.; Jäkel, C.; Hashmi, A. S. K.; Limbach, M. *Organometallics* **2010**, *29*, 4448–4458. (b) Topf, C.; Hirtenlehner, C.; Fleck, M.; List, M.; Monkowius, U. Z. Anorg. Allg. Chem. **2011**, *637*, 2129–2134. (c) See also ref 20.

⁽²⁴⁾ For examples of improved selectivity when biphasic toluene/ water systems are employed, see: (a) Genin, E.; Michelet, V.; Genêt, J.-P. *Tetrahedron Lett.* **2004**, *45*, 4157–4161. (b) Genin, E.; Michelet, V.; Genêt, J.-P. J. Organomet. Chem. **2004**, *689*, 3820–3830 and references therein.

⁽²⁵⁾ Formation of 3-acetylpropanoic acid by direct hydration **5a** was discarded since, under the same reaction conditions, complex **3** is not able to hydrate terminal alkynes, such as 1-pentyne or 1-hexyne (only at 100 °C hydration could be observed). Note also that hydrolysis of lactones is a well-known process: Olson, A. R.; Hyde, J. L. J. Am. Chem. Soc. **1941**, *63*, 2459–2461.

⁽²⁶⁾ For a review on π -activation of alkynes by Brønsted acids, see: Yamamoto, Y.; Gridner, I. D.; Patil, N. T.; Jin, T. *Chem. Commun.* **2009**, 5075–5087.

⁽²⁷⁾ Slone, C. S.; Weinberger, D. A.; Mirkin, C. A. Prog. Inorg. Chem. 1999, 48, 233–350 and references therein.

Table 2. Au(III)-Catalyzed Cycloisomerization of γ -Alkynoic Acids **5b**-**j** in a Biphasic Toluene/Water Medium^{*a*}



entry	y substrate	time (h)	yield $(\%)^b$
1	$R^1 = R^2 = Me; R^3 = H(5b)$	2	84
2	$\mathbf{R}^1 = \mathbf{CO}_2 \mathbf{Me}; \mathbf{R}^2 = \mathbf{R}^3 = \mathbf{H} \left(\mathbf{5c} \right)$	2	96
3	$R^1 = CO_2Me; R^2 = Me; R^3 = H(5d)$	2	80
4	$R^1 = CO_2Me; R^2 = Et; R^3 = H(5e)$	2	97
5	$R^1 = CO_2Et; R^2 = Bn; R^3 = H(5f)$	2	92
6	$R^1 = CO_2Me; R^2 = CH_2CH = CH_2; R^3 = H(5g)$	2	84
7	$R^1 = CO_2Me; R^2 = CH_2CH=CHPh; R^3 = H$	4	66
	(5h)		
8	$R^1 = CO_2Me; R^2 = Et; R^3 = Me(5i)$	24	94^c
9^d	$R^1 = CO_2Me; R^2 = H; R^3 = Ph(5j)$	2	28^c

^{*a*} Reactions were performed as described in ref 28. ^{*b*} Isolated yields. ^{*c*} Traces of the six-membered ring lactone were detected by ¹H NMR in the crude reaction mixture. ^{*d*} Reaction performed at 80 °C.

cyclization, were now detected in the ¹H NMR spectra of the crude mixtures.

Complex 3 proved also selective in the cyclization of more challenging bispropargylic carboxylic acids (Table 3).²⁹ Thus, starting from the symmetrical terminal 1,6-diynes $7\mathbf{a} - \mathbf{c}$ (entries 1-3), enol-lactones $8\mathbf{a} - \mathbf{c}$ containing a propargylic side arm intact could be generated in excellent yields (84-94%) under mild conditions. In contrast, desymmetrization of diynes 7d-f containing internal C=C bonds proved more difficult, and a higher temperature (80 °C) was required (entries 4-6). As in the case of 5i-j, competitive formation of the corresponding sixmembered ring lactones was again observed. The remarkable higher reactivity of terminal versus internal C=C bonds was clearly evidenced in the reactions of the nonsymmetrical 1,6-divnes 7g-h (entries 7 and 8), where only the former participated in the cyclization process leading to the exclusive formation of lactones 8g-h.³⁰ In none of

(29) For related transformations catalyzed by NHC-gold complexes in organic media see ref 17d.

Table 3. Au(III)-Catalyzed Cycloisomerization of Bispro-
pargylic Carboxylic Acids $7\mathbf{a}-\mathbf{h}$ in a Biphasic Toluene/Water
Medium^a



entry	substrate	temp	time (h)	yield $(\%)^b$
1	$R^1 = R^2 = R^3 = H(7a)$	rt	2	94
2	$R^1 = CO_2Me; R^2 = R^3 = H(7b)$	\mathbf{rt}	2	84
3	$R^1 = Ph; R^2 = R^3 = H(7c)$	\mathbf{rt}	2	88
4	$R^1 = H; R^2 = R^3 = Me(7d)$	80 °C	2	84^c
5	$R^1 = CO_2Me; R^2 = R^3 = Me(\mathbf{7e})$	80 °C	1	90^d
6	$R^1 = CO_2Me; R^2 = R^3 = Ph(7f)$	80 °C	1	40
7	$R^1 = CO_2Me; R^2 = H; R^3 = Me(7g)$	\mathbf{rt}	2	80
8	$R^{1} = CO_{2}Me; R^{2} = H; R^{3} = Ph(\mathbf{7h})$	rt	6	86

^{*a*} Reactions were performed as described in ref 28. ^{*b*} Isolated yields. ^{*c*} A mixture of **8d** and the corresponding six-membered ring lactone is formed in ca. 6:1 ratio. ^{*d*} A mixture of **8e** and the corresponding sixmembered ring lactone is formed in ca. 3:1 ratio.

these desymmetrization processes were hydration byproducts detected.

One of the major advantages of biphasic catalysis is that it is associated with the recycling of the catalytic species.³¹ In this context, we have found that, after separation of the toluene layer in which the lactone is soluble, the aqueous phase containing the catalytically active gold species can be reused. Thus, using the 1,6-diyne **7a** as model substrate, 10 consecutive reactions could be performed without any loss of activity or selectivity (quantitative yield over the 10 cycles).

In summary, we have demonstrated that non-hydrative transformations of functionalized alkynes can be efficiently and selectively performed in aqueous media by means of NHC-gold catalysis. This unprecedented fact, together with the possibility of an effective recycling, opens new research directions to expand the applications of NHC-gold complexes in synthetic organic chemistry.

Acknowledgment. This work was financially supported by MICINN (projects CTQ2010-14796/BQU, CTQ2010-17476/BQU, and CSD2007-00006) and Junta de Andalucia (project FQM-3151) of Spain and CNRS of France. E.T.-M. thanks MECD of Spain and the European Social Fund for the award of a FPU grant.

Supporting Information Available. Experimental details, characterization data, and crystallographic data for compounds 3 and 4 in CIF format. This material is available free of charge via the Internet at http://pubs.acs.org.

⁽²⁸⁾ General procedure for the catalytic reactions. To a biphasic system composed of 1 mL of toluene and 1 mL of distilled water were added 0.3 mmol of the corresponding alkynoic acid and 4.4 mg of complex 3 (0.0075 mmol; 2.5 mol %). The resulting mixture was stirred under air, at rt or 80 °C (see Tables 1–3), until complete consumption of the alkynoic acid was observed by TLC. The organic phase was then separated, the aqueous one was extracted with diethyl ether (2×2 mL), and the combined organic extracts were dried over anhydrous MgSO₄ and filtered over a short pad of silica gel using CH₂Cl₂ as eluent. The volatiles were removed under vacuum to yield the corresponding lactone in high yield and purity. When required, lactones were further purified by column chromatography over silica gel using hexane/EtOAc (9:1) as eluent.

⁽³⁰⁾ From a mechanistic point of view, the cyclization processes reported herein proceed via an *anti*-addition of the carboxylic unit to the π -coordinated alkyne. This was inferred by performing the cyclization of **7a** into lactone **8a** in a biphasic toluene/D₂O mixture, reaction that led to the major incorporation of deuterium in *E* position of the exocyclic C=C bond of **8a** (details are given in the Supporting Information).

⁽³¹⁾ Aqueous-Phase Organometallic Catalysis: Concepts and Applications; Cornils, B., Herrmann, W. A., Eds.; Wiley-VCH: Weinheim, Germany, 1998.

The authors declare no competing financial interest.