

Through-space interactions between face-to-face, center-to-edge oriented arenes: importance of polar- π effects

Franco Cozzi,^{*a} Rita Annunziata,^a Maurizio Benaglia,^a Mauro Cinquini,^a Laura Raimondi,^a Kim K. Baldrige^b and Jay S. Siegel^{*b}

^a CNR-ISTM and Dipartimento di Chimica Organica e Industriale, Università di Milano, via Golgi, 19-20133 Milano, Italy

^b Department of Chemistry and Biochemistry -0358 University of California, San Diego, 9500 Gilman Drive, La Jolla, California 92093-0358, USA

Received 11th September 2002, Accepted 5th November 2002

First published as an Advance Article on the web 6th December 2002

Two series of conformationally restricted polycyclic compounds (**1–3** and **4–7**) have been synthesized as model systems for studying the through-space interactions between face-to-face, center-to-edge (parallel-offset) oriented arenes. These compounds feature different X substituents on one of the interacting rings. By monitoring the variation of the ΔG^\ddagger for the rotation around the aryl-aryl bond in **1–7** as a function of X by 2D [¹H, ¹H] EXSY NMR spectroscopy, it was found that the barriers increase on passing from electron-donating to electron-withdrawing substituted derivatives. Quantum mechanical calculations [MP2/DVZ (2d,p)//B3LYP/DVZ(2d,p)] gave barrier values and variations in agreement with the experimental data. The results are consistent with a repulsive arene-arene interaction dominated by electrostatic effects.

Introduction

Through-space interactions between π -systems¹ are important because they play a major role in controlling disparate chemical phenomena including molecular recognition,² stereoselectivity of organic reactions,³ nucleic acid and protein macroscopic structures,⁴ aggregation of extended π -systems,⁵ and crystal packing.⁶ Several theoretical studies have been aimed at elucidating the nature of the arene-arene interactions in the simple case of the benzene dimer (Fig. 1).^{1,7} From these works emerged




Fig. 1 Relative orientations of rings in the benzene dimer.

the idea that the parallel-stacked orientation **A** is unfavorable, the edge-to-face one **B** is favorable,⁸ and the energy of the face-to-face, center-to-edge (parallel-offset) arrangement **C** depends on the contact surface area. A simple picture of the electronic distribution of benzene possessing a negatively charged core surrounded by a positively charged periphery,^{7b,7i} rationalizes these results on an electrostatic basis.

The electrostatic nature of the repulsive interaction in the parallel-stacked orientation **A** has experimentally been demonstrated by us by studying a number of variously substituted 1,8-diarylnaphthalenes,⁹ and confirmed by work carried out on other systems.¹⁰ These findings can be invoked to rationalize a variety of experimental results in different fields,^{2–5} and nicely explain the tendency of the mixed arene-perfluoroarene systems to adopt the parallel-stacked orientation.¹¹

The experimental work has also been devoted to elucidate the type **B** interaction between edge-to-face oriented arenes.^{12,13} The results of a quantitative study led Hunter and coworkers to conclude that “the interactions between the CH groups of the edge ring and the π electron density of the face ring are sensitive to changes in the local charge distribution on the two rings”.^{12a,b} In contrast, Wilcox and coworkers^{12c} designed a model system

that was unresponsive to changes in arene substitution, a result that “casts substantial doubt on the importance of electrostatic interactions in edge-to-face binding events.”^{12c}

Face-to-face, center-to-edge interactions (type **C**), also called parallel-offset interactions, have received considerably less attention, despite the fact that this structural motif can be found in many structures.¹⁴ Recently,^{15,16} some conformationally flexible derivatives of *N*-benzyl-2-phenylpyridinium bromide adopting, among others, a parallel-offset conformation in water, have been reported. Herein, we investigate type **C** interactions in two conformationally restricted polycyclic systems.

Results and discussion

Synthesis. The model compounds **1–7** (Chart 1) were selected





Chart 1 Structures of compounds **1–7**.

to study type **C** interactions. These systems feature a reference ring with an electronic distribution that is constant across the series (*i.e.* the left-hand ring of the benzophenanthrene skeleton), and a rotating phenyl ring carrying electron-donating (ED) and electron-withdrawing (EW) substituents and partially overlapping the reference ring. The synthetic sequence for their preparation (Scheme 1) was planned so that **1–7** could be obtained from a common advanced intermediate.

Ullmann reaction between commercially available 2-bromo-3-nitrotoluene and 1-bromo-2-methylnaphthalene (3 mol equiv.)¹⁷ gave the cross-coupling product **8** in 43% yield. Reduction of the nitro group (87% yield) followed by Sandmeyer



Scheme 1 Synthesis of compounds 1–7. *Reagents and conditions:* a, Cu bronze, neat, 200 °C. b, SnCl₂·H₂O, AcOEt, 80 °C, 15 h. c, H₂SO₄, NaNO₂, H₂O, 0–5 °C, 30 min; then KI 5 to 20 °C, 15 h. d, NBS cat. PhCO₂O, *hν*, CCl₄, 80 °C, 48 h. e, AgNO₃, dioxane, H₂O, 100 °C, 15 h. f, SmI₂, THF, –78 °C, 1 h. g, Pd₂(dba)₃, AsPh₃, DMF, 150 °C, 40 h. h, PySO₃, DMSO, RT, 2 h.

reaction of the resulting amine afforded the iodo compound **9** in 57% yield. Perbromination with excess NBS (5.5 mol equiv.) gave the bis(dibromomethyl) derivative (87% yield) that was then hydrolyzed to the corresponding bis(aldehyde) **10** by silver nitrate promoted reaction in dioxane–water (70% yield). Reductive intramolecular coupling carried out under McMurry conditions¹⁸ to generate 1-iodobenzo[*c*]phenanthrene led to concomitant cleavage of the carbon–iodine bond. An attempt to prepare the same compound by reaction of **10** with hydrazine in acetic acid¹⁹ failed, as did the synthesis of 1-iodo-5,6-epoxybenzo[*c*]phenanthrene by reaction of **10** with tris(dimethylaminophosphine) in toluene.²⁰ Ultimately, SmI₂²¹ promoted pinacol reaction of **10** in THF at –78 °C gave the racemic diequatorial diol **11** (*J* = 10.5 Hz) in 75% yield.²²

The reaction of **11** with 4*X*-substituted phenyl trimethylstannanes (4 mol equiv.)²³ in the presence of 0.2 mol equiv. of Pd₂(dba)₃ and 0.8 mol equiv. of AsPh₃²⁴ in degassed DMF

Table 1 ΔG^\ddagger for the rotation around the aryl–aryl bond in compounds 1–7 as determined by 2D EXSY ¹H NMR in THF-*d*₈ at the indicated temperature

Compound (type)	X substituent	ΔG^\ddagger /kJ mol ⁻¹	T/K
1 (diol)	MeO	47.9	195
2 (diol)	H	48.7	195
3 (diol)	NO ₂	50.9	210
4 (dione)	MeO	51.0	210
5 (dione)	Me	51.7	210
6 (dione)	H	52.4	205
7 (dione)	F	53.1	210

(150 °C, 40 h) gave diols **1–3**, albeit in low yields (**1**, 29%; **2**, 12%; **3**, 11%). Surprisingly, **1** and **2** were accompanied by comparable amounts of the corresponding diketones **4** (24% yield) and **6** (11% yield). Starting from 4-methylphenyl trimethylstannane, diketone **5** was the only isolated product (15% yield). Finally, fluorodiketone **7** was obtained from the corresponding crude diol **12** (that could not be obtained in pure form) by Py·SO₃ oxidation²⁵ in 12% overall yield from **11**.

Spectroscopy. The 300 MHz ¹H NMR spectra of **1–7** in THF-*d*₈ at RT revealed broad peaks for the protons of the 4*X*-substituted phenyl ring. Upon cooling the sample to about 200 K distinct resonances were observed with an average $\Delta\delta$ of 1.55 and 0.80 ppm for the *ortho* and *meta* protons, respectively. This result is consistent with rotation around the aryl–aryl bond in **1–7** occurring on an NMR timescale comparable to that of $\Delta\nu$ at RT but significantly more slowly at lower temperatures. The large $\Delta\delta$ seen for the *ortho* and *meta* protons suggests a preferred conformation where the 4*X*-substituted phenyl ring is frozen with one side of the ring above the shielding cone of the terminal ring of the benzophenanthrene system, *i.e.* a face-to-face center-to-edge conformation.²⁶

The ΔG^\ddagger for the rotation around the aryl–aryl bond in the series of compounds **1–3** and **4–7** was determined by 2D [¹H, ¹H] EXSY NMR experiments (Table 1).^{27,28} The data for the diol and, independently, the diketone series show a trend for the barrier to rotation to increase from ED to EW *X* substituents on the phenyl ring. The plots of ΔG^\ddagger values vs the σ_{para} for **1–3** and **4–7** showed excellent linear relationship (Fig. 2).




Fig. 2 Plots of ΔG^\ddagger (kJ mol⁻¹) vs. σ_{para} for compounds **1–3** ($R = 0.9999$) and **4–7** ($R = 0.9998$).

Also in type **A** interactions, the barrier to rotation around aryl–aryl bonds increased when the interacting rings were made less electron rich by the introduction of EW groups. This effect was apparently due to a decreased repulsion between the π electrons of the arenes at the ground state (GS) level.⁹ In compounds **1–7**, the distance between the interacting rings is longer and the overlap between the π clouds is less extended than in the type **A** analogs previously studied.⁹ Specifically, a typical angle between planes in **1–7** is *ca.* 55° with a center-to-center distance of 4.1 Å (closest contact 3.2 Å; furthest contact 5.2 Å). Thus, the repulsion between the π electrons is lower and the substituent effects should be less dramatic than previously seen. Indeed, the difference in the barriers between **1** and **3** is 3.0 kJ mol⁻¹,

while in the 1,8-diarylnaphthalene case a similar change in substitution led to $\Delta\Delta G^\ddagger$ values of up to 14 kJ mol⁻¹.^{9a} At the GS level compounds 1–7 should experience an attractive electrostatic interaction between the hydrogens of one ring and the π electrons of the other. However, the effect of different substituents on this interaction should be small,^{12a} and the contribution of the CH– π factor to the overall interaction likely negligible.

Computation methods. Structural computations were performed with hybrid density functional methods (HDFT) using GAUSSIAN98.²⁹ The HDFT method employed Becke's 3 parameter functional³⁰ in combination with nonlocal correlation provided by the Lee–Yang–Parr expression³¹ that contains both local and nonlocal terms, B3LYP. Dunning's DZV(2d,p)³² double- ζ valence basis set was employed. These levels of theory have been previously shown to be reliable for structural determination in these types of compounds.³³ Single point energy computations were performed on optimized structures using MP2³⁴ dynamic correlation treatment giving superior energetic analysis.

Quantum mechanical calculations [MP2/DVZ (2d,p)//B3LYP/DVZ(2d,p)] on diketone **6** showed a centroid to centroid distance of 4.3 Å and a type C conformation between the reference and rotating rings in the GS. At the transition state (TS, 1 negative eigenvalue) calculations showed that one of the *ortho* hydrogens of the rotating ring points towards the carbon atom rim of the reference ring with a distance of *ca.* 2.2 Å; the same hydrogen is 2.8 Å from the center of the ring, significantly further than the rim-atom interaction. The computational barrier (TS–GS) is 48.2 kJ mol⁻¹, in reasonable agreement with experiment (52.4 kJ mol⁻¹). Thus the computation model supports the supposition that we are studying an interaction loosely analogous to type C in the series 1–7 and that this series behaves similarly to the type A analogs studied previously.⁹

Two additional derivatives of **6** were calculated: 6-NH₂ and 6-NO₂. They provide some perspective on the trends across a Hammett series. The predicted barriers to rotation for 6-NH₂, **6**, and 6-NO₂ are 47.5, 48.2, and 49.0 kJ mol⁻¹, respectively. Albeit a small effect, the barrier increases with decreasing donor character of the *para* substituent. This trend parallels the one experimentally observed for both the diketone and diol series, and the magnitude of the barriers is in accord with those measured by NMR, showing that the type C interactions can be modeled well by this level of theory.

Conclusions

In conclusion, both theory and experiments agree in indicating that aryl rotation is responsible for the spectral changes observed by the variable temperature NMR spectroscopy, and that the through-space (type C) interactions between parallel-offset oriented arenes is strongly influenced by electrostatic effects. Since these effects were also found to be important in the type A and type B interactions between parallel-stacked and center-to-edge oriented arenes, respectively, the results herein described lend additional support to the notion that polar- π interactions are decisive factors for understanding the arene–arene interactions.

Experimental

General. ¹H NMR spectra were recorded at 300 MHz in chloroform-*d* (CDCl₃) and were referenced to tetramethylsilane (TMS) at 0.00 ppm. ¹³C NMR spectra were recorded at 75 MHz and were referenced to 77.0 ppm in CDCl₃. ¹⁹F NMR spectra were recorded at 282 MHz in CDCl₃ and were referenced to hexafluorobenzene at 0.0 ppm. Variable temperature ¹H NMR spectra were recorded at 300 MHz in THF-*d*₆. IR spectra were recorded on thin films or as a solution in CH₂Cl₂. The

4-X-phenyl trimethylstannanes required for the synthesis of 1–7 were obtained from the corresponding 4-X substituted iodo-benzenes following the reported procedure,²³ and were used as crude products.

2-Methyl-1-(2-methyl-6-nitrophenyl)naphthalene 8

To a stirred mixture of 1-bromo-2-methylnaphthalene (2.16 mL, 13.88 mmol) and copper bronze (2.93 g, 46.2 mmol) heated up to 200 °C, 2-bromo-3-nitrotoluene (1.0 g, 4.62 mmol) was added portionwise over a period of 6 h. After the end of the addition the mixture was stirred at 200 °C for 20 h, whereupon it was cooled to RT and acetone (20 mL) was added. The suspension was stirred for 1 h at RT and then filtered. The filtrate was concentrated under vacuum and the resulting crude product was purified by flash chromatography with a 95 : 5 hexanes : Et₂O mixture as eluant. The product, an orange thick oil, was isolated in 43% yield (0.556 g). IR (thin film): 3048, 2940, 1640, 1380, 1265 cm⁻¹. ¹H NMR: δ 7.87 (d, *J* = 7.9 Hz, 1H, hydrogen *ortho* to NO₂), 7.84 (dd, *J* = 8.9, 1.3 Hz, 1H, H–C5 of naphthalene), 7.81 (d, *J* = 8.9 Hz, 1H, H–C4 of naphthalene), 7.61 (d, *J* = 7.9 Hz, 1H, hydrogen *para* to NO₂), 7.50 (t, *J* = 7.9 Hz, 1H, hydrogen *meta* to NO₂), 7.47 (d, *J* = 8.9 Hz, 1H, H–C3 of naphthalene), 7.40 (dt, *J* = 8.9, 1.3 Hz, 1H, H–C6 of naphthalene), 7.31 (dt, 8.9, 1.3 Hz, 1H, H–C7 of naphthalene), 7.09 (dd, *J* = 8.9, 1.3 Hz, 1H, H–C8 of naphthalene), 2.15 (s, 3H, Me of naphthalene), 1.90 (s, 3H, Me of phenyl group); ¹³C NMR: δ 148.0, 140.1, 134.3, 133.5, 133.2, 132.0, 131.9, 131.7, 128.5, 128.3, 128.25, 128.2, 126.5, 125.1, 124.3, 121.7, 20.1, 19.8. C₁₈H₁₅NO₂ requires: C, 77.96; H, 5.45; N, 5.05%; found: C, 77.77; H, 5.58; N, 4.97%.

3-Methyl-2-(2-methyl-1-naphthyl)aniline

A mixture of compound **8** (0.979 g, 3.53 mmol) and SnCl₂ dihydrate (4.0 g, 17.65 mmol) in refluxing ethyl acetate (50 mL) was stirred for *ca.* 15 h. When TLC analysis showed the disappearance of the starting material, the mixture was cooled to RT and made slightly alkaline by the careful addition of a saturated aqueous solution of NaHCO₃. The mixture was then filtered on a Celite cake, the organic phase was separated, and the aqueous phase was extracted with Et₂O (2 × 20 mL). The combined organic phases were dried over Na₂SO₄, filtered and concentrated under vacuum. The crude product was purified by flash chromatography with a 80 : 20 hexanes : Et₂O mixture as eluant. The product, a yellow thick oil, was isolated in 87% yield (0.759 g). IR (thin film): 3469, 3388, 3053, 2950, 1612, 1265 cm⁻¹. ¹H NMR: δ 7.92 (d, *J* = 8.0 Hz, 1H, H–C5 of naphthalene), 7.87 (d, 8.4 Hz, 1H, H–C4 of naphthalene), 7.53 (d, *J* = 8.4 Hz, 1H, H–C3 of naphthalene), 7.50–7.40 (m, 3H, H–C6, H–C7 and H–C8 of naphthalene), 7.23 (t, *J* = 7.7 Hz, 1H, hydrogen *meta* to NH₂), 6.86 (d *J* = 7.7 Hz, 1H, hydrogen *para* to NH₂), 6.74 (d, *J* = 7.7 Hz, 1H, hydrogen *ortho* to NH₂), 2.29 (s, 3H, Me of naphthalene), 1.88 (s, 3H, Me of phenyl group); ¹³C NMR: δ 144.2, 138.7, 134.8, 133.4, 132.6, 132.2, 129.0, 128.3, 128.1, 127.7, 126.4, 125.2, 125.0, 120.1, 112.8, 112.2, 24.0, 19.9. C₁₈H₁₇N requires: C, 87.41; H, 6.93; N, 5.66%; found: C, 87.67; H, 7.11; N, 5.49%.

2-Methyl-1-(2-iodo-6-methylphenyl)naphthalene 9

To a stirred suspension of the above described amine (0.73 g, 2.95 mmol) in concentrated sulfuric acid (0.75 mL) and water (0.2 mL) cooled at –10 °C, a solution of sodium nitrite (0.305 g, 4.43 mmol) in water (0.2 mL) was added. After 15 min stirring at –10 °C, a solution of KI (1.45 g, 8.85 mmol) in water (0.6 mL) was slowly added. The deep red solution was stirred overnight while allowing the temperature to slowly rise to RT. The mixture was made neutral by the addition of a saturated aqueous solution of NaHCO₃, and was then washed with a saturated aqueous solution of Na₂SO₃ (10 mL). The resulting

mixture was extracted with Et₂O (2 × 20 mL) and CH₂Cl₂ (2 × 20 mL). The combined organic phases were dried over NaSO₄, filtered, and concentrated under vacuum. The residue was purified by flash chromatography with hexanes as eluant. The product, a colorless oil, was obtained in 57% yield (0.602 g). IR (thin film): 3053, 2955, 1615, 1260 cm⁻¹. ¹H NMR: δ 7.91–7.86 (m, 2H, hydrogen *ortho* to I and H–C5 of naphthalene), 7.84 (d, *J* = 8.6, 1H, H–C4 of naphthalene), 7.44 (d, *J* = 8.6, 1H, H–C3 of naphthalene), 7.42 (dt, *J* = 8.0, 1.2 Hz, 1H, H–C6), 7.35 (dt, *J* = 8.0, 1.2 Hz, 1H, H–C7 of naphthalene), 7.33 (d, *J* = 8.0 Hz, 1H, hydrogen *para* to I), 7.15 (d, *J* = 8.0 Hz, 1H, H–C8 of naphthalene), 7.04 (t, *J* = 8.0 Hz, 1H, hydrogen *meta* to I), 2.13 (s, 3H, Me of naphthalene), 1.91 (s, 3H, Me of phenyl group); ¹³C NMR: δ 144.2, 139.2, 138.7, 136.8, 132.9, 132.1, 131.3, 129.9, 129.1, 128.7, 128.0, 127.7, 126.3, 125.0, 124.6, 101.8, 20.3, 19.5. C₁₈H₁₅I requires: C, 60.35; H, 4.22%; found: C, 60.58; H, 4.29%.

1-(2-Formyl-6-iodophenyl)-2-naphthaldehyde 10

This compound was prepared in two steps as follows. *First step:* A solution of compound **9** (0.147 g, 0.41 mmol), NBS (0.404 g, 2.25 mmol), and benzoylperoxide (0.003 g) in CCl₄ (4 mL) was refluxed overnight under a 150 W lamp. An additional portion of NBS (0.073 g, 0.41 mmol) was then added, and stirring was continued for 4 h. The suspension was filtered when still hot, and the filtrate was concentrated under vacuum. The residue was filtered through silica gel with a 80 : 20 hexanes : Et₂O mixture as eluant to give the product (0.245 g), that was used without further purification in the following step. *Second step:* To a stirred, refluxing solution of the tetrabromide (0.387 g, 0.57 mmol) in dioxane (15 mL), a solution of silver nitrate (0.585 g, 3.44 mmol) in water (3.0 mL) was added dropwise. The resulting mixture was stirred at reflux overnight, and the precipitated silver bromide was then removed by filtration of the hot suspension. The filtrate was concentrated under vacuum, the residue was taken up into THF (7 mL), and a 10% aqueous solution of HCl (5 mL) was then added. The mixture was refluxed for 1 h, cooled down to RT, and extracted with Et₂O (3 × 20 mL). The combined organic phases were dried over NaSO₄, filtered, and concentrated under vacuum. The residue was dissolved in CH₂Cl₂ (20 mL) and anhydrous magnesium sulfate (1 g) was added. The suspension was then refluxed for 2 h under a Dean–Stark apparatus containing molecular sieves. The cooled solution was then concentrated under vacuum and the residue was purified by flash chromatography with a 70 : 30 hexanes : Et₂O mixture as eluant. The product, a pale yellow oil that becomes a waxy solid when digested with pentane, was obtained in 70% yield (0.154 g). IR (thin film): 3050, 1692, 1620, cm⁻¹. ¹H NMR: δ 9.79 (s, 1H, CHO of phenyl group), 9.33 (s, 1H, CHO of naphthalene), 8.30 (dd, *J* = 8.0, 1.2 Hz, 1H, hydrogen *ortho* to I), 8.16 (dd, *J* = 8.0, 1.2 Hz, 1H, hydrogen *para* to I), 8.11 (AB system, *J* = 8.0 Hz, 2H, H–C3 and H–C4 of naphthalene), 8.00 (dd, *J* = 7.9, 1.2 Hz, 1H, H–C5), 7.67 (dt, *J* = 7.9, 1.2 Hz, 1H, H–C6 of naphthalene), 7.49 (dt, *J* = 7.9, 1.2 Hz, 1H, H–C7 of naphthalene), 7.45 (t, *J* = 8.0 Hz, 1H, hydrogen *meta* to I), 7.25 (dd, *J* = 7.9, 1.2 Hz, 1H, H–C8 of naphthalene); ¹³C NMR: δ 190.7, 190.4, 144.5, 143.7, 143.4, 137.5, 136.0, 132.0, 131.8, 130.7, 130.0, 129.4, 128.7, 128.0, 127.9, 126.4, 122.8, 103.5. C₁₈H₁₁I O₂ requires: C, 55.98; H, 2.87%; found: C, 56.18; H, 2.78%.

trans-5-Iodo-9,10-dihydroxy-9,10-dihydrobenzo[*c*]phenanthrene 11

To a solution of compound **10** (0.165 g, 0.43 mmol) in dry THF (7 mL) cooled at –78 °C and kept under a nitrogen atmosphere, a freshly opened 0.1 M solution of samarium diiodide in THF (9 mL) was added by syringe. After 30 min stirring at –78 °C, an additional 9 mL aliquot of samarium diiodide solution was added to the yellow solution, whereupon the color of the

mixture became deep blue. Stirring was continued for 30 min at –78 °C and the reaction was quenched by the addition of a 10% aqueous solution of HCl (5 mL). After addition of Et₂O (20 mL) the organic phase was separated, washed with a saturated aqueous solution of Na₂SO₃ (10 mL), dried over NaSO₄, filtered, and concentrated under vacuum. The residue was purified by flash chromatography first with 40 : 60 and then with 20 : 80 hexanes : Et₂O mixtures as eluants. The product, mp 155–157 °C, was obtained in 75% yield (0.125 g). IR (CH₂Cl₂): 3650, 3055, 2960, 1610, 1425, 1265 cm⁻¹. ¹H NMR: δ 7.84 (dd, *J* = 7.8, 1.2 Hz, 1H, H–C8 of naphthalene), 7.80 (d, *J* = 8.1 Hz, 1H, hydrogen *ortho* to I), 7.74 (br s, 2H, H–C4 and H–C3 of naphthalene), 7.71 (dd, *J* = 7.8, 1.2 Hz, 1H, H–C5 of naphthalene), 7.64 (d, *J* = 8.1 Hz, 1H, hydrogen *para* to I), 7.33–7.25 (m, 2H, H–C6 and H–C7 of naphthalene), 6.95 (t, *J* = 8.1 Hz, 1H, hydrogen *meta* to I), 4.44 (B part of AB system, *J* = 10.6 Hz, 1H, CH–OH bound to naphthalene), 4.23 (A part of AB system, *J* = 10.6 Hz, 1H, CH–OH bound to phenyl); ¹³C NMR: δ 142.2, 139.9, 137.6, 136.3, 133.5, 130.2, 129.2, 129.2, 128.5, 128.4, 128.0, 125.5, 125.3, 124.3, 121.9, 96.8, 74.1, 74.0. C₁₈H₁₃I O₂ requires: C, 55.69; H, 3.37%; found: C, 55.78; H, 3.41%.

Synthesis of diols 1–3, 12 and diketones 4–6

General procedure. The synthesis of diol **1** and diketone **4** is illustrative of the procedure.

trans-5-(4-Methoxyphenyl)-9,10-dihydroxy-9,10-dihydrobenzo[*c*]phenanthrene 1 and 5-(4-methoxyphenyl)benzo[*c*]phenanthrene-9,10-dione 4

To a solution of iodo compound **11** (0.065 g, 0.167 mmol) and 4-methoxyphenyltrimethylstannane (0.204 g, 0.713 mmol) in dry, degassed DMF (5 mL) kept under nitrogen, Pd₂(dba)₃ (0.0345 g, 0.0334 mmol) and AsPh₃ (0.041 g, 0.1336 mmol), both dissolved in degassed DMF (1 mL), were added in this order. The resulting mixture was stirred at reflux for 40 h. Most of the solvent was then evaporated under vacuum and the residue was dissolved in Et₂O (15 mL). The solution was washed with water (4 × 5 mL) to remove the remaining DMF, and the organic phase was dried over NaSO₄, filtered, and concentrated under vacuum. The residue was purified by flash chromatography with pentane : Et₂O mixtures of increasing polarity as eluants (first eluant: 90 : 10; for diketone **4** isolation: 70 : 30; for diol **1** isolation: 30 : 70). Analytically pure samples were obtained by gravity chromatography of the diol and diketone compounds using flash chromatography silica gel as the stationary phase and the same eluants. Diol **1**, a white solid with mp 178–180 °C (decomp.) was obtained in 29% yield (0.0178 g). IR (CH₂Cl₂): 3650, 3054, 2987, 1422, 1264 cm⁻¹. ¹H NMR: δ 7.90–7.83 (m, 2H, H–C1 and H–C2), 7.73 (dd, *J* = 8.5, 1.3 Hz, 1H, H–C8), 7.64 (d, *J* = 8.4 Hz, 1H, H–C3'), 7.56–7.52 (m, 3H, H–C4', H–C6, and H–C7), 7.16 (t, *J* = 8.4 Hz, 1H, H–C3''), 6.93 (dt, *J* = 8.4, 1.3 Hz, 1H, H–C4''), 6.91 (d, *J* = 8.9 Hz, 2H, hydrogen *meta* to methoxy in phenyl ring), 6.51 (d, *J* = 8.9 Hz, 2H, hydrogen *ortho* to methoxy in phenyl ring), 4.83 (B part of AB system, *J* = 10.5 Hz, 1H, H–C10), 4.58 (A part of AB system, *J* = 10.5 Hz, 1H, H–C9), 3.64 (s, 3H, MeO); ¹³C NMR: δ 158.5, 140.5, 139.0, 135.7, 135.0, 133.2, 130.5, 130.0, 129.9, 128.7, 128.6, 128.5, 128.2, 127.3, 126.5, 124.9, 124.8, 121.9, 120.7, 113.5, 74.8, 74.6, 55.1. C₂₅H₂₀O₃ requires: C, 81.50; H, 5.47%; found: C, 81.37; H, 5.38%.

Diketone **4**, a red solid with mp 169–171 °C was obtained in 24% yield (0.0146 g). IR (CH₂Cl₂): 3040, 2964, 1682, 1514, 1252, 1031 cm⁻¹. ¹H NMR: δ 8.13 (dd, *J* = 8.5, 1.3 Hz, 1H, H–C8), 8.05 (B part of an AB system, *J* = 8.4 Hz, 1H, H–C1), 7.83 (dd, *J* = 8.5, 1.3 Hz, 1H, H–C6), 7.81 (B part of an AB system, *J* = 8.4 Hz, 1H, H–C2), 7.68 (dd, *J* = 8.4, 1.2 Hz, 1H, H–C4'), 7.62 (d, *J* = 8.4 Hz, 1H, H–C3'), 7.60 (t, *J* = 8.4 Hz, 1H, H–C7), 7.32 (dt, *J* = 8.4, 1.0 Hz, H–C3''), 7.00 (dt, *J* = 8.4, 1.0 Hz, H–C4''), 6.92 (broad multiplet, 2H, hydrogens *meta* to

MeO in phenyl ring), 6.55 (d, $J = 8.9$ Hz, 2H, hydrogens *ortho* to methoxy in phenyl ring), 3.66 (s, 3H, MeO); ^{13}C NMR: δ 184.0, 159.8, 142.7, 139.2, 138.2, 137.2, 135.0, 133.5, 131.9, 130.2, 130.0, 129.8, 129.4, 128.9, 128.8, 128.3, 127.7, 127.5, 126.1, 122.6, 113.2, 55.2. $\text{C}_{25}\text{H}_{16}\text{O}_3$ requires: C, 82.40; H, 4.43%; found: C, 82.37; H, 4.58%.

trans-5-Phenyl-9,10-dihydroxy-9,10-dihydrobenzo[*c*]phenanthrene 2

A white solid with mp 180–183 °C, was obtained in 12% yield. IR (CH_2Cl_2): 3655, 3055, 2986, 1422, 1264 cm^{-1} . ^1H NMR: δ 7.92–7.82 (m, 2H, H–C1 and H–C2), 7.77 (d, $J = 8.5$ Hz, 1H, H–C8), 7.66 (d, $J = 8.4$ Hz, 1H, H–C3'), 7.60–7.50 (m, 3H, H–C4'', H–C6 and H–C7), 7.15 (t, $J = 8.4$ Hz, 1H, H–C3''), 7.00–6.93 (broad multiplet, 5H of phenyl ring), 6.90 (t, $J = 8.4$ Hz, 1H, H–C4''), 4.87 (B part of AB system, $J = 10.5$ Hz, 1H, H–C10), 4.60 (A part of AB system, $J = 10.5$ Hz, 1H, H–C9); ^{13}C NMR: δ 142.5, 140.5, 139.2, 135.5, 133.0, 130.7, 130.0, 128.9, 128.8, 128.7, 128.6, 128.2, 128.0, 127.9, 127.8, 126.3, 124.9, 124.8, 122.4, 120.8, 74.6, 74.5. $\text{C}_{24}\text{H}_{18}\text{O}_2$ requires: C, 85.18; H, 5.36%; found: C, 85.41; H, 5.29%.

5-Phenylbenzo[*c*]phenanthrene-9,10-dione 6

An orange solid with mp 138–140 °C (decomp.) was obtained in 11% yield. IR (CH_2Cl_2): 3050, 2960, 1682, 1510, 1250, 1025 cm^{-1} . ^1H NMR: δ 8.15 (dd, $J = 8.0, 1.4$ Hz, 1H, H–C8), 8.07 (B part of an AB system, $J = 8.5$ Hz, 1H, H–C1), 7.86 (dd, $J = 8.0, 1.4$ Hz, 1H, H–C6), 7.81 (A part of an AB system, $J = 8.5$ Hz, 1H, H–C2), 7.68 (dd, $J = 8.4, 1.0$ Hz, 1H, H–C4'), 7.63 (t, $J = 8.0$ Hz, 1H, H–C7), 7.62 (d, $J = 8.0$ Hz, 1H, H–C3'), 7.29 (dt, $J = 8.0, 1.0$ Hz, 1H, H–C3''), 7.07–7.00 (broad multiplet, 5H of phenyl ring), 6.97 (dt, $J = 8.0, 1.0$ Hz, 1H, H–C4''); ^{13}C NMR: δ 184.0, 143.5, 140.0, 139.2, 138.4, 137.5, 135.9, 132.5, 129.9, 129.8, 129.4, 129.1, 129.0, 128.9, 128.7, 128.4, 127.7, 127.4, 127.2, 126.2, 122.6. $\text{C}_{24}\text{H}_{14}\text{O}_2$ requires: C, 86.21; H, 4.22%; found: C, 85.99; H, 4.39%.

trans-5-(4-Nitrophenyl)-9,10-dihydroxy-9,10-dihydrobenzo[*c*]phenanthrene 3

A red solid with mp >200 °C was obtained in 11% yield. IR (CH_2Cl_2): 3650, 3053, 2986, 1423, 1266, 1045 cm^{-1} . ^1H NMR: δ 7.93–7.89 (m, 2H, H–C1 and H–C2), 7.88 (d, $J = 8.5$ Hz, H–C8), 7.83 (d, $J = 8.8$ Hz, 2H, hydrogens *ortho* to NO_2 in phenyl ring), 7.67 (d, $J = 8.1$ Hz, H–C3'), 7.60 (t, $J = 8.5$ Hz, 1H, H–C7), 7.53 (d, 8.5 Hz, 1H, H–C6), 7.45 (d, $J = 8.1$ Hz, 1H, H–C4'), 7.17–7.12 (m, 3H, H–C3'' and hydrogens *meta* to NO_2 in phenyl ring), 6.92 (t, $J = 8.1$ Hz, 1H, H–C4''), 4.86 (B part of AB system, $J = 10.5$ Hz, 1H, H–C10), 4.62 (A part of AB system, $J = 10.5$ Hz, 1H, H–C9); ^{13}C NMR: δ 149.5, 146.2, 140.5, 140.0, 136.2, 133.4, 130.5, 130.3, 129.5, 129.3, 128.5, 128.1, 128.0, 125.7, 125.3, 125.2, 124.2, 124.1, 123.2, 121.1, 74.6, 74.5. $\text{C}_{24}\text{H}_{17}\text{NO}_4$ requires: C, 75.20; H, 4.47; N, 3.65%; found: C, 75.49; H, 4.56; N, 3.59%.

5-(4-Methylphenyl)benzo[*c*]phenanthrene-9,10-dione 5

An orange solid with mp 115–117 °C was obtained in 15% yield. IR (CH_2Cl_2): 3055, 2988, 1684, 1488, 1266 cm^{-1} . ^1H NMR: δ 8.12 (dd, $J = 8.5, 1.3$ Hz, 1H, H–C8), 8.05 (B part of an AB system, $J = 8.4$ Hz, 1H, H–C1), 7.82 (dd, $J = 8.5, 1.3$ Hz, 1H, H–C6), 7.80 (A part of an AB system, $J = 8.5$ Hz, 1H, H–C2), 7.68 (d, $J = 8.4$ Hz, 1H, H–C4'), 7.63 (d, $J = 8.4$ Hz, 1H, H–C3'), 7.60 (t, $J = 8.4$ Hz, 1H, H–C7), 7.28 (dt, $J = 8.4, 1.0$ Hz, H–C3''), 6.97 (dt, $J = 8.4, 1.0$ Hz, H–C4''), 6.88 (broad multiplet, 2H, hydrogens *meta* to Me in phenyl ring), 6.80 (A part of an AB system, $J = 8.8$ Hz, 2H, hydrogens *ortho* to Me in phenyl ring), 2.13 (s, 3H, Me in phenyl ring); ^{13}C NMR: δ 184.0, 143.0, 139.2, 138.5, 138.3, 137.5, 137.1, 135.2, 131.9, 129.9, 129.8, 129.4, 129.1, 129.0, 128.9, 128.8, 128.5, 127.7, 127.5, 126.1,

122.6, 22.2. $\text{C}_{25}\text{H}_{16}\text{O}_2$ requires: C, 86.19; H, 4.63%; found: C, 86.28; H, 4.66%.

5-(4-Fluorophenyl)benzo[*c*]phenanthrene-9,10-dione 7

The inseparable mixture of crude diketone 7 and diol 12, obtained as described above, was subjected to the following procedure: To a stirred mixture of 7 and 12 (0.016 g, approximately 0.045 mmol) in dry DMSO (0.5 mL) kept at RT under nitrogen, triethylamine (0.022 mL, 0.16 mmol) was added, followed by $\text{Py}\cdot\text{SO}_3$ (0.0178 g, 0.112 mmol) in 0.5 mL of dry DMSO. After 5 h stirring at RT, the reaction was quenched by the addition of water (5 mL), and the solution was extracted with Et_2O (5×5 mL). The diethyl ether solution was washed with water (5 mL) to remove the remaining DMSO, and the organic phase was dried over NaSO_4 , filtered, and concentrated under vacuum. The residue was purified by flash chromatography with pentane : Et_2O mixtures of increasing polarity as eluants (first eluant: 90 : 10; for diketone isolation: 80 : 20). Compound 7, a red solid with mp 93–95 °C was obtained in 12% overall yield from 11. IR (CH_2Cl_2): 3050, 2986, 1683, 1485, 1266 cm^{-1} . ^1H NMR: δ 8.15 (dd, $J = 7.8, 1.3$ Hz, 1H, H–C8), 8.07 (B part of an AB system, $J = 8.5$ Hz, 1H, H–C1), 7.83 (A part of an AB system, $J = 8.5$ Hz, 1H, H–C2), 7.81 (dd, $J = 7.8, 1.3$ Hz, 1H, H–C6), 7.68–7.64 (m, 2H, H–C3' and H–C4'), 7.63 (t, $J = 7.8$ Hz, 1H, H–C7), 7.34 (t, $J = 8.9$ Hz, H–C3''), 7.01 (t, $J = 8.9$ Hz, H–C4''), 6.97 (broad multiplet, 2H, hydrogen *meta* to F in phenyl ring), 6.72 (t, $J = 8.0$ Hz, 2H, hydrogens *ortho* to F in phenyl ring); ^{13}C NMR: δ 184.0, 142.3, 142.2, 139.2, 138.1, 137.8, 137.5, 135.7, 132.1, 130.8, 130.1, 130.0, 129.5, 129.4, 129.0, 128.8, 127.9, 127.3, 126.3, 122.7, 115.4; ^{19}F NMR: δ 117.0. $\text{C}_{24}\text{H}_{13}\text{FO}_2$ requires: C, 81.81; H, 3.72%; found: C, 81.59; H, 3.86%.

2D EXSY Experiments

For all the 2D EXSY experiments, the samples were dissolved in $\text{THF}-d_8$ and were degassed by several freeze–thaw cycles on a high vacuum line to remove dissolved oxygen. The spectra were recorded at ≤ 210 K temperature. Pure absorption 2D EXSY spectra were recorded employing the following parameters: spectral width 300 Hz; 256 time increments of 64 transients each; mixing times of 0.8 and 1 s for compounds 1–3 and 4–7, respectively; relaxation delays of 2.0 s (*ca.* 3 T_1 values of the relevant protons at the EXSY temperature).

Acknowledgements

National Science Foundation (CHE-9904275) and CNR-Progetto Strategico provided support for this work.

References and Notes

- (a) C. A. Hunter, *Angew. Chem. Int. Ed. Engl.*, 1993, **32**, 1584–1586; (b) C. A. Hunter, *Chem. Soc. Rev.*, 1994, 101–109.
- (a) F. Diederich, *Angew. Chem. Int. Ed. Engl.*, 1988, **27**, 362–386; (b) D. B. Smithrud, T. B. Wyman and F. Diederich, *J. Am. Chem. Soc.*, 1991, **113**, 5420–5426 and references therein.; (c) M. M. Conn, G. Deslongchamps, J. de Mendoza and J. Rebek, *J. Am. Chem. Soc.*, 1993, **115**, 3458–3557; (d) D. Philip and J. F. Stoddard, *Angew. Chem. Int. Ed. Engl.*, 1996, **35**, 1154–1196; (e) A. V. Muehldorf, D. Van Egen, J. C. Warner and A. D. Hamilton, *J. Am. Chem. Soc.*, 1988, **110**, 6561–6562; (f) S. C. Zimmermann, Z. Zeng, W. Wu and D. E. Reichert, *J. Am. Chem. Soc.*, 1991, **113**, 183–196; (g) L. T. Newcomb and S. H. Gellman, *J. Am. Chem. Soc.*, 1994, **116**, 4993–4994; (h) J. E. Cochran, T. J. Parrott, B. J. Whitlock and H. W. Whitlock, *J. Am. Chem. Soc.*, 1992, **114**, 2269–2270; (i) E. C. Breinlinger and V. M. Rotello, *J. Am. Chem. Soc.*, 1997, **119**, 1165–1166.
- Review: G. B. Jones, *Tetrahedron*, 2001, **57**, 7999–8016.
- (a) W. Saenger, *Principles of Nucleic Acid Structures*, Springer Verlag, New York, 1984; (b) S. K. Burley and G. A. Petsko, *Science*, 1985, **229**, 23–28; (c) C. A. Hunter, J. Singh and J. M. Thornton,

- J. Mol. Biol.*, 1991, **218**, 837–846; (d) C. A. Hunter, *J. Mol. Biol.*, 1993, **230**, 1025–1054; (e) O. F. Schall and G. W. Gokel, *J. Org. Chem.*, 1996, **61**, 1149–1458; (f) K. M. Guckian, B. A. Schweitzer, R. X. F. Ren, C. J. Sheils, P. L. Paris, D. C. Tanmasseri and E. T. Kool, *J. Am. Chem. Soc.*, 1996, **118**, 8182–8183; (g) T. L. Ho, P. Y. Liao and K. T. Wang, *J. Chem. Soc., Chem. Commun.*, 1995, 2437–2438; (h) D. Ranganathan, V. Haridas, R. Gilardi and J. L. Karle, *J. Am. Chem. Soc.*, 1998, **120**, 10793–10800; (i) F. Chelli, F. L. Gervasio, P. Procacci and V. Schettino, *J. Am. Chem. Soc.*, 2002, **124**, 6133–6143 and references therein.; (j) F. Chelli, F. L. Gervasio, P. Procacci and V. Schettino, *Proteins*, 2002, **48**, 117–125.
- 5 (a) C. A. Hunter, M. N. Meah and J.K.M. Sanders, *J. Am. Chem. Soc.*, 1990, **112**, 5773–5780; (b) A. S. Shetty, J. Zhang and J. S. Moore, *J. Am. Chem. Soc.*, 1996, **118**, 1019–1027 and references therein.; (c) G. W. Coates, A. R. Dunn, L. M. Henling, J. W. Ziller, E. B. Lobkovsky and R. H. Grubbs, *J. Am. Chem. Soc.*, 1998, **120**, 3641–3649.
- 6 (a) T. Dahl, *Acta Chem. Scand.*, 1994, **48**, 95–106; (b) J. H. Williams, *Acc. Chem. Res.*, 1993, **26**, 593–598; (c) H. L. Anderson, A. Bashall, K. Henrick, M. McPartlin and J. K. M. Sanders, *Angew. Chem., Int. Ed. Engl.*, 1994, **33**, 429–431; (d) I. Dance and M. Scudder, *Chem. Eur. J.*, 1996, **2**, 481–486; (e) Z. H. Wang, T. Hirose, K. Hiratani, Y. Yang and K. Kasuga, *Chem. Lett.*, 1996, 603–604; (f) C. B. Martin, B. O. Patrick and A. Cammers-Goodwin, *J. Org. Chem.*, 1999, **64**, 7568–7578; (g) Y. Nakamura, H. Suzuki, Y. Hayashida, T. Kudo and J. Nishimura, *Liebigs Ann./Recueil*, 1997, 1769–1776.
- 7 (a) W. L. Jorgensen and D. L. Severance, *J. Am. Chem. Soc.*, 1990, **112**, 4768–4774; (b) C. A. Hunter and J. K. M. Sanders, *J. Am. Chem. Soc.*, 1990, **112**, 5525–5534; (c) P. Linse, *J. Am. Chem. Soc.*, 1992, **114**, 4366–4373; (d) P. Hobza, H. L. Selzle and E. W. Schlag, *J. Am. Chem. Soc.*, 1994, **116**, 3500–3506; (e) V. E. Williams, R. P. Lemieux and G. R. Thatcher, *J. Org. Chem.*, 1996, **61**, 1927–1933; (f) C. Chipot, R. Jaffe, B. Maigret, D. A. Pearlman and P. A. Kollman, *J. Am. Chem. Soc.*, 1996, **118**, 11217–11224; (g) J. M. Seminario, A. G. Zacarias and J. M. Tour, *J. Am. Chem. Soc.*, 1998, **120**, 3970–3974; (h) S. Tsuzuki, K. Honda, T. Uchimaru, M. Mikami and K. Tanabe, *J. Am. Chem. Soc.*, 2002, **124**, 104–112; (i) M. Claessens, L. Palombini, M. L. Stein and J. Reisse, *Nouv. J. Chem.*, 1982, **6**, 595; (j) M. O. Sinnokrot, E. F. Valeev, C. D. Sherrill, *J. Am. Chem. Soc.*, 2002, **124**, in press.
- 8 The preference for the edge-to-face orientation has been demonstrated for gaseous, liquid, and crystalline benzene: (a) X. Shi and L. S. Bartell, *J. Chem. Phys.*, 1988, **92**, 5667–5673; (b) L. J. Lowden and D. J. Chandler, *J. Chem. Phys.*, 1974, **61**, 5228–5241; (c) D. Hall and D. E. Williams, *Acta Crystallogr., Sect. A: Cryst. Phys., Diffraction, Theor. Gen. Cryst.*, 1975, **31**, 56–58 and references therein.
- 9 (a) F. Cozzi, M. Cinquini, R. Annunziata, T. Dwyer and J. S. Siegel, *J. Am. Chem. Soc.*, 1992, **114**, 5729–5733; (b) F. Cozzi, M. Cinquini, R. Annunziata and J. S. Siegel, *J. Am. Chem. Soc.*, 1993, **115**, 5330–5331; (c) F. Cozzi, F. Ponzini, R. Annunziata, M. Cinquini and J. S. Siegel, *Angew. Chem., Int. Ed. Engl.*, 1995, **34**, 1019–1020; (d) F. Cozzi and J. S. Siegel, *Pure Appl. Chem.*, 1995, **67**, 683–689.
- 10 (a) J. A. Zoltewicz, N. A. Maier and W. M. F. Fablan, *J. Org. Chem.*, 1998, **63**, 4985–4990 and references therein; (b) N. Kaneta, F. Mitamura, M. Uemura, Y. Murata and K. Komatsu, *Tetrahedron Lett.*, 1996, **37**, 5835–5838.
- 11 (a) C. R. Patrick and G. S. Prosser, *Nature*, 1960, **187**, 1021; (b) J. H. Williams, J. K. Cockroft and A. N. Fitch, *Angew. Chem., Int. Ed. Engl.*, 1992, **31**, 1655–1657; (c) R. E. Gillard, J. F. Stoddardt, A. J. P. White, B. J. Williams and D. J. Williams, *J. Org. Chem.*, 1996, **61**, 4504–4505; (d) G. W. Coates, A. B. Dunn, L. M. Henling, D. A. Dougherty and R. H. Grubbs, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 248–251; (e) F. Ponzini, R. Zaghera, K. Hardcastle and J. S. Siegel, *Angew. Chem., Int. Ed.*, 2000, **39**, 2323–2325; (f) M. T. Messina, P. Metrangolo, S. Pappalardo, M.F. Parisi, T. Pilati and G. Resnati, *Chem. Eur. J.*, 2000, **6**, 3495–3500; (g) C. Y. Kim, P. P. Chandra, A. Jain and D. W. Christianson, *J. Am. Chem. Soc.*, 2001, **123**, 9620–9627 and references therein.
- 12 (a) F. J. Carver, C. A. Hunter and E. M. Seward, *J. Chem. Soc., Chem. Commun.*, 1998, 775–776; (b) F. J. Carver, C. A. Hunter, D. J. Livingstone, J. F. McCabe and E. M. Seward, *Chem. Eur. J.*, 2002, **8**, 2847–2859 and references therein; (c) E. Kim, S. Paliwal and C. S. Wilcox, *J. Am. Chem. Soc.*, 1998, **120**, 11192–11193 and references therein; (d) K. Nakamura and K. N. Houk, *Org. Lett.*, 1999, **1**, 2049–2051; (e) Review: W. B. Jennings, B. M. Farrell and J. F. Malone, *Acc. Chem. Res.*, 2001, **34**, 885–894.
- 13 For recent examples of molecular recognition between arenes based on an edge-to-face interaction, see: (a) A. P. Poisson and C. A. Hunter, *J. Chem. Soc., Chem. Commun.*, 1996, 1723–1724; (b) M.J. Cloninger and H.W. Whitlock, *J. Org. Chem.*, 1998, **63**, 6153–6159; (c) S. Superchi, M.I. Donnoli, G. Proni, G.P. Spada and C. Rosini, *J. Org. Chem.*, 1999, **64**, 4762–4767. For the influence exerted by the edge-to-face interactions on the relative orientation of different arenes, see; (d) A. Garcia-Martinez, J. Osio Barcina, A. de Fresno Cerezo and R. Gutierrez Rivas, *J. Am. Chem. Soc.*, 1998, **120**, 673–679; (e) B. Uno, T. Iwamoto and N. Okumura, *J. Org. Chem.*, 1998, **63**, 9794–9800; (f) S. V. Lindeman, D. Kosynkin and J. K. Kochi, *J. Am. Chem. Soc.*, 1998, **120**, 13268–13269; (g) For a review on the CH– π interactions, see: M. Nishio, Y. Umezawa, M. Hirota and Y. Takeuchi, *Tetrahedron*, 1995, **51**, 8665–8701.
- 14 See for instance: P.R. Ashton, D. Joachimi, N. Spencer, J. F. Stoddardt, C. Tschierske, A. J. P. White, D. J. Williams and K. Zab, *Angew. Chem., Int. Ed. Engl.*, 1994, **33**, 1503–1506.
- 15 C. B. Martin, H. R. Mulla, P. G. Willis and A. Cammers-Goodwin, *J. Org. Chem.*, 1999, **64**, 7802–7806.
- 16 M. J. Rashkin and M. L. Waters, *J. Am. Chem. Soc.*, 2002, **124**, 1860–1861.
- 17 1-Bromo-2-methylnaphthalene can be used in excess because it produces little of the homocoupling product under these conditions.
- 18 J. E. McMurry, *Chem. Rev.*, 1989, **89**, 1513–1524.
- 19 R. G. R. Bacon and W. S. Lindsay, *J. Chem. Soc.*, 1958, 1375–1381.
- 20 E. Abu-Shagra and J. Blum, *J. Heterocycl. Chem.*, 1991, **28**, 473–475.
- 21 G.A. Molander and C.R. Harris, *Chem. Rev.*, 1996, **96**, 307–338.
- 22 For similar results see: (a) K. Ohmori, M. Kitamura and K. Suzuki, *Angew. Chem., Int. Ed.*, 1999, **38**, 1226–1229; (b) N. Taniguchi, T. Hata and M. Uemura, *Angew. Chem., Int. Ed.*, 1999, **38**, 1232–1235; (c) R. Annunziata, M. Benaglia, M. Cinquini and L. Raimondi, *Eur. J. Org. Chem.*, 1999, 3369–3374.
- 23 M. Benaglia, S. Toyota, C. R. Woods and J. S. Siegel, *Tetrahedron Lett.*, 1997, **38**, 4737–4740.
- 24 (a) J. K. Stille, *Angew. Chem., Int. Ed. Engl.*, 1986, **25**, 508–524; (b) V. Farina, *Pure Appl. Chem.*, 1996, **68**, 73–78 and references therein.
- 25 A modification of the published procedure was used: R. G. Harvey, S. H. Goh and C. Cortez, *J. Am. Chem. Soc.*, 1975, **97**, 3468–3479.
- 26 For the rotation around the aryl–aryl bond in 1-phenylbenzo[*c*]phenanthrene derivatives a ΔG^\ddagger of ca. 54 kJ mol⁻¹ was determined; the enantiomerization of the benzo[*c*]phenanthrene skeleton requires a ΔG^\ddagger of ca. 67 kJ mol⁻¹: W. H. Laarhoven, W. H. M. Peters and A. H. A. Tinnemans, *Tetrahedron*, 1978, **34**, 769–777. In molecules related to diols 1–3 this process requires even higher energy (see ref. 22a).
- 27 For a review on the use of 2D EXSY NMR spectroscopy to measure chemical dynamics, see: C. L. Perrin and T. J. Dwyer, *Chem. Rev.*, 1990, **90**, 935–967.
- 28 Because of the logarithmic relationship between *k* and ΔG^\ddagger , the major contribution to the error in determining ΔG^\ddagger is the error in the temperature. All spectra were measured with the same instrumentation at similar temperatures. A liberal estimate of the error is 1 kJ mol⁻¹.
- 29 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, V. G. Zakrzewski, J. A. J. Montgomery, R. E. Stratmann, J. C. Burant, S. Dapprich, J. M. Millam, A. D. Daniels, K. N. Kudin, M. C. Strain, O. Farkas, J. Tomasi, V. Barone, M. Cossi, R. Cammi, B. Mennucci, C. Pomelli, C. Adamo, S. Clifford, J. Ochterski, G. A. Petersson, P. Y. Ayala, Q. Cui, K. Morokuma, D. K. Malick, A. D. Rabuck, K. Raghavachari, J. B. Foresman, J. Cioslowski, J. V. Ortiz, B. B. Stefanov, G. Liu, A. Liashenko, P. Piskorz, I. Komaromi, R. Gomperts, R. L. Martin, D. J. Fox, T. Keith, M. A. Al-Laham, C. Y. Peng, A. Nanayakkara, C. Gonzalez, M. Challacombe, P. M. W. Gill, B. Johnson, W. Chen, M. W. Wong, J. L. Andres, C. Gonzalez, M. Head-Gordon, E. S. Replogle, J. A. Pople, *Gaussian 98*, Revision A.6, Pittsburgh PA, 1998.
- 30 A. D. Becke, *J. Chem. Phys.*, 1993, **98**, 5648–5652. Structural computations were performed using the hybrid density functional method, B3LYP; Becke's 3 parameter functional plus the nonlocal correlation of Lee-Yang-Parr with both local and nonlocal terms. Dunning's DZV(2d,p) double- ζ valence basis set was employed. Single point energy computations were performed on optimized structures using MP2 dynamic correlation treatment for superior energetic analysis.
- 31 (a) C. Lee, W. Yang and R. G. Parr, *Phys. Rev. B*, 1988, **37**, 785–792; (b) B. Miehlich, A. Savin, H. Stoll and H. Preuss, *Chem. Phys. Lett.*, 1989, **157**, 200–207.
- 32 T. H. Dunning, Jr., P. J. Hay, *Modern Theoretical Chemistry*; Plenum, New York, 1976, pp. 1–28.
- 33 J. M. L. Martin, *Chem. Phys. Lett.*, 1996, **262**, 97–104.
- 34 C. Moller and M. S. Plesset, *Phys. Rev.*, 1934, **46**, 618–622.