

Molybdenum-catalyzed hydrostannations of allenylcarbinols†

Uli Kazmaier* and Manuela Klein

Received (in Cambridge, UK) 24th August 2004, Accepted 21st October 2004

First published as an Advance Article on the web 2nd December 2004


DOI: 10.1039/b413025a

Allenylcarbinols undergo regioselective hydrostannation in the presence of MoI_3 , a catalyst which was developed for the hydrostannation of propargyl alcohols and derivatives; allylstannanes are formed preferentially, which can easily be converted into allyl iodides.


During the last few years functionalized allenes have become more and more interesting not only as targets in natural product synthesis, but also as valuable synthetic precursors for the synthesis of complex molecules.¹ In this context, carbon–carbon and carbon–heteroatom bond formations are of major interest, triggering the development of preparatively useful methods such as additions² or palladium-catalyzed transformations.³ Besides cyclizations⁴ especially transition metal-catalyzed hydrometallations such as hydroborations,⁵ hydrozirconations⁶ or hydrostannations⁷ are synthetically useful transformations. Vinyl or allyl organometallics are obtained depending on the regioselectivity of the hydrometallation step. In principle (Scheme 1), four different products (A–D) are possible, and the product distribution depends on the reaction conditions used.

While in Lewis acid-catalyzed reactions of terminal allenes attack occurs at the internal double bond providing vinylstannanes **B**,⁸ in the presence of most palladium complexes the terminal double bond reacts preferentially giving rise to allylstannanes **C** as an *E/Z* mixture.⁹ Interestingly, the regioselectivity for the hydrostannation of the terminal double bond can be converted towards vinylstannanes **D** by switching from $\text{Pd}(\text{PPh}_3)_4$ to $\text{Pd}(\text{OH})_2/\text{C}$.¹⁰ Hydrostannations of allenes bearing suitable functional groups such as aryl halides¹¹ or further double bonds¹² allow subsequent cyclizations.

Recently, we developed a new molybdenum catalyst for hydrostannations of alkynes. MoI_3 ($\text{Mo}(\text{CO})_3(\text{CNtBu})_3$) was found to be an excellent catalyst for regioselective hydrostannations towards the internal vinylstannanes, being superior to the generally used palladium catalysts (Scheme 2).¹³



Scheme 1 Hydrostannation of allenes.




Scheme 2 Molybdenum-catalyzed hydrostannation of alkynes.

Herein, we now report on our investigations of MoI_3 -catalyzed hydrostannations of allenes. We chose allenylcarbinols **2** as substrates which could easily be obtained from the corresponding propargyl alcohols **1** according to Burgess and Jennings¹⁴ (Scheme 3).

Those allenylcarbinols were subjected to the reaction conditions optimized for the hydrostannation of alkynes (THF, Δ) and the process was monitored by TLC. In general, the reactions were completed after 4 h, and therefore allenylcarbinols **2** are significantly more reactive than the corresponding propargyl derivatives **1**. The product ratio was determined by NMR. Overall, four different products were obtained, but products of type **B** and **D** were never observed. The results obtained are summarised in Table 1.

First we investigated the hydrostannation of the phenylethyl-substituted allenylcarbinol **2a**. Addition occurred exclusively at the terminal double bond and only to give the allylstannanes **3a**, which were obtained as an *E/Z* mixture in the ratio 1 : 2. To determine if



Scheme 3 Preparation of allenylcarbinols.

Table 1 Hydrostannation of allenylcarbinols **2**

Allene	R	Temp [°C]	Yield [%]	Ratio (<i>E</i>)-3 : (<i>Z</i>)-3 : 4 : 5
2a	PhCH_2CH_2	55	68	1 : 2 : 0 : 0
2a	PhCH_2CH_2	rt	74	1 : 2 : 0 : 0
2b	$2\text{-NO}_2\text{Ph}$	55	76	12 : 12 : 1 : 0
2c	4-ClPh	55	68	1 : 1 : 0 : 0
2d	$2,6\text{-Cl}_2\text{Ph}$	55	74	1 : 2 : 0 : 0
2e	$2,4,6\text{-(MeO)}_3\text{Ph}$	55	86	1 : 0 : 1 : 2
2e	$2,4,6\text{-(MeO)}_3\text{Ph}$	rt	23 ^a	10 : 0 : 6 : 1

^a Only 1% MoI_3 , 2 equiv. Bu_3SnH used.

† Dedicated to Prof. Dr M. Veith on the occasion of his 60th birthday.

*u.kazmaier@mx.uni-saarland.de

this ratio depends on the reaction conditions, *e.g.* on the temperature, we carried out the reaction also at room temperature. Under these milder conditions the yield was higher, but no influence on the product ratio was observed.


To figure out if electronic factors in the substituent R might have an influence on the product formation, we investigated several aromatic allenylcarbinols bearing electron-withdrawing and -donating groups. In the hydrostannylation of *o*-nitrophenyl derivative **2b** the allylstannanes **3b** were by far the major products, the regioisomer **4b** was formed in trace amounts. Similar results were obtained in the reaction of *p*-chlorophenyl- and 2,6-dichlorophenyl-substituted substrates (**2c**, **2d**), but the situation changed dramatically with electron-rich aromatic systems such as the 2,4,6-trimethoxyphenyl derivative **2e**. In this case the elimination product **5e** was obtained preferentially, besides allylstannane (*E*)-**3e** and vinylstannane **4e**. Interestingly no (*Z*)-configured product was obtained in this case, and we assume that this isomer undergoes elimination according to Fig. 1.

Obviously the strong electron-donating groups facilitate the cleavage of the OH group, probably *via* stabilization of the carbenium ion intermediate. This would explain the high ratio of elimination product obtained under the standard conditions. To figure out if this proposal is reasonable, we investigated the reaction also under milder conditions. And indeed, the amount of elimination product could be reduced significantly.


The high selectivities towards the allyl and vinyl stannanes **3** and **4** can be explained by the following mechanistic rationale (Scheme 4). Probably in the first step some of the isonitrile ligands dissociate from the molybdenum opening free coordination sites for the oxidative addition of the tin hydride and coordination of the allene. Depending on the substrate used, only the terminal or both double bonds coordinate to the molybdenum giving rise to intermediates **A** and **B**. Subsequent hydrometallation should provide intermediates **A'** and **B'** with the metal fragment added to the sterically least hindered position. The products **3** and **4** were formed from these intermediates *via* reductive elimination. The *E/Z* isomers were formed by coordination to the two diastereotopic faces of the terminal double bond.

Probably this mechanism is slightly different from the one proposed for the hydrostannylation of alkynes which seems to start with a transfer of the stannyl group and a subsequent hydrogen transfer in the reductive elimination step. But both possible mechanisms were also discussed for palladium-catalyzed hydrostannations. Therefore, it is reasonable that the situation with the molybdenum complexes is similar.

To improve the synthetic potential of this protocol we subjected the mixture of (*E/Z*)-**3b** to a metal-halogen exchange (Scheme 5). Although the isomeric mixture was used, the corresponding allyl iodide (*E*)-**6** was obtained as a single isomer in high yield. Obviously under these conditions only the thermodynamically



Scheme 4 Proposed mechanism for the hydrostannylation of allenes.



Scheme 5 Metal-halogen exchange of allylstannane **3**.

most stable product is formed, which is an interesting substrate for further synthetic applications.

In conclusion we have shown that molybdenum-catalyzed hydrostannylation of allenes proceeds regioselectively towards the sterically least hindered stannylated product. Further research concerning the selectivity and the reaction mechanism is currently in progress.

Financial support by the Deutsche Forschungsgemeinschaft as well as the Fonds der Chemischen Industrie is gratefully acknowledged.

Uli Kazmaier* and Manuela Klein

Universität des Saarlandes, Institut für Organische Chemie, D-66123, Saarbrücken, Germany. E-mail: u.kazmaier@mx.uni-saarland.de; Fax: 49 681 3022409; Tel: 49 681 3023409

Notes and references

- (a) *Allenes in Organic Synthesis*, ed. H. F. Schuster, G. M. Copolla, Wiley, New York, 1984; (b) *Modern Allene Chemistry*, ed. N. Krause, A. S. K. Hashmi, Wiley-VCH, Weinheim, 2004.
- Reviews: (a) R. Zimmer, *Synthesis*, 1993, 165; (b) T. G. Back, *Tetrahedron*, 2001, **57**, 5263.
- Review: R. Zimmer, C. U. Dinesh, E. Nandan and F. A. Khan, *Chem. Rev.*, 2000, **100**, 3067.
- Reviews: (a) R. W. Bates and V. Satchavoen, *Chem. Soc. Rev.*, 2002, **31**, 12; (b) N. Krause, A. Hoffman-Röder and J. Canisius, *Synthesis*, 2002, 1759.
- Y. Yamamoto, R. Fijikawa, A. Yamada and N. Miyaura, *Chem. Lett.*, 1999, 1069.




Fig. 1 Elimination of stannylated allyl alcohols.

- 6 (a) H. Maeta, T. Hasegawa and K. Suzuki, *Synlett*, 1993, 341; (b) M. Chino, T. Matsunoto and K. Suzuki, *Synlett*, 1994, 359; (c) J. H. Pi and X. Huang, *Synlett*, 2003, 2413.
- 7 Review: N. D. Smith, J. Mancuso and M. Lautens, *Chem. Rev.*, 2000, **100**, 3257.
- 8 V. Gevorgyan, J. X. Liu and Y. Yamamoto, *Chem. Commun.*, 1998, 37.
- 9 (a) Y. Ichinose, K. Oshima and K. Utimoto, *Bull. Chem. Soc. Jpn.*, 1988, **61**, 2693; (b) K. Koerber, J. Goré and J. M. Valele, *Tetrahedron Lett.*, 1991, **32**, 1187; (c) T. N. Mitchell and U. Schneider, *J. Organomet. Chem.*, 1991, **405**, 195; (d) V. Gevorgyan, J. X. Liu and Y. Yamamoto, *J. Org. Chem.*, 1997, **62**, 2963.
- 10 M. Lautens, D. Ostrovsky and B. Tao, *Tetrahedron Lett.*, 1997, **38**, 6343.
- 11 R. Grigg and J. M. Sansano, *Tetrahedron*, 1996, **52**, 13441.
- 12 (a) J. Marco-Contelles, G. Balme, D. Bouyssi, C. Destabel, C. D. Henriët-Bernard, J. Grimaldi and J. M. Halem, *J. Org. Chem.*, 1997, **62**, 1202; (b) S. K. Kung, T. B. Baik, A. N. Kulak, Y. H. Ha, Y. Lim and J. Park, *J. Am. Chem. Soc.*, 2000, **122**, 11529.
- 13 (a) U. Kazmaier, D. Schauß and M. Pohlman, *Org. Lett.*, 1999, **1**, 1017; (b) U. Kazmaier, D. Schauß, M. Pohlman and S. Raddatz, *Synthesis*, 2000, 914; (c) U. Kazmaier, M. Pohlman and D. Schauß, *Eur. J. Org. Chem.*, 2000, 2761; (d) U. Kazmaier, D. Schauß, S. Raddatz and M. Pohlman, *Chem. Eur. J.*, 2001, **7**, 456; (e) U. Kazmaier and S. Braune, *J. Organomet. Chem.*, 2002, **641**, 26.
- 14 K. Burgess and J. D. Jennings, *J. Am. Chem. Soc.*, 1991, **113**, 6129.