### Natural Product Synthesis

## Three Rings in One Step: A Quick Approach to IKD-8344\*\*

Yang Zou and Yikang Wu\*

IKD-8344 (**1**, Scheme 1) was first isolated from an unidentified alkalophilic *Actinomycete* (strain no. 8344) in the early 1990s by Nishii and co-workers,<sup>[1]</sup> and later from *Streptomyces* 



**Scheme 1.** The structure of IKD-8344 with the relative configuration determined by single crystal X-ray crystallography and the absolute configuration depicted arbitrarily (Ref. [1b]).

sp. A6792 by Kim and co-workers.<sup>[2]</sup> This natural dilactone, which contains a 28-membered ring, represents a worthy target for synthetic studies, because of its intriguing molecular architecture and the potent activities against mouse leukemia ( $IC_{50}$  0.54 ng mL<sup>-1</sup>) and *Trichinella spiralis*.<sup>[3]</sup>

The most striking structural motifs in IKD-8344 are the 2,5-disubstituted THF rings with one of the two substituting carbon atoms being part of a hidden aldol unit. These structural motifs are similar to those found in nonactin and pamamycins. In previous studies<sup>[4,5]</sup> on the synthesis of similar THF-containing targets, we developed a novel route to enantiomerically pure THF rings by using an  $S_N^2$  type intramolecular O alkylation at a sterically hindered carbon atom in  $\beta$  position to a carbonyl group. Such an  $S_N^2$  reaction would normally be considered hopeless, because it competes with the normally very facile  $\alpha$  racemization and  $\beta$  elimination. In the beginning, en route to a synthesis of nonactin, we realized the formation of one ring at a time. Later, in the synthesis of pamamycin 621A, we achieved the formation of two rings in one step. Encouraged by these results, we decided

 [\*] Dr. Y. Zou, Prof. Dr. Y. Wu State Key Laboratory of Bioorganic and Natural Products Chemistry, Shanghai Institute of Organic Chemistry Chinese Academy of Sciences 345 Lingling Road, Shanghai 200032 (China) E-mail: yikangwu@sioc.ac.cn to take up an even greater challenge with the concurrent formation of three THF rings from a highly sensitive linear precursor. Such an approach would also circumvent the undesired complications<sup>[3]</sup> encountered in the assembly of the preformed single THF ring containing fragments in the previous total synthesis.

The outline of our retrosynthetic analysis is shown in Scheme 2. We planned to use Evans' asymmetric aldol reaction to establish the C2/C3 aldol motif and to form the



**Scheme 2.** The main features of our retrosynthetic analysis. Bn = benzyl, Ms = methanesulfonyl.

THF rings in the presence of the Evans auxiliary. The most convenient precursor for the *seco* acid would be the oxazolidinone-masked hydroxy acid (2), which in turn was expected to form from the highly sensitive/unstable linear precursor 3, which contains two carbonyl groups, three leaving groups, and four hydroxy groups, and is thus a compound that may readily undergo a range of side reactions. The C1 to C21 chain was planned to be assembled from two smaller fragments, alkenes 4 and 5, through a cross metathesis reaction.

We started the synthesis with the construction of the C3 to C11 unit (Scheme 3). The asymmetric aldol reaction between **7** (readily derived from  $6^{[6]}$ ) and acrolein under the conditions described by Crimmins et al.<sup>[7]</sup> afforded the desired enantiopure **8** in 86 % yield of isolated product. Subsequent cleavage of the chiral auxiliary with DIBAL-H<sup>[7]</sup> followed by reaction with HS(CH<sub>2</sub>)<sub>3</sub>SH in the presence of BF<sub>3</sub>·Et<sub>2</sub>O gave dithiane **9**. The free OH group was then masked as TBS ether before deprotonation of the dithiane with *t*BuLi and reaction with epoxide **11** to give alcohol **12**. Use of *n*BuLi instead of *t*BuLi led to lowered yields for **12** (42 %) in this case. However, this disadvantage was partially compensated by the convenience of performing the reaction at ambient temperature without the need to add HMPA.

Angew. Chem. Int. Ed. 2012, 51, 1-5

© 2012 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

These are not the final page numbers!

🕅 WILEY 崸

<sup>[\*\*]</sup> This work was supported by the National Basic Research Program of China (the 973 Program, 2010CB833200) and the National Natural Science Foundation of China (21172247, 21032002, 20921091, 20621062).

Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/anie.201201395.



Scheme 3. Synthesis of 15. Reagents and conditions: a)  $EtCO_2H$ , EDCI, DMAP,  $CH_2CI_2$ , 96%; b)  $TiCI_4$ , (-)-sparteine, acrolein,  $CH_2CI_2$ , 0°C, 86%; c) 1) DIBAL-H,  $CH_2CI_2$ , -78°C; 2)  $HS(CH_2)_3SH$ ,  $BF_3$ ·Et<sub>2</sub>O, 69% from 8; d) TBSOTF, Et<sub>3</sub>N,  $CH_2CI_2$ , 0°C, 90%; e) tBuLi, THF, HMPA, -78°C, 71%; f) BnBr, NaHMDS, THF, DMF, 90%; g) CSA, MeOH, -10°C, 93%; h) SO<sub>3</sub>-py, DMSO, Et<sub>3</sub>N,  $CH_2CI_2$ , 90%. CSA= camphorsulfonic acid, DIBAL-H = diisobutylaluminum hydride, DMAP = 4-dimethylaminopyridine, DMF = N,N-dimethylformamide, DMSO = dimethyl sulfoxide, EDCI = 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride, HMPA = hexamethylphosphoramide, NaHMDS = sodium hexamethyldisilazide, py = pyridine, TBS = tertbutyldimethylsilyl, Tf = trifluoromethanesulfonyl.

Benzyl protection of the newly formed hydroxy group at C6 was unexpectedly difficult. The etherification with BnBr in the presence of NaH resulted in unwanted side products in pure DMSO or DMF. Fortunately, this problem was later satisfactorily solved by using a 5:1 mixture of THF/DMF as the solvent and performing the alkylation at -10 °C.

Furthermore, the incorporation of the C1/C2 unit was planned by an Evans aldol reaction. Thus, the C3 atom had to be transformed into an aldehyde first. To this end, the TBS protecting group at the primary OH group (C3) was selectively cleaved with MeOH in the presence of a catalytic amount of CSA, while the TBS group on the secondary OH group (C10) remained intact. The resulting alcohol **14** was then treated with SO<sub>3</sub>·py in CH<sub>2</sub>Cl<sub>2</sub> to deliver the expected terminal aldehyde **15**, thus setting the stage for the incorporation of the first two carbon atoms in the C1 to C11 fragment **(4)**.

The aldol reaction was achieved (Scheme 4) under the conditions described by Crimmins et al.<sup>[8]</sup> including TiCl<sub>4</sub>/ sparteine/NMP. The enantiomerically pure **17** was then treated with PIFA<sup>[9]</sup> to deprotect the thioketal protecting group. Finally, removal of the TBS group with HF afforded alkene **4**, which is needed for the later catenation with **5**.

The synthesis of fragment **5** was started with the installation of the C16/C17 stereogenic centers (Scheme 5). The aldol reaction of **19** with **20** gave **21** in 99% yield. TBS protection of the hydroxy group with TBSOTf/2,6-lutidine followed by cleavage of the oxazolidinethione auxiliary with DIBAL-H gave the intermediate aldehyde.



**Scheme 4.** Conversion of **15** into **4**. Reagents and conditions: a) **16**, TiCl<sub>4</sub>, (+)-sparteine, NMP, CH<sub>2</sub>Cl<sub>2</sub>, 0°C, 90%; b) PIFA, MeOH/H<sub>2</sub>O, 78%; c) 40% HF, MeCN, 96%. NMP=N-methyl-2-pyrrolidinone, PIFA=phenyliodine(III) bis(trifluoroacetate).



**Scheme 5.** Synthesis of **5**. Reagents and conditions: a) TiCl<sub>4</sub>, (+)-sparteine, NMP, CH<sub>2</sub>Cl<sub>2</sub>, 0°C, 99%; b) TBSOTf, 2,6-lutidine, CH<sub>2</sub>Cl<sub>2</sub>, 0°C, 95%; c) 1) DIBAL-H, CH<sub>2</sub>Cl<sub>2</sub>, -78°C, 88%; 2) **23**, TiCl<sub>4</sub>, *i*Pr<sub>2</sub>NEt, CH<sub>2</sub>Cl<sub>2</sub>, -78°C, 77%; d) 1) MeNH(OMe)·HCl, Me<sub>3</sub>Al, CH<sub>2</sub>Cl<sub>2</sub>; 2) (CH<sub>2</sub>=CH)<sub>4</sub>Sn, MeLi, THF; 3) Me<sub>4</sub>NBH(OAc)<sub>3</sub>, AcOH, MeCN, -15°C, 71% from **24**; e) 1) Me<sub>2</sub>C(OMe)<sub>2</sub>, acetone, *p*-TsOH; 2) *n*Bu<sub>4</sub>NF, THF, 93% from **25**. Ts = toluenesulfonyl.

To extend the chain by two carbon units and install a new stereogenic center at C15, an asymmetric acetate aldol reaction that was induced by an oxazolidinethione auxiliary was performed next. Generally speaking, the diastereoselectivity for such reactions is less satisfactory than that for the corresponding propionate aldol condensations under similar conditions. Several elegant protocols<sup>[10]</sup> for dealing with these types of problems have been reported. However, the advantages of using these protocols appear to be somewhat counterbalanced by, for instance, the extra efforts required for gaining access to the special auxiliaries (with sterically more hindered substituents at the stereogenic centers). In hope to find an alternative that is more suitable to us, we

www.angewandte.org

2

© 2012 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

decided to examine the aldol condensation of **23**, which to our knowledge has never been examined to date.

The results were better than those<sup>[11]</sup> observed with the oxazolidinone counterpart of **23** (i.e., with an oxygen atom in place of the sulfur in **23**). Thus, under the same conditions, including TiCl<sub>4</sub>/*i*Pr<sub>2</sub>NEt/CH<sub>2</sub>Cl<sub>2</sub>/-78 °C, the desired diastereomer **24** was isolated in enantiopure form in 77 % yield (along with 15% of the C15 epimer).

Installation of the vinyl group and the stereogenic center at C13 was mediated by the corresponding Weinreb amide. The auxiliary in **24** was cleaved by treatment with MeNH-(OMe)-HCl/Me<sub>3</sub>Al. However, the subsequent reaction with vinyl Grignard reagent was complicated by Michael addition of the MeNH(OMe) onto the vinyl double bond. Exhaustive optimization efforts included changing the reaction conditions and procedure, but were unsuccessful. Fortunately, the annoying side reaction could be greatly suppressed under the conditions described by Fuwa and Sasaki,<sup>[12]</sup> using (CH<sub>2</sub>= CH)<sub>4</sub>Sn/MeLi. The desired conjugated enone that was thus obtained was stereoselectively reduced by using the conditions described by Evans et al., including Me<sub>4</sub>NBH(OAc)<sub>3</sub><sup>[13]</sup> in AcOH/MeCN, thus giving the 1,3-diol **25** in 71% overall yield (from **24**). Further protection with Me<sub>2</sub>C(OMe)<sub>2</sub> and desilylation with  $nBu_4NF$  provided alkene 5, which was needed for coupling with compound 4.

The coupling of **4** with **5** was most satisfactorily achieved by using **26** as the catalyst with a 1:1.2 molar ratio of **4/5** (Scheme 6). The reaction proceeded rather fast at ambient temperature (affording **27** in 44% yield within just one hour). Almost all the excess **5** that was added could be recovered and recycled. Higher temperature (heating in toluene), which was normally beneficial, did not lead to any improvements, but significantly reduced the yield of **27**.

Saturation of the C=C bond in **27** without removing the Bn groups (which is critical for the selective mesylation) was much more difficult than the similar reaction in the synthesis<sup>[5]</sup> of pamamycin 621A. The presence of the carbonyl group at C8, which greatly facilitated  $\alpha$  racemization and  $\beta$  elimination, led to a failure of the reaction with the protocol that was previously rather successful (H<sub>2</sub>/Pd-C/Et<sub>3</sub>N). Fortunately, after exhaustive tries we found that these problems could be avoided by using a PtO<sub>2</sub>-catalyzed hydrogenation.

The resulting triol 28 was converted to the corresponding trimesylated compound, which on removal of the acetonide and benzyl groups provided the key precursor 3 (crude) for the ring-closing reaction. The formation of three THF rings in



Scheme 6. Synthesis of 1. Reagents and conditions: a) 26,  $CH_2Cl_2$ , RT, 24 h, 56%; b)  $H_2$ ,  $PtO_2$ , EtOAc, 89%; c) MsCl, DMAP,  $CH_2Cl_2$ , pyridine, 99%; d) 1)  $H_2$ , Pd/C, THF; 2) CSA, MeOH; 3) 2,6-lutidine, 120°C, 2 h, 60% from 29; e) TESCl, DMAP, RT, 91%; f)  $H_2O_2$ , LiOH, THF, 0°C, 97%; g) 2, 32, DMAP,  $Et_3N$ , toluene, RT, 84%; h) 1) TFA, THF; 2)  $H_2O_2$ , LiOH, THF, 0°C, 90% from 33; i) 32, DMAP,  $Et_3N$ , toluene, reflux, 5 h, 68%. Ms = methanesulfonyl, Mes = 2,4,6-trimethylphenyl, TES = triethylsilyl, TFA = trifluoroacetic acid.

Angew. Chem. Int. Ed. 2012, 51, 1-5

© 2012 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

### einheim www.angewandte.org These are not the final page numbers!

## Angewandte Communications

one step was then achieved with 2,6-lutidine under heating. Gratifyingly, **2** was obtained in 60% overall yield (from **29**).

Coupling of 2 with 31 (prepared from 2 by protection with a TES group and removal of the chiral auxiliary) under the Yamaguchi conditions provided 33, which on removal of the TES group and the chiral auxiliary gave 34. Lactonization of 34 afforded product 1, the <sup>1</sup>H and <sup>13</sup>C NMR spectra and the specific rotation ( $[\alpha]_D^{27} = +36.33$  (c = 0.20, MeOH), lit.<sup>[14]</sup> [ $\alpha$ ]<sub>D</sub><sup>17</sup> = +40.2 (c = 1.0, MeOH)) of which were consistent with those of the natural product 1.

Only the relative configuration of natural product **1** was ever experimentally determined, although the absolute configuration that was arbitrarily depicted in the report of the structure<sup>[1b]</sup> happened to be correct. In retrospect, the first synthesis<sup>[3a,b]</sup> of **1** by Lee and co-workers would have allowed to establish the absolute configuration. But unfortunately, the Lee group measured the rotation of **1** in CHCl<sub>3</sub> rather than MeOH,<sup>[15]</sup> the absolute configuration of natural product **1** thus remained unelucidated to date.

In summary, an efficient synthesis of IKD-8344 has been achieved with a novel "three rings in one step" transformation as the key step. Apart from the formation of multiple THF rings in an unprecedented difficult and complex situation, the efficient chain assembly and the tricky saturation of the C=C bond in a highly sensitive substrate are also interesting. The absolute configuration of natural product **1**, which was arbitrarily depicted in the original report of the configuration<sup>[1b]</sup> and mistaken as proven without being noticed to date, is also verified for the first time; the corresponding hidden and potential confusions in the literature and the major online data bases are thus cleared.

Received: February 20, 2012 Published online: ■■■, ■■■

**Keywords:** cyclizations · eliminations · natural products · ring-closing reactions · total synthesis

 a) Y. Minami, K. Yoshida, R. Azuma, M. Nishii, J. Inagaki, F. Nohara, *Tetrahedron Lett.* **1992**, *33*, 7373–7376; b) T. Ishida, Y. In, M. Nishii, Y. Minami, *Chem. Lett.* **1994**, 1321–1322.

- [2] E. I. Hwang, B. S. Yun, W. H. Yeo, S. H. Lee, J. S. Moon, Y. K. Kim, S. J. Lim, S. U. Kim, J. Microbiol. Biotechnol. 2005, 15, 909–912 (No optical rotation data was given).
- [3] For the total synthesis, see: a) W. H. Kim, S. K. Hong, S. M. Lim, M.-A. Ju, S. K. Jung, Y. W. Kim, J. H. Jung, M. S. Kwon, E. Lee, *Angew. Chem.* 2006, 118, 7230–7233; *Angew. Chem. Int. Ed.* 2006, 45, 7072–7075; b) W. H. Kim, S. K. Hong, S. M. Lim, M.-A. Ju, S. K. Jung, Y. W. Kim, J. H. Jung, M. S. Kwon, E. Lee, *Tetrahedron* 2007, 63, 9784–9801. For the synthesis of fragments, see: c) W. Jiang, D. A. Lantrip, P. L. Fuchs, *Org. Lett.* 2000, 2, 2181–2184.
- [4] Y.-K. Wu, Y.-P. Sun, Org. Lett. 2006, 8, 2831–2834.
- [5] a) G.-B. Ren, Y.-K. Wu, Org. Lett. 2009, 11, 5638-5641; b) G.-B. Ren, Y.-X. Huang, Y.-P. Sun, Z.-H. Li, Y.-K. Wu, J. Org. Chem. 2010, 75, 5048-5064; c) G.-B. Ren, Y.-K. Wu, Chin. J. Chem. 2010, 28, 1651-1659.
- [6] For a practical access to 6, see: Y.-K. Wu, Y.-Q. Yang, Q. Hu, J. Org. Chem. 2004, 69, 3990–3992.
- [7] a) M. T. Crimmins, K. Chaudhary, Org. Lett. 2000, 2, 775-777;
  b) M. T. Crimmins, B. W. King, E. A. Tabet, K. Chaudhary, J. Org. Chem. 2001, 66, 894-902.
- [8] M. T. Crimmins, J. She, *Synlett* **2004**, 1371–1373.
- [9] G. Stork, K. Zhao, Tetrahedron Lett. 1989, 30, 287–290.
- [10] a) N. R. Guz, A. J. Phillips, Org. Lett. 2002, 4, 2253–2256; b) Y.
   Zhang, T. Sammakia, Org. Lett. 2004, 6, 3139–3141; c) M. T.
   Crimmins, M. Shamszad, Org. Lett. 2007, 9, 149–152.
- [11] C. Le Sann, D. M. Munoz, N. Saunders, T. J. Simpson, D. I. Smith, F. Soulas, P. Watts, C. L. Willis, *Org. Biomol. Chem.* 2005, *3*, 1719–1728.
- [12] H. Fuwa, M. Sasaki, Org. Lett. 2010, 12, 584-587.
- [13] D. A. Evans, K. T. Chapman, E. M. Carreira, J. Am. Chem. Soc. 1988, 110, 3560-3578.
- [14] A full description of the specific rotation for the natural IKD-8344 can be found in: M. Nishii (Ikeda Mohando Co., Ltd.) Jpn. Kokai Tokkyo Koho, JP 02009382, 1990; *Chem. Abstr.* 1990, 113, 103380. This patent appeared to have been overlooked during the previous synthetic studies because only the incomplete rotation data (without the solvent) from Ref. [1a] was ever cited.
- [15] Comparison of rotations measured in different solvents cannot provide any reliable clues to the absolute configuration. For instance, pamamycin 621A, also a THF-containing natural product, gave opposite rotations in CHCl<sub>3</sub> and CH<sub>2</sub>Cl<sub>2</sub> (Ref. [5a,b]). The rotation for **1** in CHCl<sub>3</sub> from this work is  $[a]_D^{27} = +20.2$  (c = 0.38, CHCl<sub>3</sub>), and thus remarkably different from that in Ref. [3a,b] ( $[a]_D^{15} = +39.7$  (c = 0.25, CHCl<sub>3</sub>), the only complete rotation data for **1** (erroneously registered with a never proven absolute configuration) that can be found in leading online databases such as SciFinder and Reaxys to date).

www.angewandte.org

These are not the final page numbers!

# Communications

Natural Product Synthesis



#### Y. Zou, Y. Wu\* \_\_\_\_\_

Three Rings in One Step: A Quick Approach to IKD-8344



A highly efficient enantioselective total synthesis of the natural antibiotic IKD-8344 is achieved through a convergent route. This route features an otherwise impossible concurrent formation of the THF rings from a linear polyketide pre-



cursor through intramolecular O alkylations of mesylates in competition with normally rather facile  $\beta$  elimination and/ or  $\alpha$  racemization reactions (see scheme, Ms = methanesulfonyl).

5