This article was downloaded by: [Aston University] On: 24 August 2014, At: 09:01 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/gpss20

Insecticidal Activity of Some New Amidophosphoric Acid Esters Containing Substituted Pyridine Moieties

Feng-Mei Sun^a & De-Qing Shi^a

^a Key Laboratory of Pesticide and Chemical Biology of Ministry of Education, College of Chemistry, Central China Normal University, Wuhan, P. R. China Published online: 27 Sep 2008.

To cite this article: Feng-Mei Sun & De-Qing Shi (2008) Insecticidal Activity of Some New Amidophosphoric Acid Esters Containing Substituted Pyridine Moieties, Phosphorus, Sulfur, and Silicon and the Related Elements, 183:10, 2615-2620, DOI: 10.1080/10426500801967997

To link to this article: http://dx.doi.org/10.1080/10426500801967997

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages,

and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions



Insecticidal Activity of Some New Amidophosphoric Acid Esters Containing Substituted Pyridine Moieties

Feng-Mei Sun and De-Qing Shi

Key Laboratory of Pesticide and Chemical Biology of Ministry of Education, College of Chemistry, Central China Normal University, Wuhan, P. R. China

Some new amidophosphoric acid esters containing substituted pyridine moieties were synthesized by the reaction of O-aryl,O-ethyl phosphoryl chlorides with 3-(aminomethyl)pyridine and 5-(aminomethyl)-2-chloropyridine. Characterization was done by ¹H NMR, ³¹P{¹H} NMR, IR, MS, and elemental analyses. The results of preliminary bioassay showed that the new compounds possess good insecticidal activity against aphides.

Keywords Amidophosphoric acid esters; insecticidal activity; substituted pyridine

INTRODUCTION

Insecticides that have the structure and mechanism similar to nicotine are called neonicotinoids.¹ Seven neonicotinoids such as imidcaloprid, acetamipyrid, and nitenpyram, which have the pyridine-like moiety, have been commercialized and many SARs (structure-activity relationships) for these neonicotinoids have been reported.^{1,2} It was found that most of the biological active nicotinic compounds contain the 3-(aminomethyl)-pyridine moiety.³ Organic phosphorus insecticides especially amidophosphoric acid ester insecticides—play an important role in pesticide industry, such as methomidophos, acephate, phenamidphos, fosthiazate, mitemate, etc., have been commercialized; some neonicotinoid derivatives containing phosphorus atom have been reported.^{4,5} Herein, we hope to report the synthesis and insecticidal

Received 2 December 2007; accepted 25 January 2008.

This work was supported by the Natural Science Foundation of China (Grant No. 20302002), and the Scientific Research Foundation for the Returned Overseas Chinese Scholars, Ministry of Education of China (Grant No. [2007 1108).

Address correspondence to Prof. Dr. De-Qing Shi, College of Chemistry, Central China Normal University, Wuhan 430079, Hubei, China. Email: chshidq@yahoo.com.cn



activity of amidophosphoric acid esters containing substituted pyridine moieties.

The synthetic route is shown in Scheme 1. Characterization was done by ¹H NMR, ³¹P{¹H}NMR, IR, MS, and elemental analyses. The results of preliminary bioassay showed that the new compounds possess good insecticidal activity against aphides.

RESULTS AND DISCUSSION

The Structures of the Title Compounds

All the products were purified by flash column chromatography on a silica gel with a mixture of petroleum ether and acetone (4:1, v/v) as the eluent, The structures of compounds **IIa**~**IIi** were confirmed by ¹H NMR, ³¹P{¹H}NMR, IR spectra, MS, and elemental analyses.

For ¹H NMR spectra, the NH proton appears as a broad singlet with variable chemical shifts, the NCH₂ protons display as a doublet, in some cases as a double doublet due to the coupling with phosphorus atom and with the proton of NH. In some cases, these protons appear with equal chemical shifts with those of OCH₂ and are difficult to be distinguished. The pyridine protons are easily differentiated from the phenyl protons. The ³¹P{¹H}NMR spectra of compounds **IIa~IIi** exhibited a singlet in the range of δ 3.0 to 3.4. Their structures were also supported by correct elemental analysis and by the mass spectra, which exhibited the anticipated molecular ion peaks. The IR spectra of all compounds showed normal stretching absorption bands indicating the existence of the N-H (~ 3200 cm⁻¹), P=O (~1250 cm⁻¹), P-O-C (~950 cm⁻¹) moiety.

Biological Activities

Compounds IIa~IIi were tested for insecticidal activity against aphides by dipping at various concentrations according to a previously reported method 6. Preliminary bioassay results show that these title compounds possess good insecticidal activity at the different dosages. For example, compounds IIb, IId, IIh, and IIi exhibited 100% inhibitory rates against aphides at the dosage of 250 mg/L. Moreover, these four compounds also showed good insecticidal activity at the low concentration (50 mg/L). Table 1 showed that the insecticidal activity of compounds II depended on the nature of substituent groups in the benzene rings; compounds containing 4-chloro and 2,4-dichloro groups displayed good insectcidal activities. However, the chloro group in the pyridine moiety has less effect on their biological activities. Further structure-activity relationships are under investigation. The results of insecticidal activities are listed in Table I.

In conclusion, we synthesized some new O-aryl, O-ethyl, N-(substitutedpyridin-methyl) amidophosphoric acid esters. The results of the preliminary bioassays indicated that most of the title compounds possess good insecticidal activity.

EXPERIMENTAL

Melting points were determined with a WRS-1B digital melting point apparatus and are uncorrected. ¹H NMR and ³¹P{¹H} NMR spectra were recorded with a Varian MERCURY-PLUS400 spectrometer with TMS and 85% H₃PO₄ as the internal and external reference, respectively, and CDCl₃ as the solvent. Mass spectra were obtained with a Finnigan TRACEMS2000 spectrometer using the EI method. IR spectra were measured by a Nicolet NEXUS470 spectrometer. Elemental analyses were performed with a Vario EL **III** CHNSO elementary analyzer. The reagents and solvents were available commercially and purified according to conventional methods before use. 5-(aminomethyl)-

 TABLE I Insecticidal Activity of IIa~IIi Against Aphides: (Death Rate %)

%Dosage	IIa	IIb	IIc	IId	IIe	IIf	IIg	IIh	IIi
250 mg/L 50 mg/L	35.6	$100 \\ 95.6$	68.5 —	100 100	53.1	40.7	71.3	$100 \\ 81.3$	$100 \\ 76.4$

2-chloro-pyridine and *O*-aryl, *O*-ethyl, phosphoryl chloride **I** were prepared according to the reported methods,^{7,8} respectively.

General Procedure for the Synthesis of O-aryl, O-ethyl, N-(substitutedpyridin-methyl) Amidophosphoric Acid Esters

Compound I (2 mmol), triethylamine (0.3 g, 3 mmol) and dichloromethane (10 mL) were added to a 50 mL three-necked reaction flask, after cooling to 5°C, the solution of 3-(aminomethyl)pyridine or 5-(aminomethyl)-2-chloro-pyridine (2 mmol) in dichloromethane (5 mL) was added dropwise slowly, the mixture was allowed to stir at room temperature for 5–6 h, until the reaction was finished (monitored by TLC). The work-up involved filtration and drying over anhydrous magnesium sulfate, the crude product was purified by flash column chromatography on a silica gel with a mixture of petroleum ether and acetone (4:1, v/v) as the eluent, giving compounds IIa~IIi as a yellow oil or solid.

O-ethyl, *O*-phenyl, *N*-(pyridin-3-yl-methyl) amidophosphoric acid ester (**IIa**): an orange oil, yield 73%; IR (KBr) υ : 3216 (N-H), 1560 (C=N), 1390–1566 (C=C), 1236 (P=O), 950 (P-O-C); {¹H}NMR (CDCl₃, 400 MHz) δ :1.30 (t, J = 6.2 Hz, 3H, CH₃), 1.72 (sb, 1H, NH), 4.08–4.16 (m, 4H, 2CH₂), 7.13–7.28 (m, 6H, Ar-H, Py-H), 7.28 (d, J = 8.4 Hz, 1H, Py-H), 7.63 (d, J = 7.6 Hz, 1H, Py-H), 8.51 (s, 1H, Py-H); ³¹P{¹H}NMR (CDCl₃, 162 MHz) δ :3.05 (s); MS (70eV) m/z (%): 294 (22.9), 292 (M⁺, 66.9), 263 (34.1), 245 (4.4), 183 (29.1), 107 (100), 92 (55.5); Anal. Calcd. for C₁₄H₁₇N₂O₃P: C 57.53, H 5.86, N 9.58; found C 57.26, H 5.65, N 9.37.

O-ethyl, *O*-(4-chlorophenyl), *N*-(2-chloropyridin-5-yl-methyl) amidophosphoric acid ester (**IIb**), an orange oil, yield 68%; IR (KBr) *v*: 3219 (N-H), 1589 (C=N), 1393–1569 (C=C), 1233 (P=O), 955 (P-O-C); {¹H}NMR (CDCl₃, 400 MHz) δ : 1.32 (t, *J* = 6.8 Hz, 3H,CH₃), 1.78 (sb, 1H, NH), 4.09–4.15 (m, 4H, 2CH₂), 7.08–7.28 (m, 5H, Ar-H, Py-H), 7.59 (d, *J* = 7.6 Hz, 1H, Py-H), 8.30 (s, 1H, Py-H); ³¹P{¹H}NMR (CDCl₃, 162 MHz) δ :3.03 (s); MS (70eV) *m*/*z* (%): 362 (2.2), 360 (M⁺, 10.5), 317 (5.3), 238 (4.3), 190 (14.0), 127 (100), 111 (41.5), 99 (86), 75 (45.5); Anal. Calcd. for C₁₄H₁₅Cl₂N₂O₃P: C 46.56, H 4.19, N 7.76; found C 46.40, H 4.31, N 7.52.

O-ethyl, O-(2-methylphenyl), N-(2-chloropyridin-5-yl-methyl) amidophosphoric acid ester (**Hc**), an orange oil, yield 63%; IR (KBr) v: 3211 (N-H), 1588 (C=N), 1392–1569 (C=C), 1233 (P=O), 947 (P-O-C); ¹H NMR (CDCl₃, 400 MHz) δ : 1.25 (t, J = 6.0 Hz, 3H, CH₃), 2.21 (s, 3H, Ar-CH₃), 2.60 (sb, 1H, NH), 4.07–4.10 (m, 2H,CH₂), 4.13 (dd, J = 10.8 Hz, J = 7.6 Hz, 2H, CH₂N), 7.03–7.58 (m, 5H, Ar-H, Py-H), 7.60

 $(d, J = 8.4 \text{ Hz}, 1H, \text{Py-H}), 8.28 (s, 1H, \text{Py-H}); {}^{31}P{}^{1}H}MR (CDCl_3, 162 \text{ MHz}) \delta:3.40 (s); Anal. calcd. for C₁₅H₁₈ClN₂O₃P: C 52.87, H 5.32, N 8.22; found C 52.94, H 5.20, N 8.17.$

O-ethyl, O-(2,4-dichlorophenyl), N-(2-chloropyridin-5-yl-methyl) amidophosphoric acid ester (**IId**), an orange solid, m.p. 52–53°C, yield 66%; IR (KBr) v: 3208 (N-H), 1579 (C=N), 1392–1578 (C=C), 1240 (P=O), 939 (P-O-C); ¹H NMR (CDCl₃, 400 MHz) δ : 1.33 (t, J = 7.2 Hz, 3H,CH₃), 4.10 (dd, J = 10.4 Hz, J = 24.4 Hz, 1H, NH), 4.18–4.22 (m, 4H,2CH₂), 7.18 (d, J = 8.4 Hz, 1H, Py-H), 7.24 (d, J = 8.0 Hz, 1H, Ar-H), 7.39–7.42 (m, 2H, Ar-H), 7.61 (d, J = 8.0 Hz, 1H, Py-H), 8.31 (s, 1H, Py-H); ³¹P{¹H}MR (CDCl₃, 162 MHz) δ :3.05 (s); MS (70eV) m/z (%): 399 (14), 396 (31.1), 394 (M⁺, 42.8), 359 (43.6), 330 (25.5), 284 (25.2), 205 (16.1), 140 (97.2), 125 (100), 89 (24.4), 75 (7.1); Anal. calcd. for C₁₄H₁₄Cl₃N₂O₃P: C 42.50, H 3.57, N 7.08; found C 42.09, H 3.25, N 7.19.

O-ethyl, O-(2,3,4,5,6-pentachlorophenyl), N-(2-chloropyridin-5-ylmethyl) amidophosphoric acid ester (**IIe**), an orange solid, m.p. 43– 45°C, yield 73%; IR (KBr) υ : 3219 (N-H), 1580 (C=N), 1390-1566 (C=C), 1239 (P=O), 954 (P-O-C); ¹H NMR (CDCl₃, 400 MHz) δ : 1.25 (t, J = 5.8Hz, 3H, CH₃), 3.90–3.94 (m, 2H,CH₂), 4.05 (d, J = 11.0 Hz, 2H, CH₂N), 4.22 (sb, 1H, NH), 7.30 (d, J = 8.0 Hz, 1H, Py-H), 7.67 (d, J = 8.0 Hz, 1H, Py-H), 8.34 (s, 1H, Py-H); ³¹P{¹H}NMR (CDCl₃, 162 MHz) δ : 3.04 (s); Anal. calcd. for C₁₄H₁₁Cl₆N₂O₃P: C 33.70, H 2.22, N 5.61; found C 33.61, H 2.17, N 5.78.

O-ethyl, *O*-(2-methylphenyl), *N*-(pyridin-3-yl-methyl) amidophosphoric acid ester (**IIf**): an orange oil, yield 75%; IR (KBr) v: 3219 (N-H), 1588 (C=N), 1389–1578 (C=C), 1241 (P=O), 948 (P-O-C); ¹H NMR (CDCl₃, 400 MHz) δ : 1.29 (t, J = 5.8 Hz, 3H,CH₃), 3.28 (sb, 1H, NH), 4.06–4.08 (m, 2H,CH₂), 4.11 (d, J = 11.0 Hz, 2H, CH₂N), 7.08–7.30 (m, 5H, Ar-H, Py-H), 7.35 (d, J = 7.8 Hz, 1H, Py-H), 7.70 (d, J = 7.8 Hz, 1H, Py-H), 8.55 (s, 1H, Py-H); ³¹P{¹H}NMR (CDCl₃, 162 MHz) δ : 3.05 (s); Anal. calcd. for C₁₅H₁₉N₂O₃P: C 58.82, H 6.25, N 9.15; found C 58.71, H 6.03, N 9.31.

O-ethyl, *O*-(2,3,4,5,6-pentachlorophenyl), *N*-(pyridin-3-yl-methyl) amidophosphoric acid ester (**Hg**): an orange oil, yield 68%; IR (KBr) υ : 3215 (N-H), 1587 (C=N), 1398–1575 (C=C), 1232 (P=O), 949 (P-O-C); ¹H NMR (CDCl₃, 400 MHz) δ : 1.32 (t, J = 5.4 Hz, 3H,CH₃), 1.96 (sb, 1H, NH), 3.94-3.98 (m, 2H,CH₂), 4.10 (dd, J = 11.4 Hz, J = 8.0 Hz, 2H, CH₂N), 7.28 (d, J = 8.2 Hz, 1H, Py-H), 7.34 (d, J = 8.2 Hz, 1H, Py-H), 7.59 (d, J = 7.4 Hz, 1H, Py-H), 8.36 (s, 1H, Py-H); ³¹P{¹H}NMR (CDCl₃, 162 MHz) δ : 3.04 (s); Anal. calcd. for C₁₄H₁₂Cl₅N₂O₃P: C 36.20, H 2.60, N 6.03; found C 36.36, H 2.44, N 5.89.

O-ethyl, *O*-(4-chlorophenyl), *N*-(pyridin-3-yl-methyl) amidophosphoric acid ester (**IIh**): a light yellow oil, yield 75%; IR (KBr) v: 3225 (N-H), 1588 (C=N), 1380–1581 (C=C), 1246 (P=O), 955 (P-O-C); ¹H NMR (CDCl₃, 400 MHz) δ : 1.29 (t, J = 5.4 Hz, 3H,CH₃), 2.10 (sb, 1H, NH), 4.05–4.08 (m, 2H,CH₂), 4.13 (dd, J = 11.6 Hz, J = 7.6 Hz, 2H, CH₂N), 7.08–7.27 (m, 5H, Ar-H, Py-H), 7.58 (d, J = 7.6 Hz, 1H, Py-H), 8.47 (d, J = 4.8 Hz, 1H, Py-H), 8.50 (s, 1H, Py-H); ³¹P{¹H}NMR (CDCl₃, 162 MHz) δ : 3.05 (s); Anal. calcd. for C₁₄H₁₆ClN₂O₃P: C 51.47, H 4.94, N 8.57; found C 51.59, H 5.12, N 8.40.

O-ethyl, *O*-(2,4-dichlorophenyl), *N*-(pyridin-3-yl-methyl) amidophosphoric acid ester (**IIi**): a light yellow solid, m.p. 47–48°C, yield 65%; IR (KBr) υ : 3223 (N-H), 1587 (C=N), 1389–1568 (C=C), 1248 (P=O), 953 (P-O-C); ¹H NMR (CDCl₃, 400 MHz) δ : 1.31 (t, J = 5.4 Hz, 3H,CH₃), 3.96 (s, 1H, NH), 4.01–4.05 (m, 2H,CH₂), 4.23 (dd, J = 11.2 Hz, J = 8.0 Hz, 2H, CH₂N), 7.17–7.44 (m, 4H, Ar-H, Py-H), 7.63 (d, J = 7.6 Hz, 1H, Py-H), 8.50 (d, J = 7.6 Hz, 1H, Py-H), 8.53 (s, 1H, Py-H); ³¹P{¹H}NMR (CDCl₃, 162 MHz) δ : 3.00 (s); Anal. calcd. for C₁₄H₁₅Cl₂N₂O₃P: C 46.56, H 4.19, N 7.76; found C 46.39, H 4.13, N 7.95.

Insecticidal Bioassay Method Against Aphis by Dipping

Two cucumber leaves ca. 8 cm in diameter were dipped into the distilled water solution of the various concentrations of the test compound for a few seconds until the leaf surface was wet. After drying, the leaves were place on soil in a pet cup. Ten 2nd-instar larvae were released into the cup. The cup was covered with a lid and stored at 25° C, 50-55% relative humidity (RH), and 14 h light/10 h dark for 5 days. The mortality was assessed after treatment. The test was run three times, the results were averaged, 0 means no effect, and 100% means excellent insecticidal activity, and given as death rate in Table I.

REFERENCES

- I. Yamamoto, Nicotinoid Insecticides and the Nicotinic Acetylcholine Receptor, I. Yamamoto and J. E. Casida, Eds. (Springer, Tokyo, 1999), pp. 3–27.
- [2] T. Wakita, K. Konoshita, N. Yasui, N. Kawahara, A. Naoi, M. Nakaya, H. Matsuno, and K. Kodaka, *Pest Manag. Sci.*, 59, 1016 (2003).
- [3] I. Yamamoto, G. Yabita, M. Tomizawa, et al., J. Pesticide Sci., 19, 335 (1994).
- [4] Z.J. Liu, L.M. Hu, X.H. Qin, H.W. He, S.Y. Xiao, S.L. Deng, et al. Chin. Patent, CN 1213664A (1999).
- [5] K. Nanjo, A. Kariya, and S. Henmi. U.S. Patent, 5405961 (1993).
- [6] K. Kiriyama, S. Kagabu, and K. Nishimura. J. Pestic. Sci., 29, 43 (2004).
- [7] X. Liu, R.Q. Huang, M.R. Cheng, et al., Chem. J. Chinese Univ., 20 (9), 1404 (1999).
- [8] O.M. Friedman and A.M. Seligman, J. Amer. Chem. Soc., 76, 655 (1954).