



## Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

### A Convenient One-Pot Synthesis of Phosphino-Dipeptide Analogs

Jean-Luc Pirat <sup>a</sup>, Agnes Coulombeau <sup>a</sup>, Arielle Genevois-Borella <sup>b</sup> & Henri-Jean Cristau <sup>a</sup>

<sup>a</sup> Laboratoire de Chimie Organique, C.N.R.S., E.N.S.C.M., Montpellier Cedex, France

<sup>b</sup> Centre de Recherche de Vitry-Alfortville, Aventis, Vitry-Sur-Seine, France

Published online: 27 Oct 2010.

To cite this article: Jean-Luc Pirat, Agnes Coulombeau, Arielle Genevois-Borella & Henri-Jean Cristau (2002) A Convenient One-Pot Synthesis of Phosphino-Dipeptide Analogs, *Phosphorus, Sulfur, and Silicon and the Related Elements*, 177:6-7, 1793-1796, DOI: [10.1080/10426500212259](https://doi.org/10.1080/10426500212259)

To link to this article: <http://dx.doi.org/10.1080/10426500212259>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>



## A CONVENIENT ONE-POT SYNTHESIS OF PHOSPHINO-DIPEPTIDE ANALOGS

Jean-Luc Pirat,<sup>a</sup> Agnes Coulombeau,<sup>a</sup> Arielle Genevois-Borella,<sup>b</sup>  
and Henri-Jean Cristau<sup>a</sup>

Laboratoire de Chimie Organique, C.N.R.S, E.N.S.C.M.,  
Montpellier Cedex, France<sup>a</sup> and Aventis, Centre de Recherche de  
Vitry-Alfortville, Vitry-Sur-Seine, France<sup>b</sup>

(Received February 28, 2001; accepted December 25, 2001)

*A general “one-pot” synthetic method is described for the preparation of phosphinodipeptides of type 1, in 60–80% overall yield, allowing the variation of the substituents in  $\alpha$  and/or  $\beta$  position to the phosphorus atom and also in  $\alpha$  position to the nitrogen atom.*


**Keywords:** Alkyl hypophosphite; aminoalkylphosphinic acids; Kabachnik–Fields reaction; Michael addition; phosphinopeptides

Analogous to  $\alpha$ -amino-carboxylic acids and their derived peptides, 1-aminoalkylphosphonic acids, and the corresponding peptides (phosphinopeptides) interfere with biological mechanisms through their hydroxyphosphonyl function.<sup>1</sup> The phosphinopeptides **1**, P–C analogs of phosphonopeptides, are better candidates for the elaboration of more stable biologically active compounds. Construction of such compounds requires the synthesis of the phosphinodipeptides analogs of type **1** as building blocks.

Compounds **1** are generally prepared in a multistep synthesis from the corresponding adequately protected 1-aminoalkylphosphonous acids.<sup>2</sup>

We developed a general “one-pot” three-step synthesis for the preparation of phosphinodipeptides of type **1** affording the possibility of variation of R<sup>1</sup> and R<sup>2</sup> groups (Figure 1). Each step of this reaction was improved separately until the yield was more than 90%; then all the steps were carried out without purification, for combinatorial or parallel synthesis.

Address correspondence to Jean-Luc Pirat, Laboratoire de Chimie Organique, UMR 5076 du C.N.R.S., E.N.S.C.M., 8 rue de l'École Normale, 34296 Montpellier, Cedex, France.



**FIGURE 1** Synthesis of phosphino-dipeptide analogs.

## RESULTS AND DISCUSSION

### Synthesis of Alkyl Hypophosphites 2


Alkyl hypophosphites, unstable but more reactive than the hypophosphorous acid, were preferred to the bis(trimethylsilyl) phosphonite, which is highly pyrophoric (Step 1, Figure 2).

In spite of many changes in the reaction conditions, the poor yield obtained in pure ethyl hypophosphite **2a** does not permit one to proceed with the synthesis using ethyl hypophosphite as starting material. However, methyl hypophosphite **2b** could be generated *in situ*<sup>3</sup> in very good yield and at good purity (95 to 99%). Therefore we decided to use methyl hypophosphite **2b** for the next step.


### Synthesis of Methyl Phosphinate 3

Improving on the Baylis method,<sup>4</sup> the intermediate 1-amino-alkylphosphonous acids **3** were prepared by a dropwise addition of an excess (1.6 equivalents) of several imines (or triazine), to methyl hypophosphite **2b**, in refluxing dry methanol (Figure 3). Thus, the methyl phosphinates **3** were obtained in good yield (79–89%), and the purity could be increased up to 85–95% by simple precipitating out of by-products.

Products **3b** were then used directly, without additional purification, in Step 3 of the synthesis.



**FIGURE 2** Preparation of alkyl hypophosphite **2** (Step 1).



**FIGURE 3** Preparation of methyl phosphinates **3** (Step 2).

## Synthesis of Phosphinodipeptide Analogs 1


The last step of the synthesis (Step 3), a Michael addition of the phosphinates **3b** to various  $\alpha$ -substituted acrylates was performed according to the method developed by Parsons et al., but by using *t*-BuOK in tetrahydrofuran (THF), an overcrowded base in catalytic amount, instead of MeONa in MeOH, as basic activating agent in order to reduce the demethylation of the phosphinate as side reaction. Compounds **1b**, from differently substituted acrylates, are obtained in 78–87% yields (Figure 4) as a mixture of diastereoisomers that can be partially separated by chromatography.

## Selective and Total Deprotection of Phosphinodipeptide Analogs 1

In order to show the value of phosphinodipeptide analogs **1** as building blocks, examples of selective deprotection of the various functional groups were performed, by acid (47% HBr) or basic hydrolysis (NaOH), by hydrogenolysis ( $\text{H}_2/\text{Pd}\cdot\text{C}$ ), or by methanolysis ( $\text{BrSiMe}_3$ ; MeOH) in almost quantitative yields.

## CONCLUSION

A general three-step synthesis has been developed for the “one-pot” preparation of phosphinodipeptides **1**, as a mixture of diastereoisomers,



**FIGURE 4** Preparation of phosphinopeptides analogs **1** (Step 3).

in 60–80% overall yield, which allows variation of the substituents on the carbon in positions  $\alpha$  and/or  $\beta$  to the phosphorus.

Selective or complete cleavages of the various protective groups were performed, in usual ways, in almost quantitative yields.

## REFERENCES

- [1] P. Kafarski and P. Mastalerz, In *Beiträge Zur Wirkstoffforschung: Aminophosphonates*, ed. P. Oehme, H. Löwe, E. Göres, and J. Axt (Institut für Wirkstoffforschung, Berlin, 1984), vol. 21, pp. 1–110.
- [2] (a) H. J. Cristau, A. Coulombeau, A. Genevois-Borella, and J. L. Pirat, *Tetrahedron Lett.*, **42**, 4491 (2001); (b) H. J. Cristau, A. Coulombeau, A. Genevois-Borella, F. Sanchez, and J. L. Pirat, *J. Organomet. Chem.*, **643–644**, 381 (2002).
- [3] (a) S. J. Fitch, *J. Am. Chem. Soc.*, **86**, 61 (1964); (b) M. Gallagher and H. Honegger, *Tetrahedron Lett.*, **34**, 2987 (1977); (c) A. W. Schwabacher and A. D. Stefanescu, *Tetrahedron Lett.*, **37**, 25 (1996).
- [4] E. K. Baylis, C. D. Campbell, and J. G. Dingwall, *J. Chem. Soc. Perkin Trans. 1*, **12**, 2845 (1984).