

Palladium-catalyzed Direct Monoarylation of Thiophene-, Benzothiophene-, and Indoleacetic Acids through Regioselective C–H Bond Cleavage


Daisuke Takeda, Mana Yamashita, Koji Hirano, Tetsuya Satoh,* and Masahiro Miura*
 Department of Applied Chemistry, Faculty of Engineering, Osaka University, Suita, Osaka 565-0871

(Received April 20, 2011; CL-110332; E-mail: satoh@chem.eng.osaka-u.ac.jp, miura@chem.eng.osaka-u.ac.jp)


Regioselective monoarylation of thiophene-3-acetic acid with aryl bromides proceeds efficiently under palladium catalysis to selectively give the corresponding C2-arylated thiophene derivatives. The procedure is also applicable to the C2-arylations of benzothiophene-3- and indole-3-acetic acids.

Since 2-arylthiophene frameworks can be seen in a wide range of biologically active agents,¹ the development of methods for their effective construction has attracted considerable attention. 2-Arylthiophene-3- and 2-arylbenzothiophene-3-acetic acids are known as their key synthetic intermediates (Scheme 1).¹

Among promising methods for making a thienyl–aryl bond is transition-metal-catalyzed direct arylation on thiophenes with aryl halides.² For example, we have reported that 3-cyanothiophene undergoes palladium-catalyzed arylation efficiently at the C2- and C5-positions (Scheme 2a).³ As in this example, it is known that diarylated together with monoarylated products are usually formed, often predominantly, in the reaction of 2,5-unsubstituted thiophenes. Thus, the regioselective monoarylation has so far been limited to only a few examples including those of 3-methoxy-⁴ and 3-alkoxycarbonylthiophenes⁵ (Schemes 2b and 2c).



Scheme 1.



Scheme 2.


In the context of our study of the direct functionalization of heteroarenes,⁶ we have undertaken the palladium-catalyzed arylation of thiophene-3- and benzothiophene-3-acetic acids with aryl bromides. As a result, the reaction has been found to take place selectively at the C2-position to produce 2-arylthiophene-3- and 2-arylbenzothiophene-3-acetic acid derivatives. The new findings are described herein.

In an initial attempt, thiophene-3-acetic acid (**1a**) (0.4 mmol) was treated with bromobenzene (**2a**) (0.5 mmol) in the presence of Pd(OAc)₂ (0.02 mmol), biphenyl-2-ylidicyclohexylphosphane (L1, 0.04 mmol), and K₂CO₃ (0.6 mmol) as catalyst, ligand, and


Table 1. Reaction of thiophene-3-acetic acid (**1a**) with aryl bromides **2a**

Entry	2	L (mmol) ^b	Temp/°C	Product(s), Yield/% ^c
1 ^d	2a : R = H	L1 (0.04)	120	3a : R = H, 81 ^e
2	2a : R = H	L1 (0.04)	120	3a : R = H, 85
3	2a : R = H	L2 (0.04)	120	3a : R = H, 94 (90)
4	2b : R = Me	L1 (0.04)	120	3b : R = Me, >99 (85)
5	2c : R = OMe	L1 (0.08)	120	3c : R = OMe, 78 (64)
6	2d : R = Cl	L1 (0.04)	100	3d : R = Cl, 89 (80)
7	2e : R = CF ₃	L2 (0.04)	85	3e : R = CF ₃ , 93 (84)
8	2f : R = CO ₂ Et	L2 (0.04)	85	3f : R = CO ₂ Et, 95 (92)
9	2g : R = CN	L2 (0.04)	85	3g : R = CN, >99 (92)
10	2h :	L1 (0.04)	120	3h , 99 (82)
11	2i :	L1 (0.08)	120	3i , 65 (65)

^aReaction conditions: 1) **1a** (0.5 mmol), **2** (0.4 mmol), Pd(OAc)₂ (0.02 mmol), K₂CO₃ (0.6 mmol), in DMF (2.5 mL) for 8 h under N₂; 2) with the addition of MeI (6 mmol) and K₂CO₃ (3 mmol) at room temperature for 3 h. ^bL1 = biphenyl-2-ylidicyclohexylphosphane, L2 = PPh₃. ^cGC yield based on the amount of **2** used. Value in parentheses indicates yield after isolation. ^d**1a** (0.4 mmol) and **2a** (0.5 mmol) were used. ^eMinor amounts of diphenylated products were also formed (total 17%).




Scheme 3.



Scheme 4.

Table 2. Reaction of benzo[*b*]thiophene-3-acetic acid (**1b**) with aryl bromides **2^a**




Entry	2	R	L (mmol) ^b	Temp / °C	Product, Yield/% ^c
1	2a	R = H	L2 (0.04)	120	5a : R = H, 94
2	2a	R = H	L2 (0.04)	100	5a : R = H, >99 (99)
3	2a	R = H	L1 (0.04)	100	5a : R = H, 90
4	2b	R = Me	L1 (0.08)	100	5b : R = Me, >99 (96)
5	2c	R = OMe	L1 (0.08)	100	5c : R = OMe, 80 (68)
6	2d	R = Cl	L1 (0.04)	100	5d : R = Cl, 81 (61)
7	2e	R = CF ₃	L2 (0.04)	100	5e : R = CF ₃ , 91 (74)
8	2e	R = CF ₃	L2 (0.04)	85	5e : R = CF ₃ , 78

^aReaction conditions: 1) **1b** (0.4 mmol), **2** (0.5 mmol), Pd(OAc)₂ (0.02 mmol), K₂CO₃ (0.6 mmol), in DMF (2.5 mL) for 8 h under N₂; 2) with the addition of MeI (6 mmol) and K₂CO₃ (3 mmol) at room temperature for 3 h. ^bL1 = biphenyl-2-ylidicyclohexylphosphane, L2 = PPh₃. ^cGC yield based on the amount of **1b** used. Value in parentheses indicates yield after isolation.

base, respectively, in DMF at 120 °C for 8 h under N₂. After the subsequent methyl esterification using iodomethane for quantification, the C2-monophenylated product **3a** was obtained in 81% yield, along with minor amounts of diphenylated products (total 17%) (Entry 1 in Table 1). In the presence of a slight excess of **1a**, formation of diphenylated products was completely suppressed to selectively give **3a** in 85% yield (Entry 2). Less expensive PPh₃ (L2) could also be employed as ligand (Entry 3). Using L1 at 120 or 100 °C, the reactions of **1a** with methyl- (**2b**), methoxy- (**2c**), and chloro- (**2d**) substituted bromobenzenes proceeded smoothly to produce the corresponding C2-arylated products **3b–3d** (Entries 4–6). In these cases, the use of L2 as ligand decreased the product yields due to contamination by phenyl groups from L2 to form small amounts of **3a**.⁷ In contrast, in the reactions of electron-deficient bromobenzenes **2e–2g** reacting with **1a**, such contamination was

Table 3. Reaction of indole-3-acetic acids **1** with aryl bromides **2^a**



Entry	1	2	Temp/°C	Product, Yield/% ^b
1	1c	2a : R = H	120	6a : R = H, >99 (93)
2		2b : R = Me	120	6b : R = Me, (94)
3		2c : R = OMe	120	6c : R = OMe, (94)
4		2d : R = Cl	120	6d : R = Cl, (91)
5		2e : R = CF ₃	120	6e : R = CF ₃ , (91)
6		2f : R = CO ₂ Et	120	6f : R = CO ₂ Et, (86)
7		2g : R = CN	120	6g : R = CN, (87)
8		2j : R = CHO	120	6h : R = CHO, (79)
9		2k : R = 4-R'C ₆ H ₄ (R' = pentyl)	120	6i : R = 4-R'C ₆ H ₄ , (84) (R' = pentyl)
10 ^c	1c	2l	140	6j , (79)
11	1c	2h	120	6k , (87)
12 ^c	1c	2i	140	6l , (60)
13 ^d	1c	2m	120	6m , (41)
14	1d	2a : R = H	140	6n : R = H, (75)
15		2n : R = Me	140	6o : R = Me, (78)
16		2o : R = Cl	140	6p : R = Cl, (74)
17	1e	2a	120	6q , (84)

^aReaction conditions: 1) **1** (0.4 mmol), **2** (0.5 mmol), Pd(OAc)₂ (0.008 mmol), biphenyl-2-ylidicyclohexylphosphane (0.016 mmol), K₂CO₃ (0.9 mmol), in DMF (2.5 mL) for 8 h under N₂; 2) with the addition of MeI (5 mmol) and K₂CO₃ (1 mmol) at room temperature for 3 h. ^bGC yield based on the amount of **1** used. Value in parentheses indicates yield after isolation.

^cbiphenyl-2-ylidicyclohexylphosphane (0.032 mmol) was used.

^dWithout biphenyl-2-ylidicyclohexylphosphane.

not observed even using L2 at 85 °C to afford **3e–3g** in good yields (Entries 7–9). Besides bromobenzenes, 2-bromonaphthalene (**2h**) and 2-bromo-5-methylthiophene (**2i**) also underwent the reaction with **1a** in the presence of L1 (Entries 10 and 11).

A plausible mechanism for the arylation of **1a** with **2** is illustrated in Scheme 3, in which neutral ligands are omitted. First, oxidative addition of **2** toward Pd⁰ species generated in situ followed by ligand exchange with **1a** gives an arylpalladium carboxylate intermediate **A**. Then, directed palladation⁸ on the thiophene ring may occur to form an aryl(thienyl)palladium species **B**. Final reductive elimination affords a C2-arylated product and regenerates Pd⁰ species. It should be noted that treatment of methyl thiophene-3-acetate with **2a** gave **3a** only in a low yield (Scheme 4). Therefore, coordination of the carboxylic group of **1a** appears to be the key for the effective palladation in **A**.⁹

Benzo[*b*]thiophene-3-acetic acid (**1b**) also underwent the phenylation upon treatment with **2a** in the presence of the Pd(OAc)₂/L2 catalyst system at 120 °C to afford the corresponding C2-phenylated product **5a** in 94% yield (Entry 1 in Table 2). At 100 °C, **5a** was obtained quantitatively (Entry 2). In the reactions of **2b–2d** with **1b**, L1 was used in place of L2 to avoid the contamination as in the reaction with **1a** (Entries 4–6). In the cases using **2b** and **2c**, addition of the increased amount of L1 improved the product yields. The reaction of **1b** with **2e** took place efficiently in the presence of the Pd(OAc)₂/L2 catalyst system at 100 °C, rather than at 85 °C (Entry 7 vs. 8).

Nitrogen-containing analogs, 2-arylindole-3-acetic acids, have also attracted attention because of their interesting biological properties.¹⁰ Therefore, we next examined their synthesis through the direct arylation of readily available indole-3-acetic acids. Treatment of *N*-methylindole-3-acetic acid (**1c**) (0.4 mmol) with **2a** (0.5 mmol) in the presence of Pd(OAc)₂ (0.008 mmol), L1 (0.016 mmol), and K₂CO₃ (0.9 mmol) in DMF at 120 °C for 8 h under N₂ and subsequent methyl esterification gave a C2-phenylated product **6a** in a quantitative yield (Entry 1 in Table 3). Under the same conditions, a number of 4-substituted bromobenzenes **2b–2k** reacted with **1c** efficiently to produce the corresponding 2-arylindole derivatives (Entries 2–9). Similarly, 2-bromotoluene (**2l**), 2-bromonaphthalene (**2h**), 2-bromo-5-methylthiophene (**2i**), and β-bromostyrene (**2m**, E:Z = 6.5:1) underwent the reaction at 120–140 °C (Entries 10–13). In the case with **2m**, the product yield was decreased by the addition of L1 or L2. The arylation of *N*-unsubstituted and *N*-phenylated indole-3-acetic acids, **1d** and **1e**, with bromides **2a**, **2n**, and **2o** took place effectively to form products **6n–6q** in 74–84% yields (Entries 14–17).

In summary, we have demonstrated that thiophene-3-, benzothiophene-3-, and indole-3-acetic acids undergo regioselective arylation upon treatment with aryl bromides under palladium catalysis to afford the corresponding C2-arylated products of pharmaceutical interest.¹¹

This work was partly supported by Grants-in-Aid from MEXT and JSPS, Japan.

This paper is in celebration of the 2010 Nobel Prize awarded to Professors Richard F. Heck, Akira Suzuki, and Ei-ichi Negishi.

References and Notes

- For example, see: a) W. F. Fobare, W. R. Solvibile, A. J. Robichaud, M. S. Malamas, E. Manas, J. Turner, Y. Hu, E. Wagner, R. Chopra, R. Cowling, G. Jin, J. Bard, *Bioorg. Med. Chem. Lett.* **2007**, *17*, 5353. b) N. Floquet, C. Richez, P. Durand, B. Maigret, B. Badet, M.-A. Badet-Denisot, *Bioorg. Med. Chem. Lett.* **2007**, *17*, 1966. c) J. G. Park, P. C. Sill, E. F. Makiyi, A. T. Garcia-Sosa, C. B. Millard, J. J. Schmidt, Y.-P. Pang, *Bioorg. Med. Chem.* **2006**, *14*, 395. d) M. J. Kukla, C. M. Woo, J. R. Kehr, A. Miller, *J. Med. Chem.* **1978**, *21*, 348.
- For selected recent reviews concerning direct C–H functionalization, see: a) L. Ackermann, *Chem. Rev.* **2011**, *111*, 1315. b) K. Hirano, M. Miura, *Synlett* **2011**, 294. c) J. Roger, A. L. Gottumukkala, H. Doucet, *ChemCatChem* **2010**, *2*, 20. d) A. A. Kulkarni, O. Daugulis, *Synthesis* **2009**, 4087. e) F. Bellina, R. Rossi, *Tetrahedron* **2009**, *65*, 10269. f) L. Ackermann, R. Vicente, A. R. Kapdi, *Angew. Chem., Int. Ed.* **2009**, *48*, 9792. g) X. Chen, K. M. Engle, D.-H. Wang, J.-Q. Yu, *Angew. Chem., Int. Ed.* **2009**, *48*, 5094. h) O. Daugulis, H.-Q. Do, D. Shabashov, *Acc. Chem. Res.* **2009**, *42*, 1074. i) G. P. McGlacken, L. M. Bateman, *Chem. Soc. Rev.* **2009**, *38*, 2447. j) F. Kakiuchi, T. Kochi, *Synthesis* **2008**, 3013. k) J. C. Lewis, R. G. Bergman, J. A. Ellman, *Acc. Chem. Res.* **2008**, *41*, 1013. l) A. Mori, A. Sugie, *Bull. Chem. Soc. Jpn.* **2008**, *81*, 548. m) T. Satoh, M. Miura, *Chem. Lett.* **2007**, *36*, 200. n) D. Alberico, M. E. Scott, M. Lautens, *Chem. Rev.* **2007**, *107*, 174. o) I. V. Seregin, V. Gevorgyan, *Chem. Soc. Rev.* **2007**, *36*, 1173. p) K. Godula, D. Sames, *Science* **2006**, *312*, 67.
- a) T. Okazawa, T. Satoh, M. Miura, M. Nomura, *J. Am. Chem. Soc.* **2002**, *124*, 5286. See also: b) L. Lavenot, C. Gozzi, K. Ilg, I. Orlova, V. Penalva, M. Lemaire, *J. Organomet. Chem.* **1998**, *567*, 49.
- a) S. Yanagisawa, T. Sudo, R. Noyori, K. Itami, *Tetrahedron* **2008**, *64*, 6073. b) S. Yanagisawa, T. Sudo, R. Noyori, K. Itami, *J. Am. Chem. Soc.* **2006**, *128*, 11748.
- B. Glover, K. A. Harvey, B. Liu, M. J. Sharp, M. F. Tymoschenko, *Org. Lett.* **2003**, *5*, 301.
- a) T. Ueyama, S. Mochida, T. Fukutani, K. Hirano, T. Satoh, M. Miura, *Org. Lett.* **2011**, *13*, 706. b) M. Kitahara, N. Umeda, K. Hirano, T. Satoh, M. Miura, *J. Am. Chem. Soc.* **2011**, *133*, 2160. c) M. Miyasaka, K. Hirano, T. Satoh, M. Miura, *J. Org. Chem.* **2010**, *75*, 5421. d) T. Kawano, K. Hirano, T. Satoh, M. Miura, *J. Am. Chem. Soc.* **2010**, *132*, 6900. e) N. Matsuyama, M. Kitahara, K. Hirano, T. Satoh, M. Miura, *Org. Lett.* **2010**, *12*, 2358. f) H. Hachiya, K. Hirano, T. Satoh, M. Miura, *Angew. Chem., Int. Ed.* **2010**, *49*, 2202. g) M. Yamashita, H. Horiguchi, K. Hirano, T. Satoh, M. Miura, *J. Org. Chem.* **2009**, *74*, 7481. h) M. Miyasaka, A. Fukushima, T. Satoh, K. Hirano, M. Miura, *Chem.—Eur. J.* **2009**, *15*, 3674. i) M. Nakano, H. Tsurugi, T. Satoh, M. Miura, *Org. Lett.* **2008**, *10*, 1851. j) A. Maehara, H. Tsurugi, T. Satoh, M. Miura, *Org. Lett.* **2008**, *10*, 1159.
- M. Miura, S. Pivsa-Art, T. Satoh, M. Nomura, M. Miura, G. Dyker, J. Heiermann, *Chem. Commun.* **1998**, 1889.
- a) T. Satoh, M. Miura, *Synthesis* **2010**, 3395. b) L. J. Gooßen, N. Rodríguez, K. Gooßen, *Angew. Chem., Int. Ed.* **2008**, *47*, 3100.
- Pd-Catalyzed oxidative ortho-arylation of phenylacetic acids with arylborates has been reported: D.-H. Wang, T.-S. Mei, J.-Q. Yu, *J. Am. Chem. Soc.* **2008**, *130*, 17676.
- For example, see: a) K. D. Dykstra, L. Guo, E. T. Birzin, W. Chan, Y. T. Yang, E. C. Hayes, C. A. DaSilva, L.-Y. Pai, R. T. Mosley, B. Kraker, P. M. D. Fitzgerald, F. DiNinno, S. P. Rohrer, J. M. Schaeffer, M. L. Hammond, *Bioorg. Med. Chem. Lett.* **2007**, *17*, 2322. b) B. W. Trotter, A. G. Quigley, W. C. Lumma, J. T. Sisko, E. S. Walsh, C. S. Hamann, R. G. Robinson, H. Bhimnathwala, D. G. Kolodin, W. Zheng, C. A. Buser, H. E. Huber, R. B. Lobell, N. E. Kohl, T. M. Williams, S. L. Graham, C. J. Dinsmore, *Bioorg. Med. Chem. Lett.* **2001**, *11*, 865. c) A. P. Kozikowski, D. Ma, J. Brewer, S. Sun, E. Costa, E. Romeo, A. Guidotti, *J. Med. Chem.* **1993**, *36*, 2908. d) A. Misato, K. Kou, M. Okada, M. Takami, M. Ishiguro, Y. Ichihara, H. Oomura, Jpn. Kokai Tokkyo Koho JP 55151505 (A), **1980**.
- Supporting Information is available electronically on the CSJ-Journal Web site, <http://www.csj.jp/journals/chem-lett/index.html>.