

A EUROPEAN JOURNAL OF CHEMICAL BIOLOGY

CHEMBIOCHEM

SYNTHETIC BIOLOGY & BIO-NANOTECHNOLOGY

Accepted Article

Title: Exploring the structural space of galectin-1-ligand interaction

Authors: Nadja Bertleff-Zieschang, Julian Bechold, Clemens Grimm,
Michael Reutlinger, Petra Schneider, Gisbert Schneider, and
Juergen Seibel

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *ChemBioChem* 10.1002/cbic.201700251

Link to VoR: <http://dx.doi.org/10.1002/cbic.201700251>

WILEY-VCH

www.chembiochem.org

A Journal of



COMMUNICATION

Exploring the structural space of galectin-1-ligand interaction

Nadja Bertleff-Zieschang[‡], Julian Bechold[‡], Clemens Grimm*, Michael Reutlinger, Petra Schneider, Gisbert Schneider and Jürgen Seibel*

Dedicated to Professor Lutz Friedjan Tietze on the occasion of his 75th birthday.

Abstract: Galectin-1 is a tumor-associated protein recognizing the Gal β 1-4GlcNAc motif of cell surface glycoconjugates. Here, we report on the stepwise expansion of a multifunctional natural scaffold based on *N*-acetylglucosamine (LacNAc). We obtained a LacNAc mimetic equipped with an alkyne function on the 3'-hydroxyl group of the disaccharide facing towards a binding pocket adjacent to the carbohydrate recognition domain. It served as an anchor motif for further expansion *via* Sharpless-Huisgen-Meldal reaction, resulting in ligands with an almost identical binding mode to the natural carbohydrate template. X-ray crystallography provided a structural understanding of the galectin-1-ligand interactions. The results of this study enable the development of bespoke ligands for members of the galectin target family.

Human galectin-1 is a homodimer of 14.5 kDa subunits and a member of carbohydrate binding lectins.^[1] It has been associated with tumor proliferation, angiogenesis and metastasis.^[2] Galectin-1 secreting tumors gain immune privilege by inducing apoptosis in T-helper cells,^[3] and its expression level correlates with the aggressiveness of certain tumors.^[4] Many of these processes are mediated by the sophisticated interplay between the lectin and the β -galactosides of complex glycan structures presented on the cell surface. Due to the overexpression of galectin-1 in numerous types of cancer cell lines, the protein has been suggested as a potential target for therapeutic intervention^[5] and a diagnostic

marker^[6]. Recent studies have demonstrated that multivalent presentation of natural β -galactosides (*N*-acetylglucosamine (LacNAc), lactose) compensates for the weak affinity (low mM range) of galectin-1 to the carbohydrate ligand and results in high-avidity binding.^[7] In this context, the development of synthetic galectin-1 ligands, based on the natural binding motif LacNAc to interfere with physiological interactions or to probe for elevated galectin-1 levels, has previously been pursued by combinatorial approaches.^[8]

The galectin-1 carbohydrate recognition domain (CRD) features a main binding site for the respective β -galactoside, (defined by His44, Asn46, Arg48, His52, Trp68, Asp54, Val59, Asn61, Glu71, Arg73, surface area depicted in blue, Figure 1a), which also accommodates the *N*-acetylglucosamine residue of LacNAc (Figure 1a). However, previous studies have shown that oligosaccharides like Neu5Ac- α -(2,3)-LacNAc^[9] bind to an adjacent pocket to the left side (Ala121, Ala122, Asp123, Val31, Leu32, Asn33, Leu34, Gly35, green surface region). The goal of this present study was to gradually expand LacNAc into a tailor-made ligand able to reach this and a previously unexplored cleft on the protein surface (Asp25, Asp126, red surface region). This part of the galectin-1 surface had not been addressed previously, and was chosen to demonstrate our rational approach to design a ligand for a specific protein or binding site.

In a preliminary experiment using isothermal titration calorimetry (ITC), we observed that the interaction between galectin-1 and LacNAc ($K_d = 81 \pm 12 \mu\text{M}$) is enthalpically driven and suffers from entropic losses (Supplementary Information, Table S1). Consequently, the predictability of synthetically extended disaccharide structures by computational methods is limited.^{[10][11]} To avoid artefacts from structural modeling, we directly crystallized the complex of human galectin-1 bound to LacNAc and solved its structure with a resolution of 1.4 Å (Figure 1a, PDB ID: 4Q26). This crystal structure features high similarity to the complex of human galectin-1 with lactose (PDB ID: 1GZW, RMSD = 0.28 Å)^[10a] as well as to bovine galectin-1 with LacNAc (PDB ID: 1SLT, RMSD = 0.55 Å)^[12]. In all three structures, the disaccharide ligand displays a canonical binding mode within the binding site (highlighted in blue, Figure 1)^[12].

The crystallographic data suggested that the functionalization of the 3'-OH of the galactosyl unit and the 1-OH of the *N*-acetylglucosamine moiety (as β -anomer) would not strongly influence the binding of the sugar and might yield a template that mimics the LacNAc binding motif. The 3'-OH group has already been proposed for modifications,^[8, 13] and apparently forms a favorable exit vector pointing away from the *N*-acetylglucosamine core, facing towards the surface pocket adjacent to the CRD. In order to provide a reactive and synthetically accessible form of the natural galectin-1 ligand, we

[*] Dr. N. Bertleff-Zieschang, J. Bechold, Prof. Dr. J. Seibel
Institute of Organic Chemistry
Julius Maximilians-Universität Würzburg
Am Hubland, 97074 Würzburg, Germany
E-mail: seibel@chemie.uni-wuerzburg.de

Dr. C. Grimm
Biozentrum der Julius Maximilians-Universität Würzburg
Am Hubland, 97074 Würzburg, Germany
E-mail: clemens.grimm@biozentrum.uni-wuerzburg.de

Dr. M. Reutlinger, Dr. P. Schneider, Prof. Dr. G. Schneider
Eidgenössische Technische Hochschule (ETH)
Department Chemie und Angewandte Biowissenschaften
Vladimir-Prelog-Weg 4, 8093 Zürich, Switzerland

[**] We thank the FCI (N.B.-Z.) and ETH Zürich (G.S.) for financial support.

[‡] The authors contributed equally to this work.

Supporting information for this article (details on synthesis, X-ray crystallography, computational methods and binding assay) is available on the WWW under <http://dx.doi.org/10.1002/anie.2017xxxxx>.

COMMUNICATION

chose the alkyne and alkene groups as modifications of the 3'-OH and 1-OH groups, respectively (Scheme 1). Both functional groups can be converted under mild conditions with excellent yields, high selectivity and compatibility with a wide scope of other functional groups^[14]. Terminal alkynes react in the Cu(I)-catalyzed Sharpless-Huisgen-Meldal click reaction with azides to 1,4-triazoles.^[15] The alkene group modification at 1-OH is able to functionalize the resulting compounds with thiols^[16], alkenes in the metathesis reaction,^[17] or with tetrazines in an inverted Diels-Alder^[18] reaction. This option offers opportunities for drug loading, microarray experiments or the formation of multivalent structures in future studies.

In this regard, 3-O-propargyl- β -D-galactopyranosyl-(1,4)-2-acetamido-1-O-allyl-2-deoxy- β -D-glucopyranoside **1** was conceived as a multifunctional LacNAc mimic. Therefore, peracetylated *N*-acetyl- β -D-glucosamine **2** was converted to partially protected *N*-acetyl- β -D-glucosamine derivative **3**. We further functionalized 1-O-methyl- β -D-galactopyranoside **4** regioselectively at the 3'-OH group with Bu₂SnO and propargyl bromide via intermediate tin complexation,^[19] and activated the galactosyl derivative with trichloroacetoneitrile to yield galactosyl donor **5**. We arrived at compound **1** by a glycosylation reaction of **3** and **5** and subsequent deprotection steps (Scheme 1).

We probed the impact of the synthetic extensions on the binding mode by X-ray crystallography. The complex of galectin-1 and LacNAc mimic **1** (Figure 1b, PDB ID: 4Q27) shows a conserved geometry. The structural superposition of the sugars (Supplementary Information, Figure 1S) demonstrates that the galactosyl moiety is captured in the protein depression, as anticipated. Key interactions are formed between His44, Arg48, Asn46 and the axial 4-OH, and between Glu71, Asn61 and the 6-OH of the carbohydrate ligand (Figure 1S). Trp68 supports this ligand orientation by π -CH-stacking with the galactosyl ring. The allyl group of the GlcNAc unit shows weak electron density, suggesting that this extension might be flexible and does not disturb the carbohydrate interactions. This observation will become relevant for galectin-1 recognition when exploiting the alkene functionality for further modification in future experiments. In contrast, the alkyne modification at 3'-OH features well-defined electron density within a distance of 3.7 Å to Trp68, which might be explained by weak T-shaped π - π -interactions.^[20] Surface plasmon resonance (SPR) studies suggested that the synthetic modifications apparently do not interfere with the natural ligand interactions and resulted in comparable dissociation constants for both disaccharides (LacNAc: $91 \pm 5 \mu\text{M}$; **1**: $107 \pm 9 \mu\text{M}$). This K_d value obtained for LacNAc is in agreement with our ITC data and a previously reported value.^[10a]

The next goal of our design strategy was to extend the ligands across the galectin-1 CRD surface region (blue) to the adjacent cavity (green) and eventually address a new surface patch (red, Figure 1). To this end, we computationally generated a library of 397 derivatives from readily available azides and halogenide precursors, and ranked them according to the heavy-atom-corrected ChemPLP score obtained by ligand docking into the galectin-1 binding site (Supplementary Information). We chemically synthesized the ten top-ranking compounds (Scheme 1 and 2). SPR experiments of the novel ligands identified three click products with increased affinity, while the

remaining compounds showed equivalent or improved affinity as LacNAc.

To elucidate their binding mode, we co-crystallized galectin-1 with the new inhibitors **6a** (PDB ID: 4Q1P), **6b** (PDB ID: 4Q1R) and **6d** (PDB ID: 4Q2F). The strong electron density of **6a** and **6b** underlines the conserved binding mode of the carbohydrate unit and its synthetic expansion towards the sub-pocket. (green patch Figure 1 and S1). The triazole fulfills its role as a linker unit and is fixated by a water-mediated interaction with Asn33 in all structures. The benzyl group of **6a** is positioned in a parallel orientation to the carboxyl group of Asp123 by π - π stacking and covers the hydrophobic area formed by Val31. In compound **6b**, the aromatic hydroxyl group undergoes hydrogen bonding with Asp125, while the acetophenone moiety is orientated for CH- π interaction with Gly124, covering the hydrophobic surface area formed by Val31.

Based on these results, we selected ligand **6a** for further elongation due to its favorable orientation on the protein surface, and identified feasible interactions that could be addressed by such an extended ligand. We focused on the introduction of a basic functionality to the ligand to address a carboxyl-rich region of the protein, which to our knowledge has not been targeted before. This part of the target pocket of galectin-1 hosts two aspartate residues (Asp26, Asp125, Figure 1c, red surface area). The main challenge we encountered was to devise a synthetic route that enables the integration of the cationic group while preserving the binding scaffold of **6a**. We extrapolated the structure and hypothesized that an alkyne connector attached to the benzyl ring in *meta*-position should be rigid enough to bridge the gap of about 5.9 Å to Asp125, and a Sonogashira coupling would allow the synthetic realization of the proposed structure.^[21] Therefore, Boc-protected propargylamine was reacted with 3-iodobenzyl alcohol **7** and Bis(triphenylphosphine) palladium(II)dichloride as catalyst in 93 % yield (Scheme 1).^[22] The subsequent steps included conversion of the alcohol into the azide **8** via the bromide compound and deprotection of the amine group, which afterwards was reacted with *N,N'*-Di-Bocthiourea to form the analogue guanidino compound **9**.^[23] Triazole formation of **8** and **9** with precursor **1** and consecutive deprotection of the amine and guanidino functionality, respectively, lead to the expanded ligands **10** and **11** (Scheme 1). We also succeeded in co-crystallizing compounds **10** (Figure 1d) and **11** (Figure 2), with the latter showing a good fit to the electron density (Figure 2b). The amino group reaches the targeted carboxylate of Asp125 and locates within a distance of 2.8 Å. The common structural elements of **6a** and the previous-generation ligand are well aligned and the alkyne extension fulfills its function as a rigid spacer, as demonstrated by the galectin-1-ligand complex. The angle between the LacNAc and the alkyne residue is about 115° (Figure 1d, 2a), which is unique for ligand-protein interactions. To further evaluate these results, we performed ITC experiments, obtaining similar ΔG values for ligand **10** and the previously probed structures **6a-d** and LacNAc respectively (Figure 3). This result is in agreement with the SPR data (Scheme 2).

In conclusion, we investigated and extended the binding modes of a growing carbohydrate ligand fragment to the tumor-associated protein galectin-1. Related combinatorial approaches have concentrated on the evaluation of ligands with a focus on their binding affinity. Here, we concentrated on the design of

COMMUNICATION

innovative ligands with a conserved orientation supported by X-ray crystallography. The alkyne-alkene modified LacNAc mimetic **1** proved to be a useful template as it allows the extension over the carbohydrate recognition site to unexplored sub-pockets in a predictable way. The alkene moiety did not interfere with sugar binding and could be used in future experiments for covalent conjugation with other compounds, e.g. drugs or biomarkers. Furthermore, it could serve as a tool to establish multivalency to compensate for the moderate affinity.

Acknowledgements

We thank the team of beamline ID14-4 and ID 21-1 of ESRF Grenoble, France for their excellent support during data collection.

Keywords: galectin-1 • X-ray crystallography • carbohydrate-lectin interaction • ligand design • fragment-based design


- [1] a) S. H. Barondes, D. N. W. Cooper, M. A. Gitt, H. Leffler, *J. Biol. Chem.* **1994**, *269*, 20807-20810; b) H. Leffler, S. Carlsson, M. Hedlund, Y. Qian, F. Poirier, *Glycoconjugate J.* **2004**, *19*, 433-440.
- [2] a) F. T. Liu, G. A. Rabinovich, *Nat. Rev. Cancer* **2005**, *5*, 29-41; b) G. A. Rabinovich, M. A. Toscano, *Nat. Rev. Immunol.* **2009**, *9*, 338-352.
- [3] a) N. L. Perillo, K. E. Pace, J. J. Seilhamer, L. G. Baum, *Nature* **1995**, *378*, 736-739; b) N. Rubinstein, M. Alvarez, N. W. Zwimer, M. A. Toscano, J. M. Illarregui, A. Bravo, J. Mordoh, L. Fainboim, O. L. Podhajcer, G. A. Rabinovich, *Cancer Cell* **2004**, *5*, 241-251.
- [4] S. Punt, V. L. Thijssen, J. Vrolijk, C. D. de Kroon, A. Gorter, E. S. Jordanova, *PLoS One* **2015**, *10*, e0129119.
- [5] a) L. Astorgues-Xerri, M. E. Riveiro, A. Tijeras-Raballand, M. Serova, C. Neuzillet, S. Albert, E. Raymond, S. Faivre, *Cancer Treat. Rev.* **2014**, *40*, 307-319; b) K. Smetana Jr., S. André, H. Kaltner, J. Kopitz, H.-J. Gabius, *Expert Opin. Ther. Targets* **2013**, *17*, 379-392; c) M. Salatino, D. O. Croci, G. A. Bianco, J. M. Illarregui, M. A. Toscano, G. A. Rabinovich, *Expert Opin. Biol. Ther.* **2008**, *8*, 45-57; d) D. J. Laderach, L. D. Gentilini, L. Giribaldi, V. C. Delgado, L. Nugnes, D. O. Croci, N. Al Nakouzi, P. Sacca, G. Casas, O. Mazza, M. A. Shipp, E. Vazquez, A. Chauchereau, J. L. Kutok, S. J. Rodig, M. T. Elola, D. Compagno, G. A. Rabinovich, *Cancer Res.* **2013**, *73*, 86-96; e) G. A. Rabinovich, *Br. J. Cancer* **2005**, *92*, 1188-1192.
- [6] a) S. Saussez, F. Lorfèvre, T. Lequeux, G. Laurent, G. Chantrain, F. Vertongen, G. Toubreau, C. Decaestecker, R. Kiss, *Oral Oncol.* **2008**, *44*, 86-93; b) M. Watanabe, I. Takemasa, N. Kaneko, Y. Yokoyama, E.-I. Matsuo, S. Iwasa, M. Mori, N. Matsuura, M. Monden, O. Nishimura, *Oncol. Rep.* **2011**, *25*, 1217-1226.
- [7] a) S. Böcker, D. Laaf, L. Elling, *Biomolecules* **2015**, *5*, 1671; b) S. Zhang, R.-O. Moussodia, C. Murzeau, H.-J. Sun, M. L. Klein, S. Vertesy, S. Andre, R. Roy, H.-J. Gabius, V. Percec, *Angew. Chem. Int. Ed.* **2015**, *54*, 4036-4040; c) J. M. Cousin, M. J. Cloninger, *Beilstein J. Org. Chem.* **2015**, *11*, 739-747.
- [8] a) H. van Hattum, H. M. Branderhorst, E. E. Moret, U. J. Nilsson, H. Leffler, R. J. Pieters, *J. Med. Chem.* **2013**, *56*, 1350-1354; b) D. Giguère, S. André, M.-A. Bonin, M.-A. Bellefleur, A. Provencal, P. Cloutier, B. Pucci, R. Roy, H.-J. Gabius, *Bioorg. Med. Chem.* **2011**, *19*, 3280-3287.
- [9] M. G. Ford, T. Weimar, T. Köhli, R. J. Woods, *Proteins: Struct., Funct., Bioinf.* **2003**, *53*, 229-240.
- [10] a) M. F. López-Lucendo, D. Solís, S. André, J. Hirabayashi, K. Kasai, H. Kaltner, H. J. Gabius, A. Romero, *J. Mol. Biol.* **2004**, *343*, 957-970; b) F. P. Schwarz, K. D. Puri, R. G. Bhat, A. Suroliia, *J. Biol. Chem.* **1993**, *268*, 7668-7677; c) C. Diehl, O. Engström, T. Delaine, M. Håkansson, S. Genheden, K. Modig, H. Leffler, U. Ryde, U. J. Nilsson, M. Akke, *J. Am. Chem. Soc.* **2010**, *132*, 14577-14589; d) H. Lis, N. Sharon, *Chem. Rev.* **1998**, *98*, 637-674.
- [11] a) D. Andres, U. Gohlke, N. K. Broeker, S. Schulze, W. Rabsch, U. Heinemann, S. Barbirz, R. Seckler, *Glycobiology* **2013**, *23*, 486-494; b) K. Saraboji, M. Håkansson, S. Genheden, C. Diehl, J. Qvist, U. Weininger, U. J. Nilsson, H. Leffler, U. Ryde, M. Akke, D. T. Logan, *Biochemistry* **2012**, *51*, 296-306.
- [12] D. I. Liao, G. Kapadia, H. Ahmed, G. R. Vasta, O. Herzberg, *Proc. Natl. Acad. Sci. U.S.A.* **1994**, *91*, 1428-1432.
- [13] a) P. Sörme, P. Arnoux, B. Kahl-Knutsson, H. Leffler, J. M. Rini, U. J. Nilsson, *J. Am. Chem. Soc.* **2005**, *127*, 1737-1743; b) P. Sörme, Y. N. Qian, P. G. Nyholm, H. Leffler, U. J. Nilsson, *ChemBioChem* **2002**, *3*, 183-189; c) M. F. Marchiori, D. E. Pires Souto, L. Oliveira Bortot, J. Francisco Pereira, L. T. Kubota, R. D. Cummings, M. Dias-Baruffi, I. Carvalho, V. L. Campo, *Bioorg. Med. Chem.* **2015**, *23*, 3414-3425; d) T.-J. Hsieh, H.-Y. Lin, Z. Tu, T.-C. Lin, S.-C. Wu, Y.-Y. Tseng, F.-T. Liu, S.-T. D. Hsu, C.-H. Lin, *Sci. Rep.* **2016**, *6*, 29457.
- [14] L. V. Lee, M. L. Mitchell, S. J. Huang, V. V. Fokin, K. B. Sharpless, C. H. Wong, *J. Am. Chem. Soc.* **2003**, *125*, 9588-9589.
- [15] a) V. V. Rostovtsev, L. G. Green, V. V. Fokin, K. B. Sharpless, *Angew. Chem. Int. Ed.* **2002**, *41*, 2596-2599; b) C. W. Tornøe, C. Christensen, M. Meldal, *J. Org. Chem.* **2002**, *67*, 3057-3064.
- [16] a) T. Posner, *Ber. Dtsch. Chem. Ges.* **1905**, *57*, 646-657; b) C. E. Hoyle, C. N. Bowman, *Angew. Chem. Int. Ed.* **2010**, *49*, 1540-1573.
- [17] a) A. Fürstner, *Angew. Chem. Int. Ed.* **2000**, *39*, 3012-3043; b) R. H. Grubbs, S. Chang, *Tetrahedron* **1998**, *54*, 4413-4450.
- [18] A.-C. Knall, C. Slugovc, *Chem. Soc. Rev.* **2013**, *42*, 5131-5142.
- [19] S. David, S. Hanessian, *Tetrahedron* **1985**, *41*, 643-663.
- [20] a) C. A. Hunter, J. K. M. Sanders, *J. Am. Chem. Soc.* **1990**, *112*, 5525-5534; b) C. Janiak, *J. Chem. Soc. Dalton Trans.* **2000**, 3885-3896.
- [21] K. Sonogashira, Y. Tohda, N. Hagihara, *Tetrahedron Lett.* **1975**, *16*, 4467-4470.
- [22] T. Ishida, R. Kobayashi, T. Yamada, *Org. Lett.* **2014**, *16*, 2430-2433.
- [23] Y. F. Yong, J. A. Kowalski, M. A. Lipton, *J. Org. Chem.* **1997**, *62*, 1540-1542.

COMMUNICATION

Entry for the Table of Contents

COMMUNICATION

Expansion of natural binding scaffold
N-acetylglucosamine (LacNAc), via a
synthetic multifunctional LacNAc
mimetic leads to expanded ligands,
which share a highly conserved
binding mode.



Nadja Bertleff-Zieschang[†], Julian
Bechold[‡], Clemens Grimm,^{*} Michael
Reutlinger, Petra Schneider, Gisbert
Schneider, and Jürgen Seibel^{*}

Page No. – Page No.

**Elucidating the structural space of
galectin-1-ligand interaction**

COMMUNICATION




Figure 1. Crystallographic complexes of galectin-1 with synthetic *N*-acetyllactosamine-based ligands. The carbohydrate binding site is colored in blue, the adjacent binding site in green, and an Asp-rich region in red. a) Galectin-1 with LacNAc (PDB ID: 4Q26, resolution: 1.40 Å). b) Galectin-1 with the multifunctional LacNAc mimic **1** (PDB ID: 4Q27, resolution: 1.27 Å), the alkyne group forms an interaction with W68. c) Galectin-1 with triazole **6a** (PDB ID: 4Q1P, resolution: 1.42 Å), the triazole forms a water mediated interaction with N33, and the benzyl moiety undergoes π - π stacking with D123. d) Galectin-1 with triazole **10** (PDB ID: 5MWX, resolution: 1.29 Å), a salt bridge is introduced between the amino residue of **9** and D125, the angle between the glucoside moiety and the alkyne group is 112°.

COMMUNICATION




Figure 2. a) Complex of galectin-1 with the synthetic *N*-acetylglucosamine-based ligand **11** (PDB ID: 5MWT, resolution: 1.71 Å). a) The carbohydrate part is colored in blue, the adjacent binding site in green, and an Asp-rich region in red. b) Simulated annealing omit map for the bound ligand **11** contoured on level, shown in blue.






Figure 3. Thermodynamic data obtained by ITC for ligands **6a**, **6b**, **6c**, **10**, and LacNac at 298 K. Colored bars give the respective changes in Gibbs free energy ΔG (red), enthalpy ΔH (black) and the entropy term $-T\Delta S$ (blue). Error bars: stddev from model fitting ($n = 1$).

COMMUNICATION



Scheme 1. Synthesis of alkyne precursor **1** and click products **10** and **11**. a) TMSOTf, CH₂Cl₂, 40 °C, 3 d, then allyl alcohol, 40 °C, 3 h, 62 %; b) NaOMe, MeOH, rt, 15 min, 95 %; c) benzaldehyde dimethylacetal, CSA, MeCN, rt, 12 h, quant.; d) acetic anhydride, pyridine, rt, 12 h, 98 %; e) TFA, TFAA, HSiEt₃, CH₂Cl₂, 0 °C to rt, 5 h, 85 %; f) Bu₂SnO, MeOH, reflux, 4 h, then propargyl bromide, dmf, 35 °C, 40 h, 67 %; g) acetic anhydride, cat. H₂SO₄, rt, 12 h, quant.; h) hydrazine acetate, dmf, 50 °C, 45 min, 69 %; i) trichloroacetonitrile, dbu, CH₂Cl₂, 0 °C, 1 h, 83 %; j) BF₃·OEt₂, CH₂Cl₂, rt, 1 h, 88 %; k) DDQ, CH₂Cl₂, 40 °C, 48 h, then H₂O, rt, 2 h, 83 %; l) NaOMe, MeOH, rt, 12 h, 87 %; m) CuSO₄, sodium ascorbate, ^tBuOH/H₂O 2:1, 36-99 %; n) *tert*-butyl-prop-2-in-1-yl-carbamate, CuI, (PPh₃)₂PdCl₂, thf/NEt₃, rt, 16 h, 93 %; o) CBr₄, PPh₃, DCM, 0 °C to rt, 16 h, 68 %; p) NaN₃, H₂O, 110 °C MW, 1 h, 44 %; q) TFA/H₂O, rt, 30 min, 99 %; r) TFA/DCM, rt, 1 h, quant.; s) *N,N*-di-bocthiourea, 2-chloro-1-methylpyridinium iodide, NEt₃, DMF, rt, 2 h.



Scheme 2. Chemical structures of computationally prioritized click products **6a-6j**. K_d values (in μM) were determined by SPR with immobilized galectin-1 in ^[a]triplicates or ^[c]duplicates. ^[b]K_d values were estimated from a single concentration screen (at 10⁻⁴ M) and measured in duplicates.