Efficient Synthesis of Ningalin C

Anucha Namsa-aid[†] and Somsak Ruchirawat^{*,†,‡,§}

Department of Chemistry, Faculty of Science, Mahidol University, Rama 6 Road, Bangkok 10400, Thailand, Laboratory of Medicinal Chemistry, Chulabhorn Research Institute, Vipavadee Rangsit Highway, Bangkok 10210, Thailand, and Program on Research and Development of Synthetic Drugs, Institute of Science and Technology for Research and Development, Mahidol University, Salaya Campus, Thailand.

somsak@tubtim.cri.or.th

Received April 24, 2002 (Revised Manuscript Received July 10, 2002)

Vol. 4, No. 16 2633-2635

ABSTRACI



A concise and efficient synthesis of the permethyl derivative of the marine alkaloid ningalin C (2) has been accomplished. The key step involves the formation of a pyrrolinone from an aminoquinone in one pot. An efficient route for the synthesis of the key aminoquinone has also been developed.

In 1997, Kang and Fenical¹ reported the isolation of four novel aromatic alkaloids, ningalin A-D (Figure 1), from an



Figure 1. Structures of ningalin A, B, C, and D.

unidentified ascidian of the genus Didemnum² collected in ascidia-rich habitats near Ningaloo Reef region at the northwest cape of western Australia. These ningalin derivatives, as well as lukianols, polycitrins, and lamellarins, appeared to be derived from the condensation of 3,4dihydroxyphenylalanine (DOPA) in the biosynthetic pathwav.3

The first total syntheses of ningalin A and B were reported by Boger et al. in 1999, utilizing the application of the versatile heteroaromatic azadiene Diels-Alder reaction.⁴ Steglich has recently reported an efficient synthesis of ningalin C using an intramolecular Friedel-Crafts acylation.5

In this paper, we report a new and efficient method for the synthesis of ningalin C as a continuation of our synthetic

[†] Mahidol University.

[‡] Chulabhorn Research Institute.

[§] Mahidol University, Salaya Campus.

⁽¹⁾ Kang, H.; Fenical, W. J. Org. Chem. 1997, 62, 3254.

^{(2) (}a) Lindquist, N.; Fenical, W.; Van Duyne, G. D.; Clady, J. J. Org. Chem. 1988, 53, 4570. (b) Carrol, A. R.; Bowden, B. F.; Coll, J. C. Aust. J. Chem. 1993, 46, 489. (c) Urban, S.; Hobbs, L.; Hooper, J. N. A.; Capon, R. J. Aust. J. Chem. 1996, 49, 711. (d) Urban, S.; Capon, R. J. Aust. J. Chem. 1995, 48, 1491.

⁽³⁾ For a review see: Bowden, B. F. In Studies in Natural Products Chemistry; Rahman, A., Ed.; Elsevier Science: New York, 2000; Vol. 23, pp 233-283.

^{(4) (}a) Boger, D. L.; Boyce, C. W.; Labroli, M. A.; Sehon, C. A.; Jin, Q. J. Am. Chem. Soc. 1999, 121, 54. (b) Boger, D. L.; Soenen, D. R.; Boyce,

C. W.; Hedrick, M. P., Jin, Q. J. Org. Chem. 2000, 65, 2479.
(5) Peschko, C.; Steglich, W. Tetrahedron Lett. 2000, 41, 9477.

program on lamellarins and other bioactive pyrrole alkaloids.⁶ Our retrosynthetic analysis of this compound is outlined in Scheme 1.



It was planned that the pyrrolinone system of the target ningalin compound 2 could be synthesized from naphthoquinone 1 via addition of the enolate derived from methyl homoveratrate and subsequent lactam bond formation. This aminoquinone could be synthesized by addition of an amine to quinone 3 derived from the oxidation of naphthol derivative 4. It was expected that an allyl carbanion prepared from 5 would undergo intramolecular acylation to give compound 4. The alkene 5 could conceivably be prepared by the reaction of eugenol methyl ether 6 with bromo compound 7 via the Heck reaction.⁷

In practice, the palladium-catalyzed coupling reaction⁸ between the methyl 2-bromoveratrate 7 and commercially available eugenol methyl ether 6 gave ester 5 in 65% yield. (Scheme 2).

Further cyclization⁹ of methyl ester **5** was accomplished by treatment with 2 equiv of lithium diisopropylamide in

(8) Amorese, A.; Arcadi, A.; Bernocchi, E.; Cacchi, S.; Cerini, S.; Fedili, W.; Ortar, G. *Tetrahedron* **1989**, *45*, 813.



^{*a*} Reagents and conditions: (a) Pd(PPh₃)₄, NaHCO₃, DMF, reflux, 24 h, 65%; (b) 2 equiv of LDA, THF, -78 °C, 2 h, then rt 2 h, 76%; (c) H₂O₂, I₂, H₂SO₄, MeOH, 62%; (d) homoveratrylamine (**9**), EtOH, rt, 24 h, 80%

THF at -78 °C to give naphthol **4** (76%), which has recently been obtained by a different route.¹⁰

Oxidation¹¹ of naphthol **4** with 30% hydrogen peroxide containing a trace amount of iodine and sulfuric acid in methanol afforded the desired naphthoquinone **3** in 62% yield as a red solid.

On the basis of the synthetic plan proposed in Scheme 1, the key aminoquinone intermediate could be prepared via the nucleophilic addition of amine to naphthoquinone derivative **3**. Indeed, it was found that treatment of naphthoquinone **3** with homoveratrylamine **9** in ethanol¹² at room temperature furnished the homologous amide, aminonaphthoquinone **1**, as a deep red solid in 80% yield (Scheme 2).

Similarly, the aminonaphthoquinone **10** was readily prepared by the addition of homoveratrylamine **9** to commercially available menadione **8** in ethanol at room temperature for 24 h. The aminoquinone derivatives have recently been shown to be versatile intermediates in the synthesis of various natural products.¹³

The construction of the pyrrolinone moiety in the ningalin skeleton was initially investigated using this readily available aminoquinone derivative **10**.

It was reasoned that the presence of the amino group would render the C-1 carbonyl less electrophilic and thus the

⁽⁶⁾ Ruchirawat, S.; Mutarapat, T. Tetrahedron Lett. 2001, 42, 1205.

⁽⁷⁾ For some recent reviews, see: (a) Ikeda, M.; El Bialy, S. A. A.; Yakura, T. *Heterocycles* **1999**, *51*, 1957. (b) Amatore, C.; Jutand, A. *Acc. Chem. Res.* **2000**, *33*, 314. (c) Beletskaya, I. P.; Cheprakov, A. V. *Chem. Rev.* **2000**, *100*, 3009. (d) Poli, G.; Giambastiani, G.; Heumann, A. *Tetrahedron* **2000**, *56*, 5959.

^{(9) (}a) Sibi, M. P.; Dankwardt, J. W.; Snieckus, V. J. Org. Chem. **1986**, 51, 273. (b) de Koning, C. B.; Michael, J. P.; Rosseau, A. L. Tetrahedron Lett. **1997**, 38, 893. (c) Hattori, T.; Takeda, A.; Suzuki, K.; Koike, N.; Koshiishi, E.; Miyano, S. J. Chem. Soc., Perkin Trans. 1 **1998**, 3661.

⁽¹⁰⁾ Estevez, R. J.; Martinez, E.; Martinez, L.; Treus, M. *Tetrahedron* **2000**, *56*, 6023.

⁽¹¹⁾ Minisci, F.; Citterio, A.; Vsaimara, E.; Fontana, F.; De Bernardinis, S. *J. Org. Chem.* **1989**, *54*, 728.

^{(12) (}a) Barret, R.; Roue, N. *Tetrahedron Lett.* **1999**, *40*, 3889. (b) Tohma, H.; Harayama, Y.; Hashizumi, M.; Iwata, M.; Egi, M.; Kita, Y. *Angew. Chem., Int. Ed.* **2002**, *41*, 348.

⁽¹³⁾ Nicolaou, K. C.; Sugita, K.; Baran, P. S.; Zhong, Y.-L. Angew. Chem., Int. Ed. 2001, 40, 207 and references therein.

nucleophile would preferentially attack C-4 carbonyl group.¹⁴ This prediction was indeed found to be the case. When aminonaphthoquinone **10** was treated with the carbanion derived from the reaction of methyl homoveratrate and 2 equiv of lithium diisopropylamide in tetrahydrofuran at -78 °C, the desired ningalin C skeleton **11** was obtained (69%) as an orange solid together with hydrated product **12**, isolated as a white solid in 9% yield (Scheme 3). It was gratifying



^{*a*} Reagents and conditions: (a) EtOH, rt, 24 h, 82%; (b) 2 equiv of LDA, THF, -78 °C; (c) 1 equiv of methyl homoveratrate, THF, -78 °C, 2 h then rt 2 h, 69%(**11**), 9%(**12**); (d) 2 M HCl, CH₂Cl₂, rt, 5 h

to see that not only was the addition of the anionic species to the right ketone group of compound **10** but also the lactam cyclization took place to generate the ultimate pyrrolinone system in one pot. We were unable to carry out the *intermolecular* amide bond formation by the reaction of aminonaphthoquinone **10** with homoveratroyl chloride in the presence of various bases (Na₂CO₃, K₂CO₃, NaH, Et₃N). This is presumably due to the fact that the lone pair of electrons on nitrogen is not readily available because the nitrogen is part of a vinylogous amide. It was thus concluded that in the key formation of the pyrrolinone the formation of the carbon–carbon bond preceded the *intramolecular* amide bond formation.

With the starting material 1 and an efficient method for the synthesis of pyrrolinones in hand, the synthesis of permethyl ningalin C 2 was then investigated. The reaction of aminoquinone 1 with methyl homoveratrate and 2 equiv of LDA gave the required permethyl ningalin C 2 (73%) and the hydrated compound 13 (15%) in a straightforward manner (Scheme 4). The hydrated compounds 12 and 13



^{*a*} Reagents and conditions: (a) 2 equiv of LDA, THF, -78 °C; (b) 1 equiv of methyl homoveratrate, THF, -78 °C, 2 h then rt 2 h, 73% (2), 15%(13); (d) 2 M HCl, CH₂Cl₂, rt, 5 h.

could be readily dehydrated by treatment with 2 M hydrochloric acid in dichloromethane at room temperature for 5 h to give the ningalin compound **11** and permethyl ningalin C **2** in quantitative yield. Permethyl ningalin C **2** was demethylated⁵ by BBr₃ in methylene chloride to give ningalin C in 72% yield. The spectral data (IR, UV, ¹H NMR, ¹³C NMR, MS) of our synthetic ningalin C are in full agreement with the published data for the natural product.^{1,5}

In conclusion, our synthetic approach to ningalin C demonstrates the utility of the one-pot reaction for the preparation of the pyrrolinone system, the core moiety of ningalin C. The synthesis of the quinone and aminoquinone also shows great promise for application in the synthesis of related quinone derivatives.

Acknowledgment. We gratefully thank the Thailand Research Fund (TRF) for generous financial support and the award of Senior Research Scholar to S.R. and the Golden Jubilee Scholarship to A.N. We acknowledge the facilities in the Department of Chemistry, Mahidol University, provided by the Postgraduate Education and Research in Chemistry (PERCH) program. We are grateful to Professor W. Steglich for copies of NMR spectra of permethyl ningalin C and ningalin C and the procedure for the demethylation.

Supporting Information Available: Detailed experimental procedures and characterization data for new compounds and spectra (IR, UV, ¹H NMR, ¹³C NMR, MS) of the synthetic ningalin C. This material is available free of charge via the Internet at http://pubs.acs.org.

OL026074S

⁽¹⁴⁾ Knolker, H.-J.; Frohner, W.; Reddy, K. R. Synthesis 2002, 557.