

Evident and latent plasticity across the rice diterpene synthase family with potential implications for the evolution of diterpenoid metabolism in the cereals

Dana MORRONE, Matthew L. HILLWIG, Matthew E. MEAD, Luke LOWRY, D. Bruce FULTON and Reuben J. PETERS¹

Department of Biochemistry, Biophysics and Molecular Biology, Iowa State University, Ames, IA 50011, U.S.A.

The evolution of natural product biosynthetic pathways can be envisioned to occur via a number of mechanisms. In the present study we provide evidence that latent plasticity plays a role in such metabolic evolution. In particular, rice (*Oryza sativa*) produces both *ent*- and *syn*-CPP (copalyl diphosphate), which are substrates for downstream diterpene synthases. In the present paper we report that several members of this enzymatic family exhibit dual reactivity with some pairing of *ent*-, *syn*- or normal CPP stereochemistry. Evident plasticity was observed, as a previously reported *ent*-sandaracopimaradiene synthase also converts *syn*-CPP into *syn*-labda-8(17),12*E*,14-triene, which can be found

in planta. Notably, normal CPP is not naturally found in rice. Thus the presence of diterpene synthases that react with this non-native metabolite reveals latent enzymatic/metabolic plasticity, providing biochemical capacity for utilization of such a novel substrate (i.e. normal CPP) which may arise during evolution, the implications of which are discussed.

Key words: metabolic evolution, phytoalexin, substrate specificity, terpene synthase, terpenoid natural product.

INTRODUCTION

Terpenoids, with over 40 000 known individual compounds, form the largest class of natural products [1]. Prominent among these are the ~7000 labdane-related diterpenoids, which are particularly prevalent in plants, where the affiliated gibberellin phytohormone production required for normal growth and development provides a genetic reservoir that has been repeatedly drawn upon for the evolution of such novel natural products [2,3]. The unifying biogenetic feature of labdane-related diterpenoid biosynthesis is its initiation by a sequential pair of cyclization and/or rearrangement reactions, and the corresponding enzymes are often found as functionally diverse gene families in plants [4,5]. The characteristic decalin ring structure is formed from the universal diterpenoid precursor GGPP [(*E,E,E*)-geranylgeranyl diphosphate] by initially acting class II diterpene cyclases, which most commonly produce a labdadienyl/CPP (copalyl diphosphate) of normal (9*S*,10*S*), *syn* (9*S*,10*R*) or *ent* (9*R*,10*R*) configuration (Scheme 1), with the corresponding enzymes termed CPSs (CPP synthases). The resulting CPP is subsequently utilized by class I diterpene synthases, which lyse the allylic diphosphate ester linkage to initiate carbocationic cyclization and/or rearrangement reactions. Because the corresponding enzyme in gibberellin biosynthesis is KS (*ent*-kaurene synthase), the derived class I diterpene synthases are often termed KSL (KS-like). However, although these enzymes do not react with GGPP and are typically assumed to be specific for particular stereoisomers of CPP, comprehensive investigation of such substrate stereospecificity (e.g. with all three commonly found CPP stereoisomers) has not been reported.

Rice (*Oryza sativa*) has become a model system for investigation of labdane-related diterpenoid biosynthesis, as it

has long been appreciated that it produces at least 20 such natural products as antimicrobial phytoalexins, and relatively early sequencing of the rice genome has led to functional characterization of the corresponding diterpene synthase gene families [4,6]. However, since rice produces only *ent*- and *syn*-CPP, as first suggested by substrate studies with cell-free enzyme assays [7,8], and then more conclusively demonstrated by functional genomics-based investigations of the rice CPS family [9–11], the rice KSL (OsKSL) was only assayed with these two stereoisomers [12–18]. This OsKSL family, with eight functional members (Scheme 2), provided an opportunity to comprehensively explore diterpene synthase substrate stereospecificity not only for individual enzymes, but also across an entire organism/genome, providing insight into the encoded metabolic plasticity. Accordingly, we have utilized our modular metabolic engineering system [19] to investigate the CPP substrate stereospecificity of the OsKSL family. The results of this substrate specificity screen revealed not only evident, but also latent, metabolic plasticity, as well as offering some insight into diterpene synthase substrate recognition, all of which are discussed below.


EXPERIMENTAL

General

Unless otherwise noted, chemicals were purchased from Fisher Scientific, and molecular biology reagents were from Invitrogen. All recombinant expression was carried out using the OverExpress C41 strain of *Escherichia coli* (Lucigen), which is well-suited to T7 promoter-based expression of plant-derived labdane-related diterpene synthases [20]. GC-MS was performed with a Varian

Abbreviations used: CPP, copalyl diphosphate; CPS, CPP synthase; CYP, cytochrome P450 mono-oxygenase; DQF-COSY, double-quantum-filtered correlation spectroscopy; FID, flame ionization detection; GGPP, (*E,E,E*)-geranylgeranyl diphosphate; GST, glutathione transferase; HMBC, heteronuclear multiple bond correlation; HSQC, heteronuclear single-quantum coherence; KS, *ent*-kaurene synthase; KSL, KS-like; MEP, methyl erythritol phosphate; OsKSL, rice (*Oryza sativa*) KSL.

¹ To whom correspondence should be addressed (email rjpeters@iastate.edu).



Scheme 1 Cyclization of GGPP to stereoisomeric CPP by CPS

Note that the position of the C5 hydrogen is always anti to the C10 methyl group due to the chair-chair conformation of the GGPP in this initial CPS-catalysed (i.e. class II) cyclization mechanism.

3900 GC with Saturn 2100 ion-trap mass spectrometer in electron ionization (70 eV) mode using a HP-5ms column. Samples (1 μ l) were injected in splitless mode at 50°C and, after holding for 3 min at 50°C, the oven temperature (unless otherwise noted) was raised at a rate of 14°C/min to 300°C, where it was held for an additional 3 min. MS data from 90 to 600 m/z were collected starting 12 min after injection until the end of the run.

Functional characterization via metabolic engineering


Functional characterization of the substrate stereospecificity of the OsKLS was accomplished by use of our previously described modular metabolic engineering system [19]. Briefly, using this system it is possible to co-express class I labdane-related diterpene synthases such as the OsKSL from pET-based pDEST expression vectors with GGPP and CPS carried on a compatible pACYC-Duet (Novagen/EMD)-derived plasmid, where pGGnC carries the D621A mutant of abietadiene synthase from *Abies grandis* that produces CPP of normal stereochemistry [21], while pGGeC carries the An2/ZmCPS2 *ent*-CPS from *Zea mays* [22], and pGGsC carries the OsCPS4 *syn*-CPS from rice [10]. Accordingly, C41 cells were transformed with the appropriate plasmid enabling functional recombinant expression of each OsKSL, as described previously [18], along with one of the three pGGxC vectors described above, in all possible pairings. These recombinant bacteria were then grown in 20 ml of liquid media cultures with shaking at 37°C to $A_{600} \sim 1.0$,

then shifted to 16°C for 1 h prior to induction with 0.5 mM IPTG (isopropyl β -D-thiogalactopyranoside), with the cultures then fermented for an additional 72 h at 16°C with continued shaking. The cultures were then extracted with an equal volume of hexane, with 1:50 (v/v) ethanol added to disrupt emulsions. Diterpene production was quantified by GC-FID (flame ionization detection) analysis, with GC-MS comparison with authentic standards enabling identification of all resulting compounds but that from co-expression of OsKSL10 with pGGsC (i.e. *syn*-CPP).

To directly determine the structure of the OsKSL10 product from *syn*-CPP, we utilized previously described methodology to improve production levels in our metabolic engineering system [23], which enabled facile generation of the milligram quantities needed for structural characterization by NMR spectroscopy. Specifically, we co-transformed pDEST15/OsKSL10 and pGGsC with the further co-compatible pIRS plasmid that increases flux into the endogenous MEP (methyl erythritol phosphate) pathway by overexpression of the *idi*, *dxs*, and *dxr* genes from *E. coli*. The resulting recombinant bacteria were grown and extracted as described above, but as a 1 litre culture supplemented with the MEP pathway precursors pyruvate (50 mM) and 2% (v/v) glycerol. The resulting 1 litre of hexane extract was reduced to ~ 10 ml by rotary evaporation, passed over a ~ 5 ml silica-gel column to remove any polar contaminants, and then dried under nitrogen gas, yielding ~ 15 mg of the unknown diterpene, which was then re-dissolved in 0.4 ml of C_2HCl_3 for NMR analysis.

Chemical structure identification

NMR spectra for the unknown *syn*-CPP-fed OsKSL10 product were recorded at 25°C on a Bruker Avance 500 spectrometer equipped with a cryogenic probe for 1H and ^{13}C . Structural analysis was performed using 1H , one-dimensional, DQF-COSY (double-quantum-filtered correlation spectroscopy), HSQC (heteronuclear single-quantum coherence), HMQC (heteronuclear single quantum correlation), HMBC (heteronuclear multiple bond correlation) and NOESY (nuclear Overhauser enhancement spectroscopy) experiment spectra acquired at 500 MHz, and ^{13}C and DEPT135 spectra (125.5 MHz) using standard experiments from the Bruker TopSpin v1.3 software. Chemical shifts were referenced to the known



Scheme 2 Rice diterpene metabolic map [17,18]

Shown are the reactions catalysed by the various rice diterpene cyclases [both class II CPS, and class I KS(L)].

Table 1 ^1H and ^{13}C NMR assignments for *syn*-labda-8(17),12*E*,14-triene

C	δ_{C} (p.p.m.)	δ_{H} (p.p.m.), multiplicity	J (Hz)
1	37.16	1.086 (m), 1.550 (m)	
2	19.39	1.434 (m), 1.607 (m)	
3	42.92	1.167 (dt), 1.403 (m)	3.6, 13.1
4	33.48		
5	46.04	1.298 (m)	
6	23.85	1.296 (m), 1.611 (m)	
7	31.88	2.065 (m), 2.146 (m)	
8	149.23		
9	58.42	1.635 (m)	
10	38.25		
11	25.89	2.164 (m), 2.409 (dt)	15.0, 5.6
12	133.36	5.365 (t)	7.1
13	133.68		
14	142.01	6.330 (dd)	10.7, 17.4
15	110.14	4.864 (d), 5.020 (d)	10.7, 17.4
16	12.12	1.707 (s)	
17	109.72	4.474 (t), 4.640 (t)	1.6, 2.2
18	33.78	0.875 (s)	
19	22.45	0.800 (s)	
20	22.40	0.909 (s)	

chloroform-*d* (^{13}C 77.23, ^1H 7.24 p.p.m.) signals offset from TMS (trimethylsilyl). Correlations from the HMBC spectra were used to propose a partial structure, whereas connections between protonated carbons were obtained from DQF-COSY to complete the partial structure and assign proton chemical shifts. The configuration of the A and B rings is predetermined by the configuration of the CPP enzyme substrate, since chemical bonds in that portion of the molecule are not altered. The configuration of the acyclic portion of the molecule (C11–C16) was determined by reference to spectra for normal labdatrienes, which have been previously characterized [24]. Specifically, the C12–C13 double-bond configuration was determined to be *E* based on the C14 ^{13}C chemical shift for *syn*-labda-8(17),12*E*,14-triene (142.01 p.p.m.). The comparison of *E* and *Z* configuration of normal labda-8(17),12,14-trienes shows a dramatic ~ 10 p.p.m. difference in the C14 chemical shift in *E* compared with *Z* configurations [~ 131 p.p.m. (*Z*) compared with ~ 141 (*E*) p.p.m.]. This reference data and our NMR analysis support the presented *syn*-labda-8(17),12*E*,14-triene double bond configuration for the OsKSL10 product. The assignments for the annotated ^{13}C and ^1H one-dimensional spectra are presented in Table 1.

Enzymatic assays

Assays were performed with the purified GST (glutathione transferase)–OsKSL10 fusion protein using enzymatic coupling, as described previously [25,26]. Briefly, GGPP was completely converted into either *ent*- or *syn*-CPP by incubation with the appropriate CPS for 5 h at room temperature (22°C). Assays were performed in duplicate with various concentrations of GGPP ranging from 0.1 μM to 100 μM , with time and GST–OsKSL10 concentration varied to ensure that measurements were taken in the linear portion of the reaction progress curve. The assays were inactivated by heating the solution to 65°C for 5 min, with the remaining substrate then dephosphorylated by treatment with CIP (calf intestinal phosphatase; New England Biolabs), enabling extraction of both the diterpene product and copalol (derived from dephosphorylation of CPP). This was done to enable correction for handling errors by measurement of the fractional turnover (i.e. comparison of the relative amounts of diterpene product and copalol by GC-FID), which was used in combination with

the known starting concentration of substrate to calculate the corresponding catalytic rate.

Extraction of rice leaves

Rice leaves that had been treated with methyl jasmonate for 36 h, as described previously [10], were frozen at -80°C . Approximately 100 g of these frozen leaves were cut up and then homogenized by grinding in a mortar and pestle under liquid nitrogen in the presence of glass beads. After grinding, the resulting powder was extracted with 75 ml of hexanes, then 75 ml of ethyl acetate, and finally 75 ml of tetrahydrofuran. The combined organic extracts were reduced by rotary evaporation, passed over silica gel, and dried under nitrogen as described above. The resulting material was then resuspended in 1 ml of ethyl acetate, with 20 μl aliquots being removed and then dried under nitrogen, and resuspended in 20 μl of hexane for analysis by GC-MS, using an extended temperature ramp where, after the initial 3 min hold at the 50°C injection temperature, the oven temperature was raised at 18°C/min to 190°C, then slowed to 2.5°C/min until reaching 275°C, following which it was raised at 20°C/min to 300°C, where it was held for 2 min.

RESULTS

Screening OsKSL substrate specificity

To investigate the substrate stereospecificity of the entire OsKSL family, we turned to our previously developed modular metabolic engineering system [19]. In particular, using this system it was possible to co-express each of the OsKSLs with a GGPP synthase and CPS producing one of the three commonly found stereoisomers of CPP (note that all of the OsKSLs have been demonstrated to not react with the acyclic GGPP precursor). Accordingly, the ability of each of the eight active OsKSLs to react with CPP of *ent*-, *syn*- or normal stereochemistry was determined by simply extracting the corresponding recombinant bacterial cultures and analysing the resulting organic extract for the presence of diterpenes by GC-MS. The previously reported activity for each of the OsKSLs, as summarized in Scheme 2, was evident. In addition, while the other OsKSLs were specific for a single stereoisomer of CPP, OsKSL4 and OsKSL11 reacted with CPP of normal stereochemistry along with their previously reported activity with *syn*-CPP, and OsKSL10 was active with *syn*-CPP as well as its previously reported activity with *ent*-CPP. Surprised by this latter result, we purified OsKSL10 and verified its reactivity with *syn*-CPP, with the same product resulting from *in vitro* reactions as found in the context of the bacterial metabolic engineering system. In all three cases, diterpene yield with the 'alternative' CPP stereoisomer approached that found with the 'native' stereoisomer, being limited to less than 3-fold decreases. As yield in our metabolic engineering system has been correlated with catalytic efficiency [25,26], this suggests that these enzymes exhibit reasonable affinity for and activity with their 'alternative' substrates.

Alternative substrate product identification

By comparison with available authentic standards, OsKSL4 activity with CPP of normal stereochemistry was found to produce a mixture of $\sim 90\%$ pimara-8(14),15-diene and $\sim 10\%$ isopimara-7,15-diene, whereas OsKSL11 reacted with the same substrate to yield essentially only isopimara-8(14),15-diene (also known as sandaracopimaradiene; Figure 1). However, it was not




Figure 1 Identification of alternative products by comparison with authentic standards

(A) GC-MS-selected ion chromatogram of OsKSL4 products from normal CPP. (B) Mass spectra of OsKSL4 major product. (C) Mass spectra of authentic pimara-8(14),15-diene. (D) GC-MS-selected ion chromatogram of OsKSL11 products from normal CPP. (E) Mass spectra of OsKSL11 major product. (F) Mass spectra of authentic sandaracopimaradiene.

possible to identify the product of OsKSL10 activity with *syn*-CPP in this fashion, as it did not match any authentic standard on-hand. This diterpene was then identified by increasing flux towards terpenoid production, via up-regulation of the endogenous isoprenoid precursor supply pathway as described previously [23], to enable the corresponding recombinant bacteria to produce enough of the novel product for NMR analysis (~15 mg, following purification from 1 litre of bacterial culture). The structure was assigned by analysis of collected HMBC, HSQC and COSY spectra, and found to be *syn*-labda-8(17),12*E*,14-triene (Table 1), completing the assignment of the observed alternative reactions (Scheme 3).


Physiological relevance of *syn*-labdatriene

Because rice produces both *ent*- and *syn*-CPP, it seemed possible that the dual substrate reactivity of OsKSL10 was physiologically relevant (i.e. both products can be found *in planta*). The labdane-related diterpenoid natural products of rice are largely thought to be anti-microbial phytoalexins [4,6], whose biosynthesis, by definition [27], is induced as part of the plant microbial defence response. This includes transcriptional up-regulation of OsKSL10, as well as OsKSL4 [14]. Accordingly,

the possibility that OsKSL10 produces *syn*-labdatriene *in planta* was examined by phytochemical analysis of methyl-jasmonate-induced and uninduced rice leaves. As previously reported [7], extracts from control (uninduced) rice leaves contained only small amounts of *ent*-kaurene, whereas numerous diterpenes were detected in extracts from induced rice leaves (Figure 2). These specifically included *syn*-labdatriene, as well as *syn*-stemodene, whose presence *in planta* had not been previously demonstrated, although the OsKSL11 responsible seems widely conserved [17]. Steady-state kinetic analysis was also consistent with physiological relevance of the ability of OsKSL10 to react with both *ent*- and *syn*-CPP, as it exhibited similar catalytic efficiency with each (Table 2). Notably, consistent with the lack of a normal CPP producing CPS in the rice genome, whereas the *syn*-CPP-derived products of OsKSL11 (as noted above), as well as OsKSL4, were readily observed, the pimaradienes produced by these diterpene synthases from normal CPP were not evident.

DISCUSSION

On the basis of the genome sequence available [28], the entire family of rice diterpene synthases is known, and subsequent biochemical characterization has assigned unique functional roles



Scheme 3 Alternative reactions catalysed by (A) OsKSL4, (B) OsKSL11 and (C) OsKSL10

Table 2 OsKSL10 steady-state kinetic constants

Substrate	K_m (μM)	k_{cat} (min^{-1})	k_{cat}/K_m ($\text{M}^{-1} \cdot \text{s}^{-1}$)
ent-CPP	0.6 ± 0.1	0.02 ± 0.01	5.6×10^3
syn-CPP	5 ± 1	0.05 ± 0.02	1.7×10^3

to each of these [15,18]. Hence it is possible to define this early part of the labdane-related diterpene metabolic network for this important crop plant (Scheme 2), with the resulting/downstream natural products largely thought to be involved in the microbial defence response of the plant [4,6]. One outcome of the studies reported in the present paper is the addition of the *syn*-labdatriene product of OsKSL10 to this metabolic network. Given the inducible nature of the transcription of *OsKSL10* [14], and the role of its already known *ent*-sandaracopimaradiene product as a precursor for the antifungal phytoalexins oryzalexins A–F, it seems likely that any *syn*-labdatriene-derived natural products will similarly function as antifungal agents, although this remains to be demonstrated. This would be consistent with similar transcriptional induction of all of the other diterpene synthases involved in biosynthesis of the >20 known rice diterpenoid phytoalexins [4,6,9–16]. Nevertheless, one intriguing alternative possibility is a role for *syn*-labdatriene in the production of an inducible signalling molecule, as has been previously indicated for a similar labdatriene-derived diterpenoid in tobacco [29]. On the other hand, the ability of OsKSL4 and OsKSL11 to react with normal CPP is not directly relevant in rice, whose genome does not contain a CPS that produces this stereoisomer [9–11].

In any case, the ability of certain rice diterpene synthases to react with two different stereoisomers of CPP raises the question of how such dual substrate recognition is achieved. OsKSL4 and OsKSL11 react with both *syn*- and normal CPP, and the




Figure 2 Detection of *syn*-labda-8(17), 12E, 14-triene in induced rice plant tissue

(A) GC-MS-selected ion chromatogram of induced rice leaf extract (* indicates the peak corresponding to *syn*-sternodene product of OsKSL11). (B) Mass spectrum of 15.53 min peak in induced rice leaf extract. (C) Mass spectrum of authenticated *syn*-labda-8(17), 12E, 14-triene (retention time = 15.51 min).

obvious structural similarity between these two stereoisomers is the occurrence of co-axial β -methyl substituents on C4 and C10, which provides steric bulk on this ‘upper face’ of the *trans*-decalin nucleus (Figure 3). Presumably, it is this common feature that enables the dual reactivity exhibited by OsKSL4 and OsKSL11. In addition, the stereochemistry of the observed products is consistent with similar conformations of both substrates in each active site (Figure 4). By contrast, the ability of OsKSL10 to react with both *ent*- and *syn*-CPP presumably arises from the common C9 α connection of the isoprenyl-diphosphate-containing ‘tail’ to the substrate nucleus (Figure 3), although the lack of cyclization with *syn*-CPP prevents any insights into the relative conformation of these two substrates within the active site.

Notably, some plasticity in substrate specificity has been previously observed in other terpene synthases, leading to suggestions of evolutionary relevance. Perhaps most strikingly, work in strawberries has demonstrated a clear evolutionary derivation of a linalool/nerolidol synthase in cultivated strawberry from monoterpene synthases found in wild strawberry plants, wherein the loss of the plastid targeting pre-sequence from a




Figure 3 Configurational comparison of CPP stereoisomers




Figure 4 Comparison of active-site conformations of the alternative CPP stereoisomers indicated by catalysed product outcome for (A) OsKSL4 and (B) OsKSL11

presumably plastic monoterpene synthase ancestor gave rise to this novel sesqui-/mono-terpene synthase [30]. Similar mono-/sesqui-terpene synthase plasticity has been uncovered in other cases as well [31].

The ability of OsKSL4 and OsKSL11 to react with normal CPP demonstrates the presence of latent diterpene metabolic plasticity in rice. Given the production of normal CPP by a diverse array of plants, ranging from gymnosperms to

angiosperms, with no obvious homology between the relevant known enzymes from these two broad groups [5,32–34], it seems likely that neofunctionalization of the corresponding class II diterpene cyclases to production of this CPP stereoisomer occurs with some appreciable frequency. Indeed, recent results demonstrate that the closely related monocot cereal crop plant wheat (*Triticum aestivum*) produces normal rather than *syn*-CPP (Y. Wu, T. Toyomasu and R.J. Peters, unpublished work). Accordingly, the latent metabolic plasticity evident in the present study from the substrate promiscuity of these two rice diterpene synthases provides a means by which such emergence of stereochemically differentiated substrates could be immediately incorporated into ‘secondary’ metabolism, which seems pertinent to such biochemical evolution in the cereal plant family. In particular, coupled with the often-observed substrate plasticity of downstream tailoring enzymes, especially CYPs (cytochrome P450 mono-oxygenases), this would then enable emergence of essentially complete biosynthetic pathways (e.g. the bioactive oryzalexins D–F are simply diterpenoid diols only two hydroxylation reactions removed from their diterpene precursor). For example, rice contains ~350 CYPs [35], with the large CYP71 and CYP76 families (74 and 28 members respectively) each containing members involved in labdane-related diterpenoid biosynthesis [36,37]. Notably, at least one of these can hydroxylate pimaradienes of normal stereochemistry (Q. Wang, M.L. Hillwig, S. Swaminathan and R.J. Peters, unpublished work). Thus, provided access to a novel metabolite substrate, the latent plasticity of terpene synthases, such as that shown in the present study, can immediately give rise to potentially bioactive natural products, providing a mechanism for rapid evolution of novel biosynthetic pathways. Although not directly selectable in and of itself, such latent plasticity would be periodically uncovered upon the introduction of novel metabolites, presumably providing a corresponding selectable advantage, albeit in punctuate fashion.

AUTHOR CONTRIBUTION

The biochemical studies described in the present study were carried out by Dana Morrone, with assistance from Matthew Mead and Luke Lowry. NMR analysis of the resulting novel diterpenes were performed by Matthew Hillwig and Bruce Fulton. The manuscript was prepared by Reuben Peters and Dana Morrone.

ACKNOWLEDGEMENTS

The authors thank Professor Robert M. Coates (University of Illinois, Urbana, IL, U.S.A.) for critically reading the manuscript, and Dr Xiaoming Chen (Iowa State University) for preparation of the rice tissues analysed in the present study.

FUNDING

This work was supported by the National Institutes of Health [grant number GM076324 (to R.J.P.)].

REFERENCES

- Buckingham, J. (2002) Dictionary of Natural Products (on-line edition). Chapman & Hall/CRC Press
- Peters, R. J. (2010) Two rings in them all: the labdane-related diterpenoids. *Nat. Prod. Rep.* **27**, 1521–1530
- Keeling, C. I., Dullat, H. K., Yuen, M., Ralph, S. G., Jancsik, S. and Bohlmann, J. (2010) Identification and functional characterization of monofunctional *ent*-copalyl diphosphate and *ent*-kaurene synthases in white spruce reveal different patterns for diterpene synthase evolution for primary and secondary metabolism in gymnosperms. *Plant Physiol.* **152**, 1197–1208

- 4 Peters, R. J. (2006) Uncovering the complex metabolic network underlying diterpenoid phytoalexin biosynthesis in rice and other cereal crop plants. *Phytochemistry* **67**, 2307–2317
- 5 Martin, D. M., Faldt, J. and Bohlmann, J. (2004) Functional characterization of nine norway spruce *TPS* genes and evolution of gymnosperm terpene synthases of the *TPS-d* subfamily. *Plant Physiol.* **135**, 1908–1927
- 6 Toyomasu, T. (2008) Recent advances regarding diterpene cyclase genes in higher plants and fungi. *Biosci. Biotechnol. Biochem.* **72**, 1168–1175
- 7 Wickham, K. A. and West, C. A. (1992) Biosynthesis of rice phytoalexins: identification of putative diterpene hydrocarbon precursors. *Arch. Biochem. Biophys.* **293**, 320–332
- 8 Mohan, R. S., Yee, N. K. N., Coates, R. M., Ren, Y. Y., Stamenkovic, P., Mendez, I. and West, C. A. (1996) Biosynthesis of cyclic diterpene hydrocarbons in rice cell suspensions: conversion of 9,10-*syn*-labda-8(17),13-dienyl diphosphate to 9 β -pimara-7,15-diene and stemar-13-ene. *Arch. Biochem. Biophys.* **330**, 33–47
- 9 Prisic, S., Xu, M., Wilderman, P. R. and Peters, R. J. (2004) Rice contains two disparate *ent*-copalyl diphosphate synthases with distinct metabolic functions. *Plant Physiol.* **136**, 4228–4236
- 10 Xu, M., Hillwig, M. L., Prisic, S., Coates, R. M. and Peters, R. J. (2004) Functional identification of rice *syn*-copalyl diphosphate synthase and its role in initiating biosynthesis of diterpenoid phytoalexin/allelopathic natural products. *Plant J.* **39**, 309–318
- 11 Otomo, K., Kenmoku, H., Oikawa, H., Konig, W. A., Toshima, H., Mitsuhashi, W., Yamane, H., Sassa, T. and Toyomasu, T. (2004) Biological functions of *ent*- and *syn*-copalyl diphosphate synthases in rice: key enzymes for the branch point of gibberellin and phytoalexin biosynthesis. *Plant J.* **39**, 886–893
- 12 Cho, E.-M., Okada, A., Kenmoku, H., Otomo, K., Toyomasu, T., Mitsuhashi, W., Sassa, T., Yajima, A., Yabuta, G., Mori, K. et al. (2004) Molecular cloning and characterization of a cDNA encoding *ent*-cassa-12,15-diene synthase, a putative diterpenoid phytoalexin biosynthetic enzyme, from suspension-cultured rice cells treated with a chitin elicitor. *Plant J.* **37**, 1–8
- 13 Nemoto, T., Cho, E.-M., Okada, A., Okada, K., Otomo, K., Kanno, Y., Toyomasu, T., Mitsuhashi, W., Sassa, T., Minami, E. et al. (2004) Stemar-13-ene synthase, a diterpene cyclase involved in the biosynthesis of the phytoalexin oryzalexin S in rice. *FEBS Lett.* **571**, 182–186
- 14 Otomo, K., Kanno, Y., Motegi, A., Kenmoku, H., Yamane, H., Mitsuhashi, W., Oikawa, H., Toshima, H., Itoh, H., Matsuoka, M. et al. (2004) Diterpene cyclases responsible for the biosynthesis of phytoalexins, momilactones A, B, and oryzalexins A-F in rice. *Biosci. Biotechnol. Biochem.* **68**, 2001–2006
- 15 Kanno, Y., Otomo, K., Kenmoku, H., Mitsuhashi, W., Yamane, H., Oikawa, H., Toshima, H., Matsuoka, M., Sassa, T. and Toyomasu, T. (2006) Characterization of a rice gene family encoding type-A diterpene cyclases. *Biosci. Biotechnol. Biochem.* **70**, 1702–1710
- 16 Wilderman, P. R., Xu, M., Jin, Y., Coates, R. M. and Peters, R. J. (2004) Identification of *syn*-pimara-7,15-diene synthase reveals functional clustering of terpene synthases involved in rice phytoalexin/allelochemical biosynthesis. *Plant Physiol.* **135**, 2098–2105
- 17 Morrone, D., Jin, Y., Xu, M., Choi, S.-Y., Coates, R. M. and Peters, R. J. (2006) An unexpected diterpene cyclase from rice: functional identification of a stemodene synthase. *Arch. Biochem. Biophys.* **448**, 133–140
- 18 Xu, M., Wilderman, P. R., Morrone, D., Xu, J., Roy, A., Margis-Pinheiro, M., Upadhyaya, N., Coates, R. M. and Peters, R. J. (2007) Functional characterization of the rice kaurene synthase-like gene family. *Phytochemistry* **68**, 312–326
- 19 Cyr, A., Wilderman, P. R., Determan, M. and Peters, R. J. (2007) A modular approach for facile biosynthesis of labdane-related diterpenes. *J. Am. Chem. Soc.* **129**, 6684–6685
- 20 Prisic, S. and Peters, R. J. (2007) Synergistic substrate inhibition of *ent*-copalyl diphosphate synthase: a potential feed-forward inhibition mechanism limiting gibberellin metabolism. *Plant Physiol.* **144**, 445–454
- 21 Peters, R. J., Ravn, M. M., Coates, R. M. and Croteau, R. B. (2001) Bifunctional abietadiene synthase: free diffusive transfer of the (+)-copalyl diphosphate intermediate between two distinct active sites. *J. Am. Chem. Soc.* **123**, 8974–8978
- 22 Harris, L. J., Saparno, A., Johnston, A., Prisic, S., Xu, M., Allard, S., Kathiresan, A., Ouellet, T. and Peters, R. J. (2005) The maize *An2* gene is induced by *Fusarium* attack and encodes an *ent*-copalyl diphosphate synthase. *Plant Mol. Biol.* **59**, 881–894
- 23 Morrone, D., Lowry, L., Determan, M. K., Hershey, D. M., Xu, M. and Peters, R. J. (2010) Increasing diterpene yield with a modular metabolic engineering system in *E. coli*: comparison of MEV and MEP isoprenoid precursor pathway engineering. *Appl. Microbiol. Biotechnol.* **85**, 1893–1906
- 24 Barrero, A. F. and Altarejos, J. (1993) ¹³C NMR data for labdane diterpenoids. *Magn. Reson. Chem.* **31**, 299–308
- 25 Xu, M., Wilderman, P. R. and Peters, R. J. (2007) Following evolution's lead to a single residue switch for diterpene synthase product outcome. *Proc. Natl. Acad. Sci. U.S.A.* **104**, 7397–7401
- 26 Wilderman, P. R. and Peters, R. J. (2007) A single residue switch converts abietadiene synthase into a pimaradiene specific cyclase. *J. Am. Chem. Soc.* **129**, 15736–15737
- 27 VanEtten, H. D., Mansfield, J. W., Bailey, J. A. and Farmer, E. E. (1994) Two classes of plant antibiotics: phytoalexins versus 'phytoanticiplins'. *Plant Cell* **6**, 1191–1192
- 28 International Rice Genome Sequencing Project (2005) The map-based sequence of the rice genome. *Nature* **436**, 793–800
- 29 Seo, S., Seto, H., Koshino, H., Yoshida, S. and Ohashi, Y. (2003) A diterpene as an endogenous signal for the activation of defense responses to infection with *tobacco mosaic virus* and wounding in tobacco. *Plant Cell* **15**, 863–873
- 30 Aharoni, A., Giri, A. P., Verstappen, F. W., Berteau, C. M., Sevenier, R., Sun, Z., Jongmsa, M. A., Schwab, W. and Bouwmeester, H. J. (2004) Gain and loss of fruit flavor compounds produced by wild and cultivated strawberry species. *Plant Cell* **16**, 3110–3131
- 31 Davidovich-Rikanati, R., Lewinsohn, E., Bar, E., Iijima, Y., Pichersky, E. and Sitrit, Y. (2008) Overexpression of the lemon basil α -zingiberene synthase gene increases both mono- and sesquiterpene contents in tomato fruit. *Plant J.* **56**, 228–238
- 32 Stofer Vogel, B., Wildung, M. R., Vogel, G. and Croteau, R. (1996) Abietadiene synthase from grand fir (*Abies grandis*). *J. Biol. Chem.* **271**, 23262–23268
- 33 Schepmann, H. G., Pang, J. and Matsuda, S. P. (2001) Cloning and characterization of *Ginkgo biloba* levopimaradiene synthase, which catalyzes the first committed step in ginkgolide biosynthesis. *Arch. Biochem. Biophys.* **392**, 263–269
- 34 Gao, W., Hillwig, M. L., Huang, L., Cui, G., Wang, X., Kong, J., Yang, B. and Peters, R. J. (2009) A functional genomics approach to tanshinone biosynthesis provides stereochemical insights. *Org. Lett.* **11**, 5170–5173
- 35 Nelson, D. R., Schuler, M. A., Paquette, S. M., Werck-Reichhart, D. and Bak, S. (2004) Comparative genomics of rice and *Arabidopsis*. Analysis of 727 cytochrome P450 genes and pseudogenes from a monocot and a dicot. *Plant Physiol.* **135**, 756–772
- 36 Swaminathan, S., Morrone, D., Wang, Q., Fulton, D. B. and Peters, R. J. (2009) CYP76M7 is an *ent*-cassadiene C11 α -hydroxylase defining a second multifunctional diterpenoid biosynthetic gene cluster in rice. *Plant Cell* **21**, 3315–3325
- 37 Shimura, K., Okada, A., Okada, K., Jikumaru, Y., Ko, K.-W., Toyomasu, T., Sassa, T., Hasegawa, M., Kodama, O., Shibuya, N. et al. (2007) Identification of a biosynthetic gene cluster in rice for momilactones. *J. Biol. Chem.* **282**, 34013–34018