


oxide; presumably the formation of carbon dioxide must also be accompanied by the formation of water. By approximate measurement the amount of carbon dioxide was that which should be expected from the amount of selenium dioxide used in excess of that required for the condensation reaction.

This reaction resembles those which Kharasch<sup>2</sup>

(2) M. S. Kharasch and M. T. Gladstone, *THIS JOURNAL*, **65**, 15 (1943).

has carried out with peroxides and it appears likely that it proceeds by a similar mechanism



RESEARCH AND DEVELOPMENT LABORATORIES  
UNIVERSAL OIL PRODUCTS COMPANY  
RIVERSIDE, ILLINOIS

RECEIVED MAY 10, 1947

## COMMUNICATIONS TO THE EDITOR

### A RAPID METHOD OF PREPARING $\text{NaC}^{14}\text{N}$ FROM $\text{BaC}^{14}\text{O}_3$

Sir:

Studies on the exchange of radiocyanide ion with cyanide complexes have been initiated recently in this Laboratory. Since radiocarbon is presently available only in the form of barium carbonate, it was necessary to investigate various means of converting it into a soluble cyanide.

A possible procedure is that of Cramer and Kistiakowsky.<sup>1,2</sup> This makes use of the reaction of gaseous ammonia and carbon dioxide with a potassium mirror. The procedure is somewhat elaborate and requires the use of a vacuum apparatus. For this reason, some simpler method of accomplishing the conversion would be highly desirable.

Several exploratory experiments were carried out on the use of electro-positive metals as reductants.<sup>3</sup> It was found that insignificant yields of cyanide resulted on heating zinc, aluminum, magnesium or sodium with barium carbonate, in a nitrogen atmosphere. This is in agreement with Loftfield's report.<sup>2</sup>

Preliminary results indicate, however, that yields of 75–80% can be obtained by heating sodium azide and barium carbonate in a nitrogen atmosphere. The method has the advantage of requiring only about thirty minutes and not necessitating the use of any special apparatus.

The procedure is: 0.1 g. of barium carbonate is mixed with 1 g. of sodium azide (Amend Drug and Chemical Co., N. Y.) in a six-inch test-tube, and a slow stream of nitrogen is directed into the mouth of the tube. The mixture is then heated carefully so as to maintain a steady but not too rapid decomposition of the azide. The fumes of sodium oxide may be drawn off into a hood or through a funnel connected to an aspirator.

(1) Cramer and Kistiakowsky, *J. Biol. Chem.*, **137**, 549 (1941).

(2) Loftfield, "The Preparation of Carbon-fourteen Labelled Hydrogen Cyanide, Alanine and Glycine," Circular C-3, Isotopes Branch, United States Atomic Energy Commission, June, 1947.

(3) A portion of these experiments was carried out by Mr. M. Volpe.

When the decomposition is complete, the test-tube is heated at a dull red heat for ten minutes. After cooling, water is added dropwise until all of the sodium present has reacted. The solution is then diluted, acidified with sulfuric acid, and the hydrogen cyanide distilled over into a slight excess of sodium hydroxide solution.

Four experiments were made with inactive barium carbonate. The resulting cyanide was determined by adding ammonia and potassium iodide and titrating with silver nitrate solution. The average yield was  $78 \pm 2\%$ . In addition, two runs were made with added  $\text{BaC}^{14}\text{O}_3$ .<sup>4</sup> The specific gravity of the radiocyanide, counted as silver cyanide, was within experimental error of the calculated value, indicating that the radiochemical yield is the same as the analytical yield.

Much of the cyanide is formed during the final heating when no sodium azide is present. This suggests that the actual reaction may involve not the sodium azide, but the sodium nitride formed by its decomposition.

Support of this investigation by a grant-in-aid from The Research Corporation is gratefully acknowledged.

(4) Supplied by the U. S. Atomic Energy Commission.

DEPARTMENT OF CHEMISTRY  
UNIVERSITY OF SOUTHERN CALIFORNIA  
LOS ANGELES, CALIFORNIA ARTHUR W. ADAMSON  
RECEIVED SEPTEMBER 10, 1947

### POLY-LYSINE

Sir:

On extending experiments concerning polymerization of amino acids<sup>1</sup> to basic amino acids, we succeeded in preparing poly-lysine. This polymer represents the first synthetic basic  $\alpha$ -poly-amide and as it is water soluble, it may serve as a useful model in protein research.

A suitable monomer was found in  $\epsilon$ -carboxybenzoxy- $\alpha$ -carboxyl-L-lysine anhydride (I)<sup>2</sup> which

(1) Frankel and Katchalski, *THIS JOURNAL*, **64**, 2264 (1942); **64**, 2268 (1942).

(2) Bergmann, Zervas and Ross, *J. Biol. Chem.*, **111**, 245 (1935).