

Green Chemistry

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: S. Das, J. Kollmann, Y. Zhang, W. Schilling, T. Zhang and D. Riemer, *Green Chem.*, 2019, DOI: 10.1039/C9GC00485H.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [author guidelines](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the ethical guidelines, outlined in our [author and reviewer resource centre](#), still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

ARTICLE

Simple Ketone as an Efficient Metal-Free Catalyst for Visible-Light-Mediated Diels-Alder and Aza-Diels-Alder Reactions

Jiri Kollmann[§], Yu Zhang[§], Waldemar Schilling, Tong Zhang, Daniel Riemer, Shoubhik Das*[§]Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Diels-Alder reactions are highly effective between electron-rich dienes and electron-poor dienophiles. However, these reactions with electron-rich dienophiles are limited and require forcing conditions. Based on this, an efficient metal-free homogeneous system has been developed for the Diels-Alder reactions between electron-rich dienophiles and dienes under visible-light conditions. Additionally, this catalyst has shown excellent reactivity for the aza-Diels-Alder reactions. This simple catalyst is commercially available, non-toxic, cheap and has shown excellent reactivity which is comparable and in some cases even better than the reported metal-based catalysts. Finally, the mechanism of this reaction has been proposed based on the experimental evidence.

Introduction

Diels-Alder (D-A) reactions are the fundamental strategy for the syntheses of versatile cyclohexene derivatives with the formation of new stereogenic centers.¹ These reactions have been well-documented in organic syntheses, in pharmaceutical industry and in biochemistry.² It should be noted that D-A reactions are highly effective between electron-rich dienes and electron-poor dienophiles, which has been explained based on the frontier molecular orbital (FMO) theory.^{3a} To enhance the conversion as well as the stereoselectivity of thermally-initiated organocatalytic Diels-Alder reactions, different strategies have been developed. Evans group established oxazolidinones as chiral organic auxiliaries which sterically hindered the attack of the diene on one face of the dienophile.^{3b} MacMillan and co-workers developed imidazolidinones as highly enantioselective chiral organocatalysts which form iminium species with enones. The iminium species lower the LUMO of the dienophile and sterically block one face from being attacked by the diene.^{3c} However, electronically mismatched coupling between dienes and dienophiles is highly challenging and require strict conditions.¹⁻² This electronically mismatched D-A reactions can be achieved by the formation of radical cation intermediates from dienophiles via one electron oxidation by using stoichiometric oxidants such as aminium salts or by photoinitiated electron transfer (PET) with photosensitizers.^{4,5}

In contrast, visible-light-mediated radical cation generation from the electron-rich dienophile is more sustainable and does not require any stoichiometric oxidant (**Figure 1**).⁶ Based on this, Yoon and co-workers developed a Ru complex which showed excellent reactivity towards electron-rich dienophiles under visible-light conditions.⁷ In this case the radical cation was generated by the Ru catalyst with the

turn over. In fact, they also investigated metal-free photocatalysts, however their efforts were not successful.^{7a} After then, Shores group and Ferreira group developed Cr complexes under UV light irradiation.⁸ However, the Cr catalyst required longer reaction time and excess of diene (10 equiv.) to achieve full conversion.^{8b}




Figure 1 Homogeneous catalysts for photocatalytic D-A reactions of electron-rich dienophiles.

Parallel to the D-A reactions, aza-Diels-Alder reactions are also efficient and powerful approaches for the syntheses of *N*-heterocycles in a straightforward way.⁹ With the same concept of the generation of radical cation intermediates from electron-rich dienophiles, iminium ions can be generated from imines with the aid of visible light.⁶ Based on this, Zhang, Arai and Ohkuma groups developed visible-light-promoted aza-Diels-Alder reactions with the Ru and Cr catalysts.¹⁰ It should be noted that in all the cases of visible-light-mediated D-A or aza-D-A reactions, only transition metal based photocatalysts exist. In contrast, organic dyes are commercially available, cheaper in price, non-toxic in nature and are an interesting avenue to explore.¹¹ Based on all the above information and to

Institut für Organische und Biomolekulare Chemie, Georg-August-Universität Göttingen, Tammannstraße 2, 37077 Göttingen, Germany

Email: Shoubhik.Das@chemie.uni-goettingen.de

§ J. K. and Y. Z. contributed equally.

Electronic Supplementary Information (ESI) available: See DOI: 10.1039/x0xx00000x

aid of the visible-light and air/oxygen was utilized for the catalyst




develop our own interest in metal-free catalysis, herein we describe an efficient metal-free homogeneous catalyst for visible-light-mediated Diels-Alder and aza-Diels-Alder reactions.¹²

Results and discussion

At the outset of our reaction, we focused on the organic dyes which have high oxidation potential (e.g. in case of fluorenone +1.7 V vs SCE)^{11a} to generate radical cation from trans-anethole (**1a**) (+1.1 V vs SCE).^{7a} We started the optimization using trans-anethole (**1a**) as a dienophile and 2,3-dimethyl-1,3-butadiene (**1b**) as diene under air (Table S1; see supporting information). Eight different organic dyes were applied using nitromethane (MeNO₂) as solvent and MgSO₄ as desiccant under the irradiation of 12 W blue LED for 4 h. Among them, fluorenone showed 81% yield of the corresponding Diels-Alder product. Subsequently, other aprotic polar solvents such as THF, DMF, DMSO and ACN were investigated but did not show any improvement of the yield. Nitromethane was the best solvent in this reaction as aprotic polar solvents with higher dipole moment at comparable polarities can solvate the formed radical cation intermediate which enhances the product formation.¹³

Scheme 1 Scope of the fluorenone-catalyzed Diels-Alder reactions under visible-light irradiation^{a-c}.



[a] Reaction conditions: air, 12 W blue LED, dienophiles (0.5 mmol, 1 eq.), dienes (3 eq.), photocatalyst (3 mol%), 5 mL solvent, room temperature. [b] Yield was determined by NMR yield using iodoform as an internal standard. [c] Isolated yields.

With these optimized reaction conditions in hand, we extended the scope of this metal-free system to other electron-rich dienophiles and dienes (Scheme 1; entries 1c–11c). The formation of [4+2] cycloadducts proceeded smoothly using 2,3-dimethyl-1,3-butadiene as a diene and other electron-rich dienophiles such as myrcene, isoprene etc. (Scheme 1; entries 2c–4c). In addition to these, 1,2-dimethoxy-4-propenylbenzene was also effectively explored as a dienophile with a series of dienes (Scheme 1; entries 5c–7c). Furthermore, 2,5-furandione was also suitable in our system and generated corresponding cyclohexene derivatives in excellent yields in 2 h (Scheme 1; entries 8c–9c). It should be noted that for 2,5-furandione, expensive metal-based catalysts are required to obtain


the desired product.¹⁴ However, with our cheap and metal-free fluorenone catalyst, these cyclohexene derivatives can be achieved with excellent yields in only 2 h. Additionally, 9-vinylcarbazole was also capable of acting as a dienophile and resulted **11c** with 2,3-dimethyl-1,3-butadiene as a diene.

Inspired by the excellent reactivity of our catalyst in [4+2] cycloaddition reactions, we sought to apply these reaction conditions to aza-D-A reactions. Our main target was to synthesize important *N*-heterocycles in a straightforward way. For this purpose, we aimed to synthesize 2,3-dihydropyridin-4(1*H*)-ones derivatives as these have wide applications for the treatment of emergent infectious diseases.¹⁵ Additionally, these compounds have been utilized for the syntheses of piperidine containing natural products and bioactive molecule synthesis.¹⁶ In general, syntheses of 2,3-dihydropyridin-4(1*H*)-ones derivatives require either expensive metal based catalysts or longer reaction time.¹⁷ Gratifyingly, fluorenone showed an excellent yield within 2 h when (*E*)-1-(4-methoxyphenyl)-*N*-phenylmethanimine (**1aa**) as a dienophile and Danishefsky's diene (**1ab**) were applied to synthesize 2-(4-methoxyphenyl)-1-phenyl-2,3-dihydropyridin-4(1*H*)-one (**1ac**) as the product (Table S2). Different ratios of diene and dienophile were also investigated and to our delight, ratio 2:1 proceeded smoothly with the formation of 96% of the desired product in 2 h.

With these optimized reaction conditions, we applied fluorenone catalyst to other imines to obtain diverse 2,3-dihydropyridin-4(1*H*)-ones derivatives (Scheme 2). Various electron-rich imines showed high reactivities with the formation of desired products in 2 h up to 96% yield. The catalyst showed high tolerance towards alkyl, aryl and heteroaryl substituents (Scheme 2; entries 2ac–10ac). Additionally, six new 2,3-dihydropyridin-4(1*H*)-ones derivatives have been synthesized whose biological activities should be further investigated considering the promising lead compounds against human infectious diseases (Scheme 2; entries 3ac–4ac, 7ac–10ac).¹⁶ Expediently, with this fluorenone catalyst, we have been able to synthesize previously reported bioactive derivatives which were evaluated in vitro against wide spectrum of viruses (Scheme 3; entries 1ac, 11ac–12ac).^{15a} The scope of the fluorenone-catalyzed aza-Diels-Alder reactions was not only limited to the synthesis of 2,3-dihydropyridin-4(1*H*)-one derivatives but also applied to synthesize other important heterocycles (Scheme 4). For example, when the Danishefsky's diene was replaced by 1-methoxy-1,3-butadiene, it generated 1,2,3,6-tetrahydropyridine derivative (Scheme 4; entry 13ac). In addition to this, 2,3-dimethyl-1,3-butadiene reacted with 1,4-benzoquinone to generate 3,4-dimethyl-1-oxaspiro[5.5]undeca-3,7,10-trien-9-one (Scheme 4; entry 14ac).

Scheme 2 Scope of the fluorenone-catalyzed aza-Diels-Alder reactions under visible light irradiation^{a-c}.






[a] Reaction conditions: air, 12 W blue LED, imines (0.10 mmol, 1 eq.), diene (2 eq.), photocatalyst (3 mol%), 5 mL solvent, room temperature. [b] Yield was determined by NMR yield using iodoform as an internal standard. [c] Isolated yields.

Considering the broad scope of this catalyst, we sought to enquire the role of the catalyst, air, and light source in this reaction. Control experiments showed no product formation in the absence of light or photocatalyst (see supporting information). Notably, 60% yield was achieved under nitrogen, which clearly indicates that there is a chain propagation mechanism similar to the earlier report of Ru complex.^{7a} The singlet oxygen in the air acts as an indispensable electron mediator which further increases the yield. Furthermore, the effect of different quenchers was investigated to recognize the reactive oxygen species from air and possible intermediates (Table 1).^{12c-e} When 2,6-di-tert-butyl-4-methylphenol (BHT) or 2,2,6,6-tetramethyl-1-piperidinyloxy (TEMPO) were added to the reaction mixture, the reaction was completely inhibited, which proved a radical pathway. Furthermore, addition of CuCl₂ to the reaction mixture showed lower yields, which showed the involvement of single electron processes in this photocatalytic system. The addition of sodium azide also showed a decreased yield, indicating the involvement of singlet oxygen, or activated oxygen species.


To acquire further information about the reaction mechanism, Stern-Volmer quenching experiments were carried out (see supporting information). The excited state of the photocatalyst was quenched by trans-anethole.¹⁸ As shown in Figure S3, a clear decrease of light emission was observed with the increasing concentration of trans-anethole (1a) and no change was observed with different concentrations of 2,3-dimethyl-1,3-butadiene.

Scheme 3 Syntheses of bioactive derivatives using fluorenone catalyst.



Reaction conditions: air, 12 W blue LED, imines (0.10 mmol, 1 eq.), diene (2 eq.), photocatalyst (3 mol%), 5 mL solvent, room temperature.

Scheme 4 Fluorenone-catalyzed hetero-Diels-Alder reactions under visible light irradiation.



Reaction conditions: air, 12 W blue LED, imine or 1,4-benzoquinone (0.10 mmol, 1 eq.), dienes (2 eq.), photocatalyst (3 mol%), 5 mL solvent, room temperature.

Combining all the mechanistic experiments, we have been able to propose the mechanism for visible-light-mediated D-A and aza-D-A reactions in Figure 2. At first, the photocatalyst was excited to the photo-excited state by the irradiation of visible light and underwent a single electron transfer (SET) with trans-anethole (1a). In fact the redox potential value of the catalyst (+1.7 V vs SCE)^{11a} is enough to oxidize 1a (+1.1 V vs SCE).^{14d} The resulting radical cation 1a^{•+} reacted with the diene via [4+2] cycloaddition to afford radical cation product 1c^{•+}. This product 1c^{•+} accepted one electron mainly from another equivalent of 1a with a chain propagation step to form the final product (1c).^{7a} Meanwhile, oxygen will also be involved partly in electron donating. Here the role of air/oxygen was to turn over the reduced photocatalyst to the original state. The reported value for the reduction potential of the excited-state fluorenone resides at -0.61 V vs SCE,^{11d} which is sufficient for the reduction of molecular oxygen to its superoxide radical form (O₂/O₂^{•-}) with the reduction potential residing at -0.56 V vs SCE.¹⁹

Table 1 Quenching experiments for fluorenone catalyzed cycloaddition reactions.



ARTICLE

Journal Name




Reaction conditions: air, 12 W blue LED, 0.5 mmol dienophiles (1 eq.), 1.5

Quencher	Equivalents	Yield [%]	Scavenger for
BHT	1.0	0	radical
TEMPO	1.0	0	radical
CuCl ₂	1.0	73	single electron
NaN ₃	1.0	63	singlet oxygen

mmol dienes (3 eq.), photocatalyst (3 mol%), 5 mL solvent, room temperature.

Scheme 5 (a) Examples of bioactive cyclobutane lignan natural products. (b) Our work on [2+2] cycloaddition reactions.



Reaction conditions: air, 12 W blue LED, dienophiles (0.5 mmol, 1 eq.), dienes (3 eq.), photocatalyst (3 mol%), 5 mL solvent, room temperature.

Figure 2 Proposed reaction mechanism for the fluorenone catalyzed Diels-Alder reactions.

The [2+2] cycloaddition was also proceeded well via SET with product **1a^{•+}** in the absence of diene.²⁰ In fact, cyclobutane derivatives have wide applications in natural product syntheses and many of them already have become promising candidates as anticancer, antiviral and antifungal drugs.^{20b} Additionally, lignan and neolignan type compounds also contain a cyclobutane motif (**Scheme 5a**).^{20b} Based on all this information, we applied fluorenone catalyst to promote [2+2] cycloaddition reactions to obtain cyclobutane derivatives (**Scheme 5b**; entries **1d–2d**). To our delight, the catalyst not only showed the activity for homodimerization of alkenes (**1d**) but also showed an excellent reactivity towards heterodimerization of alkene (**2d**).

Conclusions

In summary, we have developed fluorenone as a metal-free catalyst for the visible-light-mediated cycloaddition reactions. This catalyst is very cheap and commercially available and has shown excellent substrates scope for [4+2] and [2+2] cycloaddition reactions with electron-rich dienophiles. In addition, we have been able to synthesize bioactive molecules via aza-Diels Alder reactions. We believe this protocol can be further applied and extended for the syntheses of pharmaceuticals and natural products. Additionally, detailed mechanistic studies revealed the role of the catalyst and oxygen and led to the proposed mechanism of this reaction.



Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We thank Fonds der Chemischen Industrie (FCI, Liebig-Fellowship to S.D.) and Chinese Scholarship Council (CSC funding to Y.Z.) for the financial support. We are highly thankful to Prof. Dr. Lutz Ackermann for his support behind our work.

Notes and references

- (a) X. Jiang and R. Wang, *Chem. Rev.*, 2013, **113**, 5515. (b) C. K. Nicolaou, A. Snyder, T. Montagnon and G. Vassilikogiannakis, *Angew. Chem. Int. Ed.*, 2002, **41**, 1668.
- (a) G. Desimoni and G. Faita, P. Quadrelli, *Chem. Rev.*, 2015, **115**, 9922; (b) T. Gatzemeier, M. van Gemmeren, Y.-W. Xie, D. Höfler, M. Leutzsch and B. List, *Science*, 2016, **351**, 949; (c) G.-L. Li, T. Liang, L. Wojtas and J. C. Antilla, *Angew. Chem., Int. Ed.*, 2013, **52**, 4628; (d) R. F. Higgins, S. M. Fatur, S. G. Shepard, S. M. Stevenson, D. J. Boston, E. M. Ferreira, N. H. Damrauer, A. K. Rappe and M. P. Shores, *J. Am. Chem. Soc.*, 2016, **138**, 5451.
- (a) K. N. Houk, *Acc. Chem. Res.*, 1975, **8**, 361. (b) D. A. Evans, K. T. Chapman and J. Bisaha, *J. Am. Chem. Soc.*, 1984, **106**, 4261. (c) K. A. Ahrendt, C. J. Borths and D. W. MacMillan, *J. Am. Chem. Soc.*, 2000, **122**, 4243.
- (a) D. J. Bellville, D. W. Wirth and N. L. Bauld, *J. Am. Chem. Soc.*, 1981, **103**, 718; (b) N. L. Bauld, D. J. Bellville, R. Pabon, R. Chelsky and G. Green, *J. Am. Chem. Soc.*, 1983, **105**, 2378; (c) D. J. Bellville, N. L. Bauld, R. Pabon and S. A. Gardner, *J. Am. Chem. Soc.*, 1983, **105**, 3584.
- (a) C. R. Jones, B. J. Allman, A. Mooring and B. Saphic, *J. Am. Chem. Soc.*, 1983, **105**, 652; (b) R. A. Pabon, D. J. Bellville, and N. L. Bauld, *J. Am. Chem. Soc.*, 1983, **105**, 5158; (c) G. C. Calhoun and G. B. Schuster, *J. Am. Chem. Soc.*, 1984, **106**, 6870.
- (a) C. K. Prier, D. A. Rankic and D. W. MacMillan, *Chem. Rev.*, 2013, **113**, 5322; (b) Y.-B. Zhao and M. Antonietti, *Angew. Chem. Int. Ed.*, 2017, **56**, 9336.
- (a) S. Lin, M. A. Ischay, C. G. Fry and T. P. Yoon, *J. Am. Chem. Soc.*, 2011, **133**, 19350. (b) A. E. Hurtley, M. A. Cismesia, M. A. Ischay and T. P. Yoon, *Tetrahedron*, 2011, **67**, 4442.
- (a) S. M. Stevenson, R. F. Higgins, M. P. Shores and E. M. Ferreira, *Chem. Sci.*, 2017, **8**, 654; (b) S. M. Stevenson, M. P. Shores and E. M. Ferreira, *Angew. Chem. Int. Ed.*, 2015, **54**, 6506; (c) R. F. Higgins, S. M. Fatur, S. G. Shepard, S. M. Stevenson, D. J. Boston, E. M. Ferreira, N. H. Damrauer, A. K. Rappe and M. P. Shores, *J. Am. Chem. Soc.*, 2016, **138**, 5451.
- (a) J. M. Eagan, M. Hori, J.-B. Wu, K. S. Kanyiva and S. A. Snyder, *Angew. Chem. Int. Ed.*, 2015, **54**, 7842; (b) G. Masson, C. Lalli, M. Benohouda and G. Dagousseta, *Chem. Soc. Rev.*, 2013, **42**, 902; (c) H. Mandai, K. Mandai, M. L. Snapper and A. H. Hoveyda, *J. Am. Chem. Soc.*, 2008, **130**, 17961.
- (a) N. Arai and T. Ohkuma, *J. Org. Chem.*, 2017, **82**, 7628; (b) W.-H. Wang, Y. Yuan, X.-S. Gao, B. Hu, X.-M. Xie and Z.-G. Zhang, *ChemCatChem.*, 2018, **10**, 2878.
- (a) N. A. Romero and D. A. Nicewicz, *Chem. Rev.*, 2016, **116**, 10075; (b) M. Majek and A. Von Wangelin, *J. Acc. Chem. Res.*, 2016, **49**, 2316; (c) Y. Zhang, D. Riemer, W. Schilling, J. Kollmann and S. Das, *ACS Catal.*, 2018, **8**, 6659; (d) W. Schilling, D. Riemer, Y. Zhang, N. Hatami and S. Das, *ACS Catal.*, 2018, **8**, 5425. (e) D. Riemer, W. Schilling, A. Goetz, Y. Zhang, S. Gehrke, I. Tkach, O. Holoczeki and S. Das, *ACS Catal.*, 2018, **8**, 11679; (f) M. Majek and A. J. Von Wangelin, *Angew. Chem. Int. Ed.*, 2015, **54**, 2270.
- (a) D. Riemer, B. Mandaviya, W. Schilling, A. C. Götz, T. Köhl, M. Finger and S. Das, *ACS Catal.*, 2018, **8**, 3030; (b) D. Riemer, P. Hirapara and S. Das, *ChemSusChem.*, 2016, **9**, 1916; (c) P. Hirapara, D. Riemer, N. Hazra, J. Gajera, M. Finger and S. Das, *Green Chem.*, 2017, **19**, 5356.
- V. Santacroce, R. Duboc, M. Malacria, G. Maestri, G. Masson, *Eur. J. Org. Chem.*, 2017, **15**, 2095.
- (a) K. Mori, T. Hara, T. Mizugaki, K. Ebitani and K. Kaneda, *J. Am. Chem. Soc.*, 2003, **125**, 11460. (b) Y. Ogasawara, S. Uchida, K. Yamaguchi and N. Mizuno, *Chem. Eur. J.*, 2009, **15**, 4343.
- (a) A. Peduto, A. Massa, A. D. Mola, P. de Caprariis, P. L. Colla, R. Loddo, S. Altamura, G. Maga and R. Filosa, *Chem. Biol. Drug Des.*, 2011, **77**, 441; (b) J. Sun, E.-Y. Xia, Q. Wu and C.-G. Yan, *ACS Combi. Sci.*, 2011, **13**, 421; (c) A. R. Ranade and G. I. Georg, *J. Org. Chem.*, 2014, **79**, 984; (d) S. Fustero, J. Miro, M. Sanchez-Rosello and C. Del Pozo, *Chem. Eur. J.*, 2014, **20**, 14126.
- (a) M. Ali, S. H. Ansari and J. S. Qadry, *J. Nat. Prod.*, 1991, **54**, 1271; (b) J. Matsuo, R. Okado and H. A. Ishibashi, *Org. Lett.*, 2010, **12**, 3266.
- (a) N. S. Josephsohn, M. L. Snapper and A. H. Hoveyda, *J. Am. Chem. Soc.*, 2003, **125**, 4018. (b) S. Kobayashi, S. Komiyama and H. Ishitani, *Angew. Chem. Int. Ed.*, 1998, **37**, 979; (c) S. Kobayashi, M. Ueno, S. Saito, Y. Mizuki, H. Ishitani and Y. Yamashita, *Pro. Nat. Acad. Sci.*, 2004, **101**, 5476. (d) S. Kaneko, Y. Kumatabara, S. Shimizu, K. Maruoka and S. Shirakawa, *Chem. Commun.*, 2017, **53**, 119; (e) Y. Takeda, D. Hisakuni, C.-H. Lin and S. Minakata, *Org. Lett.*, 2015, **17**, 318.
- D. M. Arias-Rotondoa and J. K. McCusker, *Chem. Soc. Rev.*, 2016, **45**, 5803.
- W. Huang, B.-C. Ma, H. Lu, R. Li, L. Wang, K. Landfester and K. A. I. Zhang, *ACS Catal.*, 2017, **7**, 5438.
- (a) M. A. Ischay, M. E. Anzovino, J.-N. Du and T. P. Yoon, *J. Am. Chem. Soc.*, 2008, **130**, 12886; (b) M. Riemer and D. A. Nicewicz, *Chem. Sci.*, 2013, **4**, 2625; (c) R. Li, B. C. Ma, W. Huang, L. Wang, D. Wang, H. Lu, K. Landfester and K. A. I. Zhang, *ACS Catal.*, 2017, **7**, 3097; (d) M. A. Ischay, M. S. Ament, T. P. Yoon, *Chem. Sci.*, 2012, **3**, 2807.

