

Solution Structure and Kinetic Study of Metal-Phosphine and -Phosphite Complexes. I. The Silver(I) System

E. L. Muetterties* and C. W. Alegranti

Contribution No. 1901 from the Central Research Department,
E. I. du Pont de Nemours and Company, Experimental Station,
Wilmington, Delaware 19898. Received January 26, 1972

Abstract: The character of the solution interaction of silver(I) salts with the phosphorus ligands (L), tri-*p*-tolylphosphine and triethyl phosphite, has been elucidated primarily through analysis of ^{31}P nmr data. The L_4Ag^+ cationic complex prevails in solution at low temperatures for all salts except cyanide, iodide, bromide, and chloride, provided that at least 4 equiv of phosphorus ligand per silver salt are present. The ligands are labile in L_4Ag^+ as shown by the loss of the Ag-P coupling at temperatures above $\sim -70^\circ$. The exchange is independent of free ligand concentration but dependent upon the nature of the counteranion. Ligand lability is higher in the phosphine than in the phosphite system. In the bromide and chloride systems the equilibrium, $\text{L}_4\text{Ag}^+\text{X}^- \rightleftharpoons \text{L}_3\text{AgX} + \text{L}$, was detected by adding excess ligand to the solution. In 3L-AgX solutions at $\sim -90^\circ$, either $\text{L}_3\text{Ag}^+\text{X}^-$ or L_3AgX forms; the determinant factor is the complexation character of the anion. $\text{L}_3\text{Ag}^+\text{X}^-$ was the predominant form for the PF_6^- , BF_4^- , and ClO_4^- salts and tetrahedral L_3AgX for the Cl^- , Br^- , I^- , NCO^- , CN^- , and S_2PF_2^- derivatives. Surprisingly, some L_3AgX compounds underwent disproportionation in solution at -90° : $2\text{L}_3\text{AgX}$ (or $\text{L}_3\text{Ag}^+\text{X}^-$) $\rightleftharpoons \text{L}_4\text{Ag}^+\text{X}^- + \text{L}_2\text{AgX}$. This behavior was found for the nitrate, triborohydride, and trifluoroacetate derivatives in the phosphine system and for the perchlorate and nitrate in the phosphite system. There was some evidence at 25° for dissociation of L_3AgX to L_2AgX and L. For the 2:1 solutions, L_2Ag^+ cationic complexes dominated in the PF_6^- , BF_4^- , and ClO_4^- group and neutral L_2AgX complexes for most other derivatives. Ligand lability in L_3Ag^+ was greater than that in L_2Ag^+ , and the same ordering obtained for the neutral L_3AgX and L_2AgX complexes. In the neutral L_3AgX and L_2AgX series, phosphorus ligand lability was sensitive to the binding power of the anion, with the cyanide complex most labile.

Trivalent phosphorus compounds as exemplified by phosphines and phosphites serve as a very important class of ligands in metal complex chemistry.^{1a} Especially distinctive are the catalytically active phosphine complexes which are a common feature of modern developments in homogeneous catalysis.^{1b} Explication of mechanistic details in these catalytic systems requires an understanding of the solution structure of the complexes, stereochemistry, and a knowledge of the kinetic and thermodynamic parameters of the phosphorus ligands. We are making a systematic study of such solution-state, metal complex chemistry with primary reliance on nuclear magnetic resonance data. A preliminary report on coinage metal-phosphine complexes² has appeared as well as a study of phosphino-metalloboranes.³ We report here extensive data and general qualitative conclusions about structural and kinetic features of phosphine and phosphite complexes of silver(I) salts.⁴

(1) (a) G. Booth, *Advan. Inorg. Chem. Radiochem.*, **6**, 1 (1964); (b) E. L. Muetterties, Ed., "Transition Metal Hydrides," Marcel Dekker, New York, N. Y., 1971.

(2) E. L. Muetterties and C. W. Alegranti, *J. Amer. Chem. Soc.*, **92**, 4114 (1970).

(3) E. L. Muetterties, W. G. Peet, P. A. Wegner, and C. W. Alegranti, *Inorg. Chem.*, **9**, 2447 (1970).

(4) For background literature and for leading references to work on silver(I) complexes of this type, see ref 5-8. Most of the solution studies were on dilute solutions of the complexes at room temperature. Colligative properties, electronic spectra, and conductivities were the primary data collected.

(5) C. Kowala and J. M. Swan, *Aust. J. Chem.*, **19**, 555 (1966).

(6) H. C. Brinkhoff, A. G. Matthijssen, and C. G. Oomes, *Inorg. Nucl. Chem. Lett.*, **7**, 87 (1971).

(7) K. J. Coskran, R. D. Bertrand, and J. G. Verkade, *J. Amer. Chem. Soc.*, **89**, 4535 (1967), report ^{31}P data at 25° for solutions of L_4AgClO_4 and L_4AgNO_3 with phosphite ligands.

(8) F. A. Cotton and D. M. L. Goodgame, *J. Chem. Soc.*, 5267 (1960), isolated tetrakis(phosphine)silver(I) salts with ClO_4^- , BrO_3^- , and NO_3^- as counterions.

Reference ligands (L) are tri-*p*-tolylphosphine and triethyl phosphite. These two ligands were selected primarily for the good, low-temperature solubility characteristics of the silver complex derivatives.

$[(\text{C}_6\text{H}_5)_3\text{P}]_4\text{Ag}^+$ and $[(\text{C}_2\text{H}_5\text{O})_3\text{P}]_4\text{Ag}^+$ Complexes

Isolable crystalline L_4AgX compounds were obtained in the reaction of most silver salts with four or more equivalents of tri-*p*-tolylphosphine.⁸ These tetrakis complexes dissolve in nonpolar solvents to give essen-

Table I. ^{31}P Nmr Parameters for $[(\text{C}_6\text{H}_5)_3\text{P}]_4\text{AgX}$ and $[(\text{C}_2\text{H}_5\text{O})_3\text{P}]_4\text{AgX}$ Complexes

X	$[(\text{C}_6\text{H}_5)_3\text{P}]_4\text{AgX}^a$		$[(\text{C}_2\text{H}_5\text{O})_3\text{P}]_4\text{AgX}^b$	
	$J_{^{31}\text{P Ag-P}}$	δ^c	$J_{^{31}\text{P Ag-P}}$	δ^c
F	224	25.3		
Cl	225	25.3	341	153
Br	230	25.5	340	153.1
I			340	153
SCN			342	153.1
B_2H_6	230	25.6		
S_2PF_2	225	25.6		
OOCFCF_3	230	25.7		
NO_3	224	25.5	342	153.2
PF_6	224	25.6		
ClO_4	225	25.6	341	153

^a Parameters were obtained at -80° . The solvent was dichloromethane. ^b Parameters were obtained at -100° . The solvent was 80/20 dichloromethane-toluene mixture. ^c Chemical shifts are reported in parts per million with reference to an internal capillary of $(\text{C}_6\text{H}_5)_3\text{P}$; tri-*p*-tolylphosphine is at 11.1 ppm and triethyl phosphite at 159.9 ppm.

tially the same ^{31}P spectrum (Figures 1 and 2 and Table I) which below -60° consists of two doublets arising




Figure 1. The ^{31}P 40.5-Mc spectrum of tetrakis(tri-*p*-tolylphosphine)silver hexafluorophosphate at -86° in dichloromethane.

from $^{107}\text{Ag}-^{31}\text{P}$ and $^{109}\text{Ag}-^{31}\text{P}$ spin-spin coupling.⁹ The analogous phosphite complexes yielded similar spectra. All chemical shifts for the phosphine set fell in a narrow range, and those for the phosphites fell in another narrow range (Table I). Nmr intensity relations between free and bound ligand for mixtures of known amounts of $\text{L}_4\text{Ag}^+\text{X}^-$ and L were determined to definitely establish that four phosphorus ligands are bound to the metal nucleus. The counterion in not bound. This state was inferred from the invariability of the nmr parameters for the phosphine (and phos-

Table II. Conductivity Data

Compound ^a	Equivalent conductance ($\Omega^{-1}\text{ cm}^2\text{ equiv}^{-1}$) in dichloromethane	
	10^{-2} equiv/l.	10^{-3} equiv/l.
(<i>n</i> -C ₃ H ₇) ₄ NCl	9.2	19.0
(<i>n</i> -C ₃ H ₇) ₄ NS ₂ PF ₂	14.6	28.0
(C ₆ H ₅) ₄ AsCl	35.0	52.0
(C ₆ H ₅) ₄ AsB ₃ H ₈	40.0	60.0
(C ₆ H ₅) ₄ AsNO ₃	34.9	55.0
(C ₆ H ₅) ₄ AsNO ₃ (-78°)	10.5	16.0
(C ₆ H ₅) ₄ AsOCCF ₃	36.0	56.0
(C ₆ H ₅) ₄ AsOCCF ₃ (-78°)	13.9	17.0
L' ₂ AgCl	0.1	
L' ₂ AgCl (-78°)	0.1	
L' ₃ AgCl	0.2	
L' ₃ AgCl (-78°)	0.07	
L' ₂ AgNO ₃	0.7	0.9
L' ₂ AgNO ₃ (-78°)	0.08	
L' ₄ AgNO ₃	20.0	
L' ₄ AgNO ₃ (-78°)	10.8	
L ₂ AgCl	0.2	
L ₃ AgCl	0.4	
L ₃ AgI	0.2	
L ₂ AgS ₂ PF ₂	0.7	1.0
L ₃ AgS ₂ PF ₂	3.5	
L ₂ AgO ₂ CCF ₃	2.4	4.0
L ₃ AgO ₂ CCF ₃	15.0	17.0
L ₃ AgO ₂ CCF ₃ (-78°)	6.9	9.7
L ₂ AgNO ₃	1.8	3.9
L ₃ AgNO ₃	26.0	28.0
L ₃ AgNO ₃ (-30°)	21.0	
L ₄ AgNO ₃		37.0
L ₄ AgNO ₃ (-30°)	28.0	
L ₂ AgPF ₆		60.0
L ₂ AgPF ₆ (-78°)	10.0	14.0
L ₃ AgPF ₆	46.0	58.0
L ₃ AgPF ₆ (-30°)	32.0	
L ₄ AgPF ₆	50.1	66.0
L ₃ AgClO ₄	45.1	55.0
L ₂ AgB ₃ H ₁₂ S	5.2	10.0
L ₃ AgB ₃ H ₁₂ S	33.0	39.2

^a L = (C₇H₇)₃P; L' = (C₂H₅O)₃P.

(9) The multiplets were identified from the relative peak intensities and the magnitude of the coupling constant. Ratios of $J(^{109}\text{Ag}-^{31}\text{P})/J(^{107}\text{Ag}-^{31}\text{P})$ were consistently close to the $\mu(^{109}\text{Ag})/\mu(^{107}\text{Ag})$ ratio of 1.149.





Figure 2. The ^{31}P 40.5-Mc spectrum of the tris(triethyl phosphite)silver nitrate at -90° in 80/20 dichloromethane-toluene. This tris complex partially disproportionates to give the bis and tetrakis complexes.

phite) complexes with change in counterion and was confirmed by conductivity studies which showed the complexes to have conductivities comparable to several tetraphenylarsonium salts (Table II).

Silver(I) chloride, bromide, and cyanide did not form isolable L_4AgX complexes, but the tetrakis complexes were detected for some salts in the nmr experiment by addition of phosphine to the L_3AgX solutions. Equilibrium constants calculated for (1) at low temperatures




from nmr intensities, are shown in Table III. The value

Table III. Equilibrium Data for $\text{L}_4\text{AgX} \rightleftharpoons \text{L}_3\text{AgX} + \text{L}$ in Dichloromethane

Anion	Ligand	K , mol/l.	Temp, $^\circ\text{C}$
F	(C ₇ H ₇) ₃ P	0.7 (est)	-100
Cl	(C ₂ H ₅ O) ₃ P	0.03	-100
Cl	(C ₇ H ₇) ₃ P	0.3	-86
Br	(C ₂ H ₅ O) ₃ P	0.04	-100
Br	(C ₇ H ₇) ₃ P	0.7	-86
I	(C ₂ H ₅ O) ₃ P	0.7	-100
I	(C ₇ H ₇) ₃ P	$>10^4$ (est)	-100
CN	(C ₂ H ₅ O) ₃ P	$>10^4$ (est)	-100
CN	(C ₇ H ₇) ₃ P	$>10^4$ (est)	-100

given for the fluoride is only an estimate. A competing fluorination reaction (2) made quantitative measure-




ments difficult. The marked tendency of cyanide and halide ions to bind, to the exclusion or near exclusion of a fourth phosphorus ligand, to L_3Ag^+ more strongly than other counterions employed (Table I) in this study is not unexpected.

Lability of the phosphorus ligands in L_4Ag^+ was evident in the temperature dependence of the ^{31}P nmr spectra. Broadening of the ^{31}P multiplets commenced at temperatures above $\sim -70^\circ$ and was followed by coalescence at -40 and -15° for phosphine and phosphite complexes, respectively. Exchange rates (Table IV) were essentially identical for the relatively nonpolarizable anions, e.g., NO_3^- , ClO_4^- , and PF_6^- in the phosphine series and in the phosphite series. Added phosphorus compound did not detectably alter the ligand exchange rate. Absence of a marked anion effect suggests the ligand exchange mechanism in these tetrahedral, closed shell (18 electron) complexes^{1b} closely approximates a simple dissociative $\text{S}_\text{N}1$ mechanism (3) where $k_1 \ll k_2$. Similar mechanistic con-

Table IV. Rate Data for Ag-P Bond Rupture in Silver(I) Complexes


Compound ^a	Approximate rate, ^{b,c} sec ⁻¹
L ₂ AgCl ^d	3612 (-45)
L ₂ AgNO ₃ ^d	4478 (+10)
L ₂ AgPF ₆ ^d	4731 (+25)
L ₂ AgPF ₆ ^d	4400 (+20)
L ₂ AgCl ^e	5245 (-45)
L ₂ AgNO ₃ ^e	6490 (-10)
L ₂ AgCl ^d	2675 (-50)
L ₂ AgPF ₆ ^d	3037 (-10)
L ₂ AgCl ^e	3894 (-45)
L ₂ AgClO ₄ ^e	4509 (-15)
L ₄ AgNO ₃ ^{d,f}	75 (-70)
	175 (-60)
	2000 (-40)
	13,000 (-20)
L ₄ AgPF ₆ ^d	2094 (-40)
L ₄ AgClO ₄ ^d	2094 (-40)
L ₄ AgNO ₃ ^e	3265 (-30)
L ₄ AgClO ₄ ^{e,f}	7 (-70)
	17 (-60)
	150 (-40)
	4500 (-10)
L ₄ AgNO ₃ ^d	3265 (-30)
L ₄ AgNO ₃ ^g	3265 (-60)
L ₄ AgNO ₃ ^h	3265 (-40)
L ₄ AgNO ₃ ^e	2095 (-40)
L ₄ AgNO ₃ ⁱ	2095 (-50)

^a L = (C₇H₇)₃P; L' = (C₂H₅O)₃P. ^b Temperatures in parentheses, accuracy $\pm 5^\circ$. ^c Based on coalescence temperature. ^d Dichloromethane. ^e 80/20 dichloromethane-toluene. ^f Based on line-shape analysis of nmr data. The E_a for dissociation of phosphine is 9 ± 1 kcal/mol and for phosphite is 11 ± 1 kcal/mol in the tetrakis complexes. ^g Toluene. ^h Acetonitrile. ⁱ 50/50 dichloromethane-toluene.



clusions^{10,11} have been reached for the isoelectronic ML₄ complexes in the nickel triad. The studies of Meier, *et al.*,¹⁰ showed conclusively that in ML₄ substitution reactions the ligand dissociation step (3) is rate determining.

In [(C₂H₅O)₃P]₄Ag⁺SCN⁻, the exchange rate is substantially higher than in the analogous perchlorate and hexafluorophosphate salts. Since the thiocyanate is a much more effective nucleophile than these other anions, it is possible that the ligand exchange mechanism for the thiocyanate more closely approximates an S_N2 (4) than an S_N1 (3) mechanism.^{12,13}



(10) M. Meier, F. Basolo, and R. G. Pearson, *Inorg. Chem.*, **8**, 795 (1969).

(11) C. A. Tolman, *J. Amer. Chem. Soc.*, **92**, 2956 (1970).

(12) It was noted above that added phosphorus compound did not affect the ligand exchange rates. In such reaction systems, a reaction intermediate or transition state analogous to L₄AgX in (4) with L substituted for X is much less likely than in the anion case because of steric factors and charge considerations.

(13) Addition of a tenfold molar excess of (C₂H₅)₃N⁺SCN⁻ to the L₄Ag⁺SCN⁻ solution did not affect the temperature dependence of the nmr line shapes. However, this addition of thiocyanate ion probably did not substantially raise the activity of the ion in solution because of ion pairing and because of the necessarily high concentrations employed in the nmr studies.

Ligand exchange rates for the L₄Ag⁺ series were slightly higher in the phosphine than in the phosphite complexes (Table IV). However, the dissociation constants K (k_1/k_2 in eq 1) must be just slightly higher for phosphite than phosphine in L₄Ag⁺. Triethyl phosphite addition¹⁴ to tetrakis(tri-*p*-tolylphosphine)silver salts led to the appearance of the ³¹P resonance line characteristic of free phosphine and of multiplets due to various Ag(I)-phosphine-phosphite mixed complexes. In the converse experiment, addition of phosphine¹⁴ gave a similar result; slightly more free phosphite appeared than free phosphine in the former experiment. This relative stability does not parallel the trend in the equilibrium data for eq 1 (Table III). In the isoelectronic nickel series, dissociation constants are much higher in a triarylphosphine than in triethyl phosphite complex.¹¹ We suggest that the small difference, if any, between the two ligands in the silver case reflects the formal positive charge of these complexes; phosphine should be far more effective than phosphite in reducing the charge on the metal atom. This effect nearly outweighs the steric factor that dominates ligand dissociation in the nickel complexes.^{11,15,16}

The rates of ligand exchange in L₄Ag⁺ are solvent dependent (Table IV); *e.g.*, the rate increases for [(C₂H₅O)₃P]₄Ag⁺ in going from dichloromethane to acetonitrile to toluene. These rates do not correlate with the dielectric constant of the medium; specific solvation effects must be operative.

Tetrahedral and Planar Tris(phosphine)- and Tris(phosphite)-Silver(I) Complexes

The L₃AgX complexes are of two limiting types: neutral, tetrahedral L₃AgX and ionic L₃Ag⁺X⁻. Included in the ionic phosphine set are the PF₆⁻ and ClO₄⁻ salts. Conductivities in dichloromethane were high (Table II). The nmr chemical shifts were at the lowest field in the tris series and coupling constants (Table V)

Table V. ³¹P Nmr Parameters for [(C₇H₇)₃P]₃AgX and [(C₂H₅O)₃P]₃AgX Complexes

X	[(C ₇ H ₇) ₃ P] ₃ AgX ^a $J_{107Ag-P}$	δ^c	[(C ₂ H ₅ O) ₃ P] ₃ AgX ^b $J_{107Ag-P}$	δ^c
F	273	22.5		
Cl	280	22.7	408	155.5
Br	278	21.7	398	153.2
I	266	18.8	390	150.1
CN	230	24.2	324	158.7
SCN	~270	~25	397	155.4
B ₃ H ₈	300	26.8		
S ₂ PF ₂	308	26.5		
OOCF ₃	310	27.3		
NO ₃	312	27.3	450	154.8
PF ₆	321	30.6		
ClO ₄			472	149.1
B ₃ H ₁₂ S	316	28.9		

^a Parameters were obtained at -80° . The solvent was dichloromethane. ^b Parameters were obtained at -100° . The solvent was 80/20 dichloromethane-toluene mixture. ^c Chemical shifts are reported in parts per million with reference to an internal capillary of (C₂H₅)₃P; tri-*p*-tolylphosphine is at 11.1 ppm and triethyl phosphite at 159.9 ppm.

(14) 0.5 mol of free ligand per L₄Ag⁺.

(15) C. A. Tolman, *J. Amer. Chem. Soc.*, **92**, 2953 (1970).


(16) Furthermore, Ag⁺ is larger than Ni²⁺.

were close to the estimated 303 Hz for sp^2 silver.¹⁷ Ionic tris(triethyl phosphite)silver derivatives included the nitrate and perchlorate which had coupling constants (^{107}Ag) comparable to the calculated value of 449 Hz.¹⁷

A tetrahedral, neutral form was unequivocally established as the predominant solution species for the cyanide, cyanate, and halide complexes in the phosphine series. Conductivity was near zero (Table II), the ^{107}Ag coupling constants approached the calculated sp^3 value of 227 Hz (Table V), and the established equilibria (3) require ion binding. Similar arguments apply to the cyanide, halide, and thiocyanate complexes in the phosphite series.

Intermediate in character were the trifluoroacetate, octahydridotriborate, and nitrate derivatives. The relatively large conductivities and coupling constants (Tables I and V) for these complexes suggest at least partial dissociation to $\text{L}_3\text{Ag}^+\text{X}^-$. However, the conductivities are not relevant because of the disproportionation these derivatives undergo (*vide infra*). In the case of the S_2PF_2^- derivative, the conductivity was quite low and the ^{19}F and ^{31}P nmr data for the anion were comparable to those for bound, unidentate S_2PF_2^- in transition metal complexes.¹⁹

Unexpectedly, some L_3AgX species show extensive disproportionation, even at very low temperatures.



Estimates for the equilibrium constants were obtained by intensity measurements from the low-temperature ^{31}P nmr spectra and are given in Table VI. There was

Table VI. Equilibrium Constants for the Reaction $2\text{L}_3\text{AgX} \rightleftharpoons \text{L}_4\text{AgX} + \text{L}_2\text{AgX}$

	Phosphite (-90°) ^a	Phosphine (-86°) ^b
NO_3^-	0.1	0.15
ClO_4^-	0.06	$<10^{-4}$ (not detected)
OOCFF_3^-		0.40
B_3H_8		0.64

^a Dichloromethane-toluene in 8:2 volume ratio. ^b Dichloromethane.

no nmr evidence of disproportionation for any L_3AgX complexes in which the counterion is tightly bound, *e.g.*, cyanide and halide.²⁰


(17) The calculated value, simply an estimate of s character in the Ag-P bond, was not adjusted for changes in the effective nuclear charge on the metal. The model is crude but seems applicable to these cases. A similar relationship is not found for the zerovalent platinum-phosphine complexes.¹⁸ In the platinum complexes, there is a significant change in color with change in coordination number, indicating large changes in ΔE , the mean excitation energy (average energy of all triplet excited states interacting with the ground state). ΔE is an explicit term in the expression for the coupling constant. If ΔE changes substantially, then linear correlation of J and s character may not follow. There was no evidence of color change with coordination number change in the silver series.

(18) D. H. Gerlach, A. R. Kane, G. W. Parshall, J. P. Jesson, and E. L. Muetterties, *J. Amer. Chem. Soc.*, **93**, 3543 (1971).

(19) F. N. Tebbe and E. L. Muetterties, *Inorg. Chem.*, **9**, 629 (1970).

(20) A trigonal L_3Ag^+ complex is coordinately unsaturated. Some energy reduction in solution may be achieved by formation of the coordinately saturated $\text{L}_4\text{Ag}^+\text{X}^-$ especially if in the coformed L_2AgX species the counteranion is bound. In fact, the counteranion appears to be bound in L_2AgX where X is nitrate, trifluoroacetate, and octahydridotriborate. These anions are potentially bidentate; thus L_2AgX could also be coordinately saturated in these derivatives.

All L_3AgX complexes are kinetically labile. Phosphorus ligand exchange was fast on the nmr time scale above -70 to -50° . In the phosphine series, the ligand exchange rates for the neutral, tetrahedral complexes are just slightly higher than that for the L_4Ag^+ complexes with NO_3^- , PF_6^- , and ClO_4^- counteranions (Table IV). Lower rates were found for the cationic trigonal $[(\text{C}_6\text{H}_5)_3\text{P}]_3\text{Ag}^+\text{PF}_6^-$ complex. These rates for the coordinately unsaturated $\text{L}_3\text{Ag}^+\text{X}^-$ complex are also lower than those for the coordinately saturated $\text{L}_4\text{Ag}^+\text{X}^-$ complexes. However, this expected rate reduction is small in contrast to the relatively large difference between L_4Pt and L_3Pt .¹⁸ A dissociative process is the only plausible mechanism for ligand exchange in the coordinately saturated L_3AgX complexes. Ligand exchange in the ionic set may be more complex since there is the possibility of an $\text{S}_\text{N}2$ contribution (6).²¹



Variation in ligand lability for the phosphite system paralleled that for the phosphines. Rate of ligand exchange decreased in the series $\text{CN} > \text{Cl} \sim \text{Br} > \text{SCN} > \text{NO}_3 > \text{ClO}_4$.

Trigonal and Linear Bis(phosphine)- and Bis(phosphite)-Silver(I) Complexes

Nmr parameters for the L_2AgX complexes are listed in Table VII. The estimated¹⁷ coupling constants for

Table VII. ^{31}P Nmr Parameters for $[(\text{C}_6\text{H}_5)_3\text{P}]_2\text{AgX}$ and $[(\text{C}_2\text{H}_5\text{O})_3\text{P}]_2\text{AgX}$ Complexes

X	$J_{^{107}\text{Ag-P}}$	δ^c	$J_{^{107}\text{Ag-P}}$	δ^c
F	450	28.1		
Cl	378	21.4	550	154.3
Br			536	151.5
CN	278	23.8		
B_3H_8	390	29.3		
S_2PF_2	410	27.4		
OOCFF_3	451	29.5		
NO_3	470	29.8	680	149.5
PF_6	496	33.4		
ClO_4	503	33.1	756	143.6
$\text{B}_3\text{H}_7\text{S}$	420	33.9		


^a Parameters were obtained at -80° . The solvent was dichloromethane. ^b Parameters were obtained at -100° . The solvent was an 80/20 dichloromethane-toluene mixture. ^c Chemical shifts are reported in parts per million with reference to an internal capillary of $(\text{C}_6\text{H}_5)_3\text{P}$; tri-*p*-tolylphosphine is at 11.1 ppm and triethyl phosphite at 159.9 ppm.

sp linear L_2Ag^+ complexes are 452 and 676 Hz for phosphine and phosphite ligands, respectively; and the nmr data suggest that the PF_6^- , BF_4^- , and ClO_4^- derivatives are ionic $\text{L}_2\text{Ag}^+\text{X}^-$ complexes. This is consistent with the high conductivities of these derivatives in dichloromethane (Table II).

In the phosphine series, the halide and pseudohalide complexes are neutral L_2AgX complexes in solution at

(21) The high exchange rates observed for ionic $\text{L}_3\text{Ag}^+\text{X}^-$ complexes are probably not due to very small concentrations of free ligand. We found nmr line-shape changes with temperature to be virtually identical for solutions of $\text{L}_3.0\text{Ag}^+\text{X}^-$ and $\text{L}_2.8\text{Ag}^+\text{X}^-$ (synthetically prepared from $\text{L}_2\text{AgX} + 0.8\text{L}$).

low temperature. The coupling constants were low compared to the linear sp estimate;¹⁷ and conductivities were very low (Tables II and VII). Molecular weight data taken at 25° indicated that these complexes are monomeric with *possibly* some ligand dissociation. A number of derivatives had coupling constants that fell in the range of 380–470 Hz. Conductivities of these derivatives were measurable but low (Table II). Thus, it would appear that some derivatives in this intermediate group dissociate slightly in solution (7).



Some neutral L_2AgX complexes may be tetrahedral complexes with a bidentate anion, *e.g.*, CF_3COO^- , NO_3^- , $B_3H_8^-$, and $S_2PF_2^-$.^{20,22} Nmr evidence supports this hypothesis for $S_2PF_2^-$. The δ_{19F} and J_{PF} values for $[(C_7H_7)_3P]_2AgS_2PF_2$ are close to the characteristic values¹⁹ for the ion bound through both sulfur atoms, to give a tetrahedral, neutral complex.

The bisphosphite complexes of silver(I) were not extensively examined, but their behavior generally parallels that of the phosphine analogs. The perchlorate derivative is best characterized as $L_2Ag^+ClO_4^-$ and the halide and cyanide derivatives as neutral complexes based on the observed coupling constants and conductivities (Tables II and VII).

The ligands are labile in the trigonal L_2AgX and linear L_2Ag^+ complexes. In the phosphine and phosphite series, the neutral L_2AgX complexes undergo ligand exchange at a rate (Table IV) comparable to those of the tetrahedral L_3AgX complexes. It is rather surprising that the rates for the bromide and chloride derivatives are so similar to those for the L_3AgX and L_2AgX species in both the phosphine and phosphite series. The linear L_2Ag^+ complexes are kinetically the most stable species, yet the rate differences between these coordinately deficient molecules and L_3Ag^+ are small (Table IV) compared to differences found in the platinum series.¹⁸

Solution Structure at 25°

Because of the kinetic lability of the silver–phosphorus bond in the phosphine and phosphite systems, the nmr spectra provide no information concerning solution structure or composition of the silver complexes at 25°. All silver phosphite and phosphine complexes show an averaged chemical shift that is temperature dependent, whereas the free phosphorus ligands have essentially temperature-independent chemical shifts. The temperature dependence of chemical shift for the complexes could be taken as evidence for significant dissociation of ligand. This may be the case for $L_2Ag^+X^-$ species and perhaps other specific L_3Ag^+ or L_3AgX complexes. However, L_3AgNO_3 cannot be significantly dissociated to L_2AgNO_3 because the conductivity of the former at 25° is substantially higher than that of L_2AgNO_3 . Molecular-weight studies of the

complexes suggested that dissociation was a common phenomenon. However, we are reluctant to interpret these data since the concentrations employed for the nmr and the molecular-weight studies are so different as to make comparative conclusions virtually meaningless. Also, where there is the possibility of ion dissociation as well as neutral ligand dissociation interpretation of “molecular weight” data is somewhat speculative.

Experimental Section

Nmr Studies. The ^{31}P spectra were obtained on a Varian Associates HA-100 with a variable temperature controller at 40.5 MHz. The mode of operation employed a homonuclear lock on triethylphosphine with random noise decoupling of the protons and use of the CAT. The ^{19}F spectra were obtained on the HA-100 at 94.1 MHz with a trichlorofluoromethane lock and on the HR-60 at 56.4 MHz with trichlorofluoromethane internal reference. The temperature readings were accurate to $\pm 6^\circ$. The results are tabulated in Tables I, V, and VII.

Conductivity Studies. The conductivities were measured on dichloromethane solutions at concentrations of 10^{-2} and 10^{-3} M. An Industrial Instruments conductivity bridge Model RC16B2 and a cell of constant 1.0 were used. The measurements were done at 25, –30, and -78° . Low-temperature determinations were carried out in a dry nitrogen atmosphere (Vacuum Atmospheres Dribox Model Dri-Lab He-43).

Molecular Weight Studies. Molecular weight studies were done in a dry nitrogen atmosphere. Ethanol was removed from chloroform for the vapor pressure osmometry measurements. The dichloromethane was Fisher Scientific and used as received. The freezing point determinations were carried out in benzene in a nitrogen atmosphere.

Preparations of Solutions of the Triethyl Phosphite–Silver Complexes. Since the triethyl phosphite–silver complexes generally melted below room temperature, they were not isolated. The commercial samples of the silver salts were analyzed prior to use. The triethyl phosphite was distilled in a nitrogen atmosphere prior to use. The solutions were prepared by adding the appropriately weighted amounts of the reactants to the solvent. All operations were effected in a dry nitrogen atmosphere (Vacuum Atmospheres Dribox).

Synthesis of Tri-*p*-tolylphosphine–Silver Complexes. The tri-*p*-tolylphosphine–silver complexes were isolatable, colorless, crystalline solids except for the tetrakis(tri-*p*-tolylphosphino)silver halides and the tris- and tetrakis(tri-*p*-tolylphosphino)silver difluorodithiophosphates which existed only in solution. These were prepared in solution by adding the appropriate amount of the phosphine to the $[(C_7H_7)_3P]_2AgX$ and $[(C_7H_7)_3P]_2AgS_2PF_2$ complexes.

$[(C_7H_7)_3P]_2AgF$. Silver fluoride (1 g, 7.9×10^{-2} mol) was dissolved in 25 ml of water and a solution of 4.7 g (1.5×10^{-2} mol) of tri-*p*-tolylphosphine in 75 ml of acetone was added. During this time the solution became brown. The reaction mixture was stirred 30 min, filtered, and evaporated to dryness. The glass-like solid was dissolved in acetonitrile and filtered to remove metallic silver and the filtrate evaporated to dryness to give a brown gum. Diethyl ether was slurried with the gum. The gum dissolved and a colorless solid separated, mp 162–166° dec.

Anal. Calcd for $C_{42}H_{42}P_2F$: C, 68.6; H, 5.8; F, 2.6; P, 8.4. Found: C, 66.0; H, 6.1; F, 2.6; P, 7.8.

Obtaining a pure compound proved difficult since in a competing reaction tri-*p*-tolylphosphine reduced Ag(I) to Ag(0) in solution. A pure $[(C_7H_7)_3P]_3AgF$ was not obtained. The nmr parameters were obtained from nmr solutions of mixtures of the bis and tris complexes.

$[(C_7H_7)_3P]_3AgCl$ and $[(C_7H_7)_3P]_3AgCl$. These complexes were prepared from silver chloride and tri-*p*-tolylphosphine in acetonitrile or chloroform using 1:2 and 1:3.5 molar ratios of reactants. The reaction solutions were evaporated to dryness. Recrystallization of crude L_2AgCl and L_3AgCl from acetone gave pure colorless crystals.

Anal. Calcd for $C_{42}H_{42}P_3Cl$: C, 67.1; H, 5.6; mol wt, 752.5. Found: C, 67.6; H, 5.7; mol wt VPO ($CHCl_3$), 662; VPO (CH_2Cl_2), 413; cryoscopic (C_6H_6), 792; mp 188–190°. Calcd for $C_{63}H_{63}P_3Cl$: C, 71.6; H, 6.0; Cl, 3.4; Ag, 10.2; mol wt, 1056. Found: C, 71.4; H, 6.1; Cl, 3.3; Ag, 10.3; mol wt VPO ($CHCl_3$), 540; VPO (CH_2Cl_2) 522; cryoscopic (C_6H_6), 498; mp 154–156°.

(22) If this is the case for the triborohydride ion, then the phosphorus atom environments would be inequivalent and this should be evident in the ^{31}P nmr. An AB ^{31}P pattern was observed below -70° for the copper analog;^{21,8} and the phosphorus environments equilibrated through a fast intramolecular rearrangement of the CuB_3 framework above -70° . There was no evidence of phosphorus atom inequivalence in $[(C_7H_7)_3P]_2AgB_3H_8$ at temperatures down to -100° . However, we find it difficult to visualize the $B_3H_8^-$ ligand being anything but a bifunctional moiety in this case and suggest that the absence of ^{31}P inequivalence under the conditions of our experiments simply reflects a very rapid rearrangement process.

$[(C_7H_7)_3P]_2AgBr$. Silver bromide (1.88 g, 1×10^{-2} mol) was added to a slurry of 6.08 g of tri-*p*-tolylphosphine (2×10^{-2} mol) in 100 ml of acetonitrile. The reaction mixture was heated at reflux ~ 6 hr until almost all the AgBr dissolved and then filtered. On cooling, colorless crystals appeared, which were recrystallized from acetone, mp 188–189°.

Anal. Calcd for $C_{42}H_{42}P_2BrAg$: C, 63.3; H, 5.3; P, 7.8; Br, 10.0. Found: C, 62.9; H, 5.5; P, 7.3; Br, 9.8.

$[(C_7H_7)_3P]_3AgBr$. The reaction was run as above but with 3 equiv of $(C_7H_7)_3P$ per equivalent of silver bromide. The colorless solid was recrystallized from acetone, mp 176–177°.

Anal. Calcd for $C_{63}H_{63}P_3BrAg$: C, 68.7; H, 5.8; Br, 7.3; Ag, 9.8. Found: C, 68.1; H, 5.8; Br, 8.4; Ag, 9.9.

$(C_7H_7)_3PAgI$ and $[(C_7H_7)_3P]_3AgI$. Silver iodide (2.34 g, 1×10^{-2} mol) was added to a solution of 6.08 g (2×10^{-2} mol) of tri-*p*-tolylphosphine in 40 ml of toluene. The reaction mixture was stirred at room temperature about 3 hr. During this time the silver iodide dissolved and a colorless solid formed. The reaction was filtered and the colorless solid was recrystallized from dichloromethane to give $LAgl$, mp 263–265°.

The filtrate from the above reaction was evaporated to dryness and the colorless solid recrystallized from acetone to give L_3AgI , mp 156–158°.

Anal. Calcd for $C_{21}H_{21}PIAg$: C, 46.8; H, 3.9. Found: C, 45.9; H, 4.1. Calcd for $C_{63}H_{63}P_3IAg$: C, 65.9; H, 5.5; Ag, 9.4. Found: C, 65.7; H, 5.4; Ag, 10.0.

No L_2AgI was isolated from the reaction.

$[(C_7H_7)_3P]_2AgCN$ and $[(C_7H_7)_3P]_3AgCN$. Silver cyanide (2.68 g, 2×10^{-2} mol) was added to a solution of 12.2 g (4×10^{-2} mol) of tri-*p*-tolylphosphine in 100 ml of toluene. The silver cyanide immediately dissolved. The solution was evaporated to dryness, and the colorless solid was recrystallized from acetone, mp 154–156°. The infrared CN stretching frequency was ~ 2110 cm^{-1} .

Anal. Calcd for $C_{43}H_{43}NP_2Ag$: C, 69.5; H, 5.7; N, 1.9; P, 8.3; Ag, 14.5; mol wt, 742.6. Found: C, 69.5; H, 5.7; N, 2.0; P, 8.2; Ag, 14.6; mol wt VPO (CH_2Cl_2), 447.

The L_3AgCN complex was synthesized similarly using 3 equiv of $(C_7H_7)_3P$ per AgCN. The colorless solid was recrystallized from diethyl ether, mp 128–133°. The CN stretch in the ir (Nujol spectrum) was at ~ 2110 cm^{-1} .

Anal. Calcd for $C_{64}H_{64}NP_3Ag$: C, 73.4; H, 6.1; N, 1.3; P, 8.9. Found: C, 71.3; H, 6.0; N, 1.8; P, 8.8.

$[(C_7H_7)_3P]_2AgB_3H_8$. Synthesis procedure was that of Muetterties, *et al.*³ The product is light sensitive.

$[(C_7H_7)_3P]_2AgS_2PF_6$. The reaction was run in a dry nitrogen atmosphere. Tri-*p*-tolylphosphine (6 g, 2×10^{-2} mol) was dissolved in 130 ml of acetonitrile and a solution of 2.4 g (1×10^{-2} mol) of AgS_2PF_6 dissolved in 25 ml of acetonitrile was added. A colorless solid precipitated which was isolated by filtration. The solid was recrystallized from toluene, mp 198–205°.

Anal. Calcd for $C_{42}H_{42}F_3S_2P_3Ag$: C, 59.3; H, 5.0; F, 4.5; P, 10.9; Ag, 12.7. Found: C, 59.7; H, 5.1; F, 4.8; P, 11.2; Ag, 12.3.

$[(C_7H_7)_3P]_3AgS_2PF_6$. Attempts to isolate $[(C_7H_7)_3P]_3AgS_2PF_6$ by the procedure above but using 3 equiv of tri-*p*-tolylphosphine per equivalent of AgS_2PF_6 gave only $[(C_7H_7)_3P]_2AgS_2PF_6$. A solution of $[(C_7H_7)_3P]_3AgS_2PF_6$ was prepared by dissolving equivalent amounts of $(C_7H_7)_3P$ and $[(C_7H_7)_3P]_2AgS_2PF_6$ in dichloromethane.

$[(C_7H_7)_3P]_2AgOCCF_3$ and $[(C_7H_7)_3P]_3AgOCCF_3$. To a solution of tri-*p*-tolylphosphine (6 g, 2×10^{-2} mol) dissolved in

40 ml of toluene was added silver trifluoroacetate (2.2 g, 1×10^{-2} mol). The silver trifluoroacetate quickly dissolved. After stirring for 1 hr, the reaction mixture was filtered and the filtrate was evaporated to dryness to give a colorless solid which was recrystallized from acetone, mp 197–199°.

Anal. Calcd for $C_{44}H_{44}O_2F_3P_2Ag$: C, 63.7; H, 5.1; P, 7.5; F, 6.9; Ag, 13.0. Found: C, 63.4; H, 5.1; P, 7.6; F, 6.9; Ag, 13.0.

The synthesis of $[(C_7H_7)_3P]_3AgOCCF_3$ was carried out by the same procedure as above using 6 g (2×10^{-2} mol) of tri-*p*-tolylphosphine and 1.47 g (6.7×10^{-3} mol) of silver trifluoroacetate, mp 174–176°.

Anal. Calcd for $C_{65}H_{65}O_2F_3P_3Ag$: C, 68.9; H, 5.6; F, 5.0. Found: C, 67.7; H, 5.6; F, 5.4.

$[(C_7H_7)_3P]_2AgNO_3$, $[(C_7H_7)_3P]_3AgNO_3$, and $[(C_7H_7)_3P]_4AgNO_3$. The silver nitrate-tri-*p*-tolylphosphine complexes were prepared by dissolving 1 equiv of silver nitrate in acetonitrile and adding it to a slurry of 2, 3, or 4 equiv of $(C_7H_7)_3P$ in acetonitrile. The tri-*p*-tolylphosphine dissolved immediately on addition of the silver nitrate solution. All the complexes were recrystallized from acetone.

Anal. Calcd for $C_{42}H_{42}NO_3P_2Ag$: C, 64.8; H, 5.4; N, 1.8; P, 8.0; Ag, 13.9; mol wt, 778.6. Found: C, 64.7; H, 5.7; N, 1.8; P, 7.5; Ag, 13.9; mol wt VPO ($CHCl_3$), 805; VPO (CH_2Cl_2), 783; cryoscopic (C_6H_6), 439; mp 185° coalesces, 199–202° melts. Calcd for $C_{63}H_{63}NO_3P_3Ag$: C, 69.9; H, 5.9; N, 1.3; P, 8.6; Ag, 10.0; mol wt, 1082.9. Found: C, 69.3; H, 5.9; N, 1.2; P, 8.6; Ag, 9.5; mol wt VPO ($CHCl_3$), 595; cryoscopic (C_6H_6) 609; mp 193–196°. Calcd for $C_{84}H_{84}NO_3P_4Ag$: C, 72.7; H, 6.1; N, 1.0; mol wt, 1387.3. Found: C, 72.6; H, 6.2; N, 1.1; mol wt VPO ($CHCl_3$), 554; cryoscopic (C_6H_6), 415; mp 147° coalesces, 177–190° melts.

$[(C_7H_7)_3P]_2AgPF_6$, $[(C_7H_7)_3P]_3AgPF_6$, and $[(C_7H_7)_3P]_4AgPF_6$. The hexafluorophosphate complexes were obtained by metathesis of NH_4PF_6 with the appropriate tri-*p*-tolylphosphine-silver nitrate complex in an acetonitrile-water solution or ethanol-water solution. The recrystallization solvent was a diethyl ether-tetrahydrofuran mixture for L_2AgPF_6 , and acetone for L_3AgPF_6 and L_4AgPF_6 .

Anal. Calcd for $C_{42}H_{42}P_3F_6Ag$: C, 58.6; H, 4.9. Found: C, 60.2; H, 5.9; mp 229–234°. Calcd for $C_{63}H_{63}P_4F_6Ag$: C, 64.9; H, 5.4. Found: C, 64.0; H, 5.4; mp 190° coalesces, 225–226° melts. Calcd for $C_{84}H_{84}P_5F_6Ag$: C, 68.6; H, 5.8; P, 10.5. Found: C, 68.4; H, 5.8; P, 10.2; mp 256–258° cloudy, 262° clear.

$[(C_7H_7)_3P]_2AgClO_4$, $[(C_7H_7)_3P]_3AgClO_4$, and $[(C_7H_7)_3P]_4AgClO_4$. No attempt was made to isolate the perchlorate complexes because of possible instability. Solutions were prepared by adding the appropriate ratios of tri-*p*-tolylphosphine and silver perchlorate to dichloromethane.

$[(C_7H_7)_3P]_2AgB_3H_{12}S^8$ and $[(C_7H_7)_3P]_3AgB_3H_{12}S$. The $L_2AgB_3H_{12}S$ was synthesized according to Muetterties, *et al.*³ The $L_3AgB_3H_{12}S$ was not isolated but studied in solution by adding 1 equiv of tri-*p*-tolylphosphine to $L_2AgB_3H_{12}S$.

Acknowledgment. We would like to thank Mrs. Jean Read and Mr. Lou Walther for the ^{31}P spectra and Mr. W. G. Peet for preparing solutions of the triethyl phosphine-silver complexes.