

5-Oxazolonium Perchlorates from α -Acylamino-acids


By G. V. BOYD

(Department of Chemistry, Chelsea College of Science and Technology, London, S.W.3)

THE action of phosphorus halides on α -acylamino-acids¹ or of hydrogen halides on simple 5-oxazolones ("saturated azlactones")² results in the formation of highly reactive unstable compounds which have been extensively studied. The view¹ that they are 5-oxazolonium halides (I; X = Br or Cl) has been confirmed by cryoscopic measurements³ and i.r. spectroscopy;⁴ the presence of a strong absorption band near 1880 cm.⁻¹ is irreconcilable with the alternative acyl halide structure.

It has now been found that treatment of α -acylamino-acids with acetic anhydride and 70% perchloric acid at or below room temperature

gives high yields of the corresponding 5-oxazolonium perchlorates:



Thus, hippuric acid affords the salt (I; R¹ = Ph, R² = H, X = ClO₄), 89%, m.p. 171° (decomp.)[†]

[†] Satisfactory analytical and spectroscopic data have been obtained for all new compounds whose melting points are given.

ν_{\max} 1880 cm^{-1} , which is stable when kept dry. It rapidly reacts at room temperature with water,

perchlorates (III) are readily prepared from *N*-alkyl- and *N*-aryl- α -acylamino-acids:

N-Substituted 5-oxazolonium perchlorates (III)

Compound	R ¹	R ²	Yield (%)	M.p. (decomp.)	ν_{\max} (cm^{-1})	Chemical shift of CH ₂ -protons (τ) in CF ₃ ·CO ₂ H
(a)	Ph	Ph	85	163°	1890	4.61
(b)	Ph	Me	82	167	1880	4.90
(c)	Me	Ph	90	143	1880	4.78
(d)	<i>p</i> -NO ₂ C ₆ H ₄	Ph	100	170	1887	5.01


methanol, aniline, and sodium glycinate to yield, respectively, hippuric acid, methyl hippurate, hippuranilide, and benzoylglycylglycine. It is deprotonated by triethylamine with the formation of 2-phenyloxazol-5-one (II; R¹ = Ph, R² = H), from which perchloric acid regenerates the salt; cold aqueous sodium carbonate produces a mixture of the oxazolone and sodium hippurate. The latter does not arise from the azlactone since this is stable under these conditions. The perchlorate condenses with benzaldehyde in hot acetonitrile to yield the bright yellow benzylidene derivative (I; R¹ = Ph, R² = :CHPh, X = ClO₄), m.p. 198° (decomp.), which gives the labile geometrical isomer⁵ of 4-benzylidene-2-phenyloxazol-5-one on treatment with water.

Cyclisation of *N*-benzoyl-L-phenylalanine gave the optically active salt (I; R¹ = Ph, R² = CH₂Ph, X = ClO₄), 76%, m.p. 132° (decomp.), i.r. 1880 cm^{-1} , $[\alpha]_D^{25}$ -37.8° (*c* 2.4, acetonitrile), from which the optically active oxazolone (II; R¹ = Ph, R² = CH₂Ph), $[\alpha]_D^{25}$ -70.3° (*c* 0.37, dioxan) was obtained.[†] Water hydrolysed the salt to give the parent acid of 94% optical purity.

The formation of an optically active oxazolonium salt suggests strongly that these compounds exist in the Δ^2 -form (I) rather than as the Δ^3 -tautomers (Ia); conclusive evidence for structure (I) is provided by the n.m.r. spectra of the two salts: the first shows a methylene singlet at τ 4.98, and the second a methine triplet at 4.51 and a benzylic methylene doublet at 6.37. α -Acetaminophenylacetic acid was then used in the synthesis as it was thought that the 4-phenyl substituent might stabilise structure (Ia). However, the resulting oily perchlorate, i.r. 1885 cm^{-1} , is clearly (I; R¹ = Me, R² = Ph, X = ClO₄) since its n.m.r. spectrum exhibits a methyl singlet (τ 7.30) and a methine doublet (due to coupling with N⁺H) at 4.15.

N-Substituted oxazolonium salts have not been reported previously; the colourless hygroscopic

Some of the salts are listed in the Table. They all show the characteristic high-frequency i.r. absorption, the appearance of methylene signals again confirms the Δ^2 -structure (III). These salts are of current interest because they can be deprotonated to give mesoionic oxazolones, the "münchnones" (IV),⁶ for which they are con-



venient precursors. The mesoionic compounds are too unstable to be isolated but their formation is manifest by the appearance of yellow or orange colours and they can be trapped by suitable reagents. Thus, when the diphenyl-compound (IIIa) is treated with triethylamine, or solid sodium carbonate, in the presence of trifluoroacetic anhydride the known⁷ stable trifluoroacetyl-derivative (V) is obtained in 90% yield; dimethyl acetylenedicarboxylate yields the pyrrole (VI), 80%, m.p. 122.5°, by cycloaddition and subsequent loss of carbon dioxide,⁸ and benzylamine gives the benzylamide of *N*-benzoyl-*N*-phenylglycine, 88%, m.p. 162°. The perchlorates (III) are instantly decomposed by cold water to the parent acylamino-acids without change of colour, but hydrolysis with aqueous sodium carbonate may proceed wholly or partly *via* the münchnones since transient colours are observed. *N*-*p*-Nitrobenzoyl-L-proline⁸ yields an oily optically inactive perchlorate, i.r. 1890 cm^{-1} , which gives the (\pm)-acid on treatment with water.

[†] The highest rotation recorded (M. Goodman and L. Levine, *J. Amer. Chem. Soc.*, 1964, **86**, 2918.) for this azlactone is -71.2°.

The acetic anhydride-perchloric acid method when applied to 1-carboxymethyl-2-pyridone afforded the exceptionally stable condensed oxazononium salt (VII), 94%, colourless prisms,

m.p. 179° (decomp.), i.r. 1890 cm.⁻¹, CH₂-singlet at τ 4.35.

(Received, August 19th, 1968; Com. 1148.)

¹ J. W. Cornforth, in "The Chemistry of Penicillin", ed. H. T. Clarke, Princeton University Press, Princeton, 1949, pp. 731, 746.

² E. Mohr and F. Stroschein, *Ber.*, 1909, **42**, 2521.

³ J. L. O'Brien and C. Niemann, *J. Amer. Chem. Soc.*, 1957, **79**, 80.

⁴ C. W. Smith and R. S. Rasmussen, *J. Amer. Chem. Soc.*, 1949, **71**, 1080; H. E. Carter and J. W. Hinman, *J. Biol. Chem.*, 1949, **178**, 403.

⁵ K. Brocklehurst, H. S. Price and K. Williamson, *Chem. Comm.*, 1968, 884.

⁶ R. Huisgen, in "Aromaticity", *Chem. Soc. Special Publ. No. 21*, 1967, p. 51; R. Huisen, H. Gotthardt, H. O. Bayer, and F. C. Schaefer, *Angew. Chem.*, 1964, **76**, 185; R. Huisen, E. Funker, F. C. Schaefer, and R. Knorr, *ibid.*, 1967, **79**, 321.

⁷ G. Singh and S. Singh, *Tetrahedron Letters*, 1964, 3789.

⁸ M. W. Williams and G. T. Young, *J. Chem. Soc.*, 1964, 3701.