

Fibrin-Stabilizing Factor Inhibitors. 12. 5-Dibenzylaminopentylamine and Related Compounds, a New Type of FSF Inhibitors

Kurt-Jürgen Hoffmann, Pål Stenberg, Christine Ljunggren, Uno Svensson, J. Lars G. Nilsson,*

Faculty of Pharmacy, University of Uppsala, S-751 23 Uppsala, Sweden

Olle Eriksson, Ann Hartkoorn, and Ragnar Lundén

AB Kabi, Fack, S-104 25 Stockholm 30, Sweden. Received September 24, 1974

A series of ω -dibenzylaminoalkylamines and related compounds have been prepared and tested as inhibitors of fibrin cross-linking. This structural type was chosen in an attempt to develop noncompetitive inhibitors of fibrinolytic. By the combination of the dibenzylamino moiety at one end and the primary amino group at the other end of a polymethylene chain, the same compound could function both as a pseudo donor substrate and as a noncompetitive alkylating inhibitor. Some of the compounds, notably 74–79, are among the most active fibrinolytic inhibitors described. However, the data indicate that the compounds probably function only as pseudo donor inhibitors.

The final step in the formation of a fibrin clot is the stabilization of the protein, *i.e.*, the cross-linking of the fibrin molecules. This reaction is a transamidation, catalyzed by the thrombin and Ca^{2+} activated fibrin-stabilizing factor (FSF; F XIIIa) or fibrinolytic. The inhibition of this reaction may lead to development of anticoagulants. We have therefore considered it worthwhile to explore the possibilities for therapeutic use of such inhibitors in various coagulation diseases.


A large number of compounds have already been tested *in vitro* as potential inhibitors of fibrin cross-linking.^{1,2} These inhibitors may be considered as pseudo substrates for the enzyme, functioning either as "donors" or "acceptors" in the transamidation reaction.³ Monodansylcadaverine¹ is the most studied inhibitor of the donor type while recently reported thiol esters^{4,5} function as acceptor substrates.

We have previously reported⁶ that an inhibitor of the donor type should have the general structure $\text{ArX}(\text{CH}_2)_5\text{NH}_2$, where the X group is an electron-withdrawing moiety, *e.g.*, a sulfonyl group imparting a relative low electron density to the aromatic nucleus. It has been proposed² that a pseudo donor of this type is aligned in the proximity of the active center of the enzyme *via* a charge-transfer type complex between the aromatic part of the substrate and the tryptophan residue adjacent to the active site cysteine of the enzyme. The side chain carrying the nucleophilic amino group must then have a specific length (preferably five methylene groups) to permit the amino group to attack the thiol ester of the acyl-enzyme intermediate and form an amide bond between the fibrin and the inhibitor, thus blocking the cross-linking of fibrin.

We now wish to report on a new structural type of fibrinolytic inhibitors. We have found that primary *n*-alkylamines, particularly *n*-pentylamines carrying an aryl-methylamino group at the other end of the polymethylene chain, are highly active as inhibitors of fibrin cross-linking. Similar to the sulfonamides, the specificity of these compounds is based on the structure of the aromatic moiety of the molecule. The inhibitors and their activities are presented in Table II and intermediates used in the synthesis are listed in Table I.

Chemistry. The compounds presented in Table II were prepared using mainly four different methods designated A–D in Scheme I. Most of the compounds tested were prepared by alkylation of the appropriate secondary amine, in most cases a dibenzylamine. These amines (compounds 1–


Scheme I



13) were synthesized *via* two different routes as shown in Scheme I.

* Address correspondence to this author at AB Hässle, Fack, S-431 20 Mölndal 1, Sweden.

Table I. Physical Data for Nitriles and Phthalimido Derivatives

Compd no.	R ₁	R ₂	n	R ₃	R ₁ R ₂ N(CH ₂) _n R ₃	Yield, %	Mp and/or bp (mm), °C	Formula	Recrystn solvent
14	β -C ₁₀ H ₇ CH ₂ ⁻	CH ₃ ⁻	4	-CN	C ₁₇ H ₂₀ N ₂	80	a	C ₁₇ H ₂₀ N ₂	2-Propanol
15	C ₆ H ₅ ⁻	C ₃ H ₅ ⁻	4	-CN	C ₁₃ H ₁₈ N ₂	30	145 (0.7)	C ₁₃ H ₁₈ N ₂	Absolute ethanol-ether
16	C ₆ H ₅ CH ₂ ⁻	C ₆ H ₅ CH ₂ ⁻	2	-CN	C ₁₇ H ₁₈ N ₂	64	152-155 (0.01), 45-46	C ₁₇ H ₁₈ N ₂	2-Propanol
17	C ₆ H ₅ CH ₂ ⁻	C ₆ H ₅ CH ₂ ⁻	3	-CN	C ₁₈ H ₂₀ N ₂	64	167-169 (0.01), 45-49	C ₁₈ H ₂₀ N ₂	Water
18	C ₆ H ₅ CH ₂ ⁻	C ₆ H ₅ CH ₂ ⁻	4	-CN	C ₁₉ H ₂₂ N ₂	77	170-174 (0.5)	C ₁₉ H ₂₂ N ₂	Acetone
19	C ₆ H ₅ CH ₂ ⁻	C ₆ H ₅ CH ₂ ⁻	5	-CN	C ₂₀ H ₂₄ N ₂	65	180-182 (0.01), 40-41	C ₂₀ H ₂₄ N ₂	Absolute ethanol-ether
20	C ₆ H ₅ CH ₂ ⁻	C ₆ H ₅ CH ₂ ⁻	6	-CN	C ₂₁ H ₂₆ N ₂	70	185-187 (0.01)	C ₂₁ H ₂₆ N ₂	Absolute ethanol
21	CH ₂ =CHCH ₂ ⁻	CH ₂ =CHCH ₂ ⁻	4	-CN	C ₁₁ H ₁₈ N ₂	70	135-136 (15)	C ₁₁ H ₁₈ N ₂	Toluene
22	C ₃ H ₇ ⁻	C ₃ H ₇ ⁻	4	-CN	C ₁₁ H ₂₂ N ₂	88	72-75 (1)	C ₁₁ H ₂₂ N ₂	Toluene
23	C ₆ H ₅ CH ₂ ⁻	HOCH ₂ CH ₂ ⁻	4	-CN	C ₁₄ H ₂₀ N ₂ O	80	153-155 (0.05)	C ₁₄ H ₂₀ N ₂ O	Petroleum ether
24	4-ClC ₆ H ₄ CH ₂ ⁻	C ₆ H ₅ CH ₂ ⁻	4	-CN	C ₁₃ H ₂₁ ClN ₂ •C ₂ H ₅ O ₄	38	120-122	C ₁₃ H ₂₁ ClN ₂ •C ₂ H ₅ O ₄	Ethanol-water
25	4-(CH ₃) ₂ N•C ₆ H ₅ CH ₂ ⁻	C ₆ H ₅ CH ₂ ⁻	4	-CN	C ₂₁ H ₂₇ N ₃ •2HCl	51	153-155	C ₂₁ H ₂₇ N ₃ •2HCl	2-Propanol
26	4-CH ₃ C ₆ H ₅ CH ₂ ⁻	C ₆ H ₅ CH ₂ ⁻	4	-CN	C ₂₀ H ₂₄ N ₂ •C ₂ H ₅ O ₄	45	126-128	C ₂₀ H ₂₄ N ₂ •C ₂ H ₅ O ₄	Water
27	4-CH ₃ OC ₆ H ₅ CH ₂ ⁻	C ₆ H ₅ CH ₂ ⁻	5	Phthalimido	C ₂₃ H ₃₀ N ₂ O ₃ •C ₂ H ₅ O ₄	55	232-233	C ₂₃ H ₃₀ N ₂ O ₃ •C ₂ H ₅ O ₄	Acetone
28	C ₆ H ₅ CH ₂ CH ₂ ⁻	C ₆ H ₅ CH ₂ CH ₂ ⁻	5	Phthalimido	C ₂₃ H ₃₂ N ₂ O ₂ •C ₂ H ₅ O ₄	63	142-145 dec	C ₂₃ H ₃₂ N ₂ O ₂ •C ₂ H ₅ O ₄	Absolute ethanol-ether
29	3,4-Cl ₂ C ₆ H ₄ CH ₂ ⁻	C ₆ H ₅ CH ₂ ⁻	4	-CN	C ₁₉ H ₂₀ Cl ₂ N ₂ •C ₂ H ₅ O ₄	49	238-239 dec	C ₁₉ H ₂₀ Cl ₂ N ₂ •C ₂ H ₅ O ₄	Absolute ethanol
30	4-CH ₃ CONHC ₆ H ₅ CH ₂ ⁻	C ₆ H ₅ CH ₂ ⁻	5	Phthalimido	C ₂₃ H ₃₁ N ₃ O ₃ •C ₂ H ₅ O ₄	68	212-214	C ₂₃ H ₃₁ N ₃ O ₃ •C ₂ H ₅ O ₄	Toluene
31	4-CNC ₆ H ₅ CH ₂ ⁻	C ₆ H ₅ CH ₂ ⁻	5	Phthalimido	C ₂₃ H ₂₇ N ₃ O ₂	45	68-69	C ₂₃ H ₂₇ N ₃ O ₂	Toluene
32	4-NO ₂ C ₆ H ₅ CH ₂ ⁻	C ₆ H ₅ CH ₂ ⁻	4	-CN	C ₂₇ H ₂₇ N ₃ O ₄	40	76-78	C ₂₇ H ₂₇ N ₃ O ₄	Petroleum ether
33	4-FC ₆ H ₅ CH ₂ ⁻	C ₆ H ₅ CH ₂ ⁻	4	-CN	C ₁₉ H ₂₁ FN ₂	58	a	C ₁₉ H ₂₁ FN ₂	Ethanol-water
34	α-C ₁₀ H ₇ CH ₂ ⁻	C ₆ H ₅ CH ₂ ⁻	4	-CN	C ₂₅ H ₂₄ N ₂	69	a	C ₂₅ H ₂₄ N ₂	Absolute ethanol
35	4-BrC ₆ H ₅ CH ₂ ⁻	C ₆ H ₅ CH ₂ ⁻	5	Phthalimido	C ₂₇ H ₂₇ BrN ₂ O ₂	55	54-56	C ₂₇ H ₂₇ BrN ₂ O ₂	Petroleum ether
36	4-ClC ₆ H ₅ CH ₂ ⁻	4-ClC ₆ H ₅ CH ₂ ⁻	4	-CN	C ₁₉ H ₂₀ ClN ₂ •C ₂ H ₅ O ₄	59	132-133	C ₁₉ H ₂₀ ClN ₂ •C ₂ H ₅ O ₄	Ethanol-water
37					C ₁₉ H ₂₀ N ₂ •HCl	40	230-233	C ₁₉ H ₂₀ N ₂ •HCl	Absolute ethanol
38	HC≡CCH ₂ CH ₂ ⁻	phthalimido			C ₁₂ H ₉ NO ₂	88	134-135	C ₁₂ H ₉ NO ₂	Glacial acetic acid
39	(C ₆ H ₅ CH ₂) ₂ NCH ₂ C≡CCH ₂ CH ₂ ⁻	phthalimido			C ₂₇ H ₂₄ N ₂ O ₂	50	85-86	C ₂₇ H ₂₄ N ₂ O ₂	70% ethanol
40	(C ₆ H ₅ CH ₂) ₂ NCH ₂ CH ₂ OCH ₂ CN				C ₁₈ H ₂₀ N ₂ O•HCl	58	154-155	C ₁₈ H ₂₀ N ₂ O•HCl	Ethanol-ether
41	(C ₆ H ₅ CH ₂) ₂ NCO(CH ₂) ₄ ⁻	phthalimido			C ₂₇ H ₂₆ N ₂ O ₃	80	84-86	C ₂₇ H ₂₆ N ₂ O ₃	Ethyl acetate

^aOil which could not be distilled; hydrochloride and oxalate not crystalline. ^bM. V. Gittos and N. Wilson, *J. Chem. Soc.*, 2371 (1955).

Table II. Physical Data and Inhibitory Activities of the Compounds Tested^a

Compd no.	R ₁	R ₂	X	Method	Yield, %	Mp and/or bp (mm), °C	Formula	Derivative	Activity ^b
R ₁ R ₂ NXNH ₂									
Monodansylcadaverine									
42	C ₂ H ₅ -	C ₂ H ₅ -	-(CH ₂) ₅ -	d	68	90-93 (10)	C ₃ H ₂₂ N ₂	Base	100
43	C ₆ H ₅ CH ₂ -	H-	-(CH ₂) ₄ -	C	64	125-127 (5)	C ₁₁ H ₁₈ N ₂	Base	1
44	C ₆ H ₅ CH ₂ -	H-	-(CH ₂) ₅ -	C	66	110-112 (0.4)	C ₁₂ H ₂₀ N ₂	Base	5
45	C ₆ H ₅ CH ₂ -	H-	-(CH ₂) ₆ -	C	69	136-140 (1)	C ₁₃ H ₂₂ N ₂	Base	6
46	C ₆ H ₅ CH ₂ -	CH ₃ -	-(CH ₂) ₅ -	d		114-115 (1.5)	C ₁₃ H ₂₂ N ₂	Base	10
47	β-C ₁₀ H ₇ CH ₂ -	CH ₃ -	-(CH ₂) ₅ -	A	67	160-164 (0.8)	C ₁₇ H ₂₄ N ₂	Base	11
48	C ₆ H ₅ -	C ₂ H ₅ -	-(CH ₂) ₅ -	A	65	134 (1)	C ₁₃ H ₂₂ N ₂	Base	19
49	C ₆ H ₅ CH(CH ₃)-	H-	-(CH ₂) ₅ -	D	33	110-112 (0.7)	C ₁₃ H ₂₂ N ₂	Base	1
50	α-C ₁₀ H ₇ CH(CH ₃)-	H-	-(CH ₂) ₅ -	D	52	136-140 (0.3)	C ₁₇ H ₂₄ N ₂	Base	7
51	(C ₆ H ₅) ₂ CH-	H-	-(CH ₂) ₅ -	C	65	256-257	C ₁₈ H ₂₄ N ₂	Base	5
52	C ₆ H ₅ CH ₂ -	C ₆ H ₅ CH ₂ -	-(CH ₂) ₃ -	A	79	137-139 (0.01), 104-105	C ₁₇ H ₂₂ N ₂	2HCl	40
53	C ₆ H ₅ CH ₂ -	C ₆ H ₅ CH ₂ -	-(CH ₂) ₄ -	A	76	145-146 (0.01)	C ₁₈ H ₂₄ N ₂	2HCl	2
54	C ₆ H ₅ CH ₂ -	C ₆ H ₅ CH ₂ -	-(CH ₂) ₅ -	A	63	200-202	C ₁₉ H ₂₆ N ₂	2HCl	5
55	C ₆ H ₅ CH ₂ -	C ₆ H ₅ CH ₂ -	-(CH ₂) ₆ -	A	78	158-160 (0.01), 198-199	C ₂₀ H ₂₈ N ₂	HCl	110
56	C ₆ H ₅ CH ₂ -	C ₆ H ₅ CH ₂ -	-(CH ₂) ₅ CH ₂ CH ₂ CH ₂ CH ₂ -	A	80	146-148	C ₁₈ H ₂₄ N ₂ O	HCl	58
57	C ₆ H ₅ CH ₂ -	C ₆ H ₅ CH ₂ -	-(CH ₂) ₅ CH ₂ CH ₂ CH ₂ CH ₂ -	C	85	140-142	C ₁₈ H ₂₄ N ₂ S	0.5 Fumarate	3
58	C ₆ H ₅ CH ₂ -	C ₆ H ₅ CH ₂ -	-(CH ₂) ₅ C≡CCH ₂ CH ₂ -	B	67	156-158	C ₁₉ H ₂₂ N ₂	Fumarate	2
59	C ₆ H ₅ CH ₂ -	C ₆ H ₅ CH ₂ -	-COCH ₂ CH ₂ CH ₂ CH ₂ -	C	91	139-140	C ₁₈ H ₂₂ N ₂ OS	Fumarate	7
60	C ₆ H ₅ CH ₂ -	C ₆ H ₅ CH ₂ -	-COCH ₂ CH ₂ CH ₂ CH ₂ CH ₂ -	C	86	Oil	C ₁₉ H ₂₀ N ₂ O	Base	2
61	CH ₂ =CHCH ₂ -	CH ₃ =CHCH ₂ -	-(CH ₂) ₅ -	A	55	120-123 (20)	C ₁₄ H ₂₂ N ₂	Base	5
62	n-C ₃ H ₇ -	n-C ₃ H ₇ -	-(CH ₂) ₅ -	A	70	64-65 (0.5)	C ₁₁ H ₂₀ N ₂	Base	4
63	C ₆ H ₅ CH ₂ CH ₂ -	C ₆ H ₅ CH ₂ CH ₂ -	-(CH ₂) ₅ -	B	87	138-140	C ₂₁ H ₃₀ N ₂	Base	1
64	C ₆ H ₅ CH ₂ -	-CH ₂ CH ₂ OH	-(CH ₂) ₅ -	A	76	133-135 (0.1)	C ₁₄ H ₂₄ N ₂ O	Fumarate	40
65	4-NO ₂ C ₆ H ₄ CH ₂ -	C ₆ H ₅ CH ₂ -	-(CH ₂) ₅ -	B	75	152-153	C ₁₉ H ₂₅ N ₂ O ₂	Base	6
66	4-CNC ₆ H ₄ CH ₂ -	C ₆ H ₅ CH ₂ -	-(CH ₂) ₅ -	B	51	146-147	C ₂₀ H ₂₅ N ₃	Fumarate	51
67	4-HOCOC ₆ H ₄ CH ₂ -	C ₆ H ₅ CH ₂ -	-(CH ₂) ₅ -	C	37	202-205	C ₂₀ H ₂₆ N ₂ O ₂	2HCl	26
68	4-CH ₃ CONHC ₆ H ₄ CH ₂ -	C ₆ H ₅ CH ₂ -	-(CH ₂) ₅ -	B	86	Oil	C ₂₁ H ₂₉ N ₃ O	Base	43
69	4-C ₂ H ₅ NHC ₆ H ₄ CH ₂ -	C ₆ H ₅ CH ₂ -	-(CH ₂) ₅ -	C	51	Oil	C ₂₁ H ₃₁ N ₃	Base	30
70	4-CH ₃ OC ₆ H ₄ CH ₂ -	C ₆ H ₅ CH ₂ -	-(CH ₂) ₅ -	B	63	149-152	C ₂₀ H ₂₂ N ₂ I	2HCl	55
71	4-(CH ₃) ₂ NC ₆ H ₄ CH ₂ -	C ₆ H ₅ CH ₂ -	-(CH ₂) ₅ -	A	85	Oil	C ₂₁ H ₃₁ N ₃	Base	56
72	4-NH ₂ CH ₂ C ₆ H ₄ CH ₂ -	C ₆ H ₅ CH ₂ -	-(CH ₂) ₅ -	C	92	149-150	C ₂₀ H ₂₉ N ₃	Fumarate	40
73	4-CH ₃ C ₆ H ₄ CH ₂ -	C ₆ H ₅ CH ₂ -	-(CH ₂) ₅ -	A	77	175-177	C ₂₀ H ₂₈ N ₂	2HCl	92
74	4-FC ₆ H ₄ CH ₂ -	C ₆ H ₅ CH ₂ -	-(CH ₂) ₅ -	A	82	128-130	C ₁₉ H ₂₅ FN ₂	2HCl	105
75	4-ClC ₆ H ₄ CH ₂ -	C ₆ H ₅ CH ₂ -	-(CH ₂) ₅ -	A	83	177-179	C ₁₉ H ₂₅ ClN ₂	2HCl	150

chain (cf. compounds 52–55). A similar specificity for the chain length was observed for the dansyl derivatives previously studied.⁹ In the monobenzyl compounds 43–45 the length of the side chain is not very critical for the effect.

To study further the possibility that the benzyl compounds might function as alkylating inhibitors, as discussed above, compounds 61–63, 81, and 82 were synthesized. A diallylamine (61) may function in the same manner as the dibenzylamine⁸ and alkylate the attacking nucleophilic species, whereas the closely related dipropylamine derivative 62 or the diphenethylamine derivative 63 would not have these properties. Of these compounds, 61 and 62 have only low activity while 63 has an activity of 40 as compared to 110 for 5-dibenzylaminopentylamine (54). When the primary amino group of 54 is methylated to a tertiary amine as in 81 or when the dibenzylamino nitrogen is quaternized as in 82, all the activity is lost. The results thus indicate that the compounds probably are pseudo donors and not alkylating inhibitors as assumed in our working hypothesis. This is particularly well illustrated by compound 63 which should be inactive as an alkylating inhibitor and by 81 and 82 which both theoretically should be potent inhibitors.

The data also show that potent FSF inhibitors of the dibenzyl type should have a primary amino group, a pentamethylene chain, and a basic tertiary nitrogen substituted with preferentially two benzyl groups. The importance of the basic tertiary nitrogen is also illustrated by compound 60 which is an amide. This substance has only low activity.

In a previous study we demonstrated¹⁰ that monodansyl derivatives of 3-oxacadaverine and of 3-thiacadaverine were considerably more active inhibitors of fibrin cross-linking than monodansylcadaverine itself. This is probably due to the low basic strength of the primary amine in these compounds. The low basicity would increase the amount of the primary amine present in the base form at physiological pH and as a consequence also increase the activity of the compound when it functions as a nucleophile in the reaction with the acyl-enzyme intermediate. When we tested the same idea for the dibenzyl compounds 56–59 these inhibitors had only low activity. This may indicate that the specificity of the dansyl compounds is accomplished in a different way than that for the dibenzyl derivatives. The latter compounds could be bound to the receptor structure of the enzyme in a different way than the dansyl compounds.² An electron-rich structure, an oxygen (56), a sulfur (57), or a triple bond (58) in the chain of the dibenzyl derivatives may bind to some macromolecular functional group at the receptor and divert the nucleophilic H₂N group from reacting with the thiol ester of the acyl-enzyme intermediate. It is also possible that these side-chain modifications might affect the protonation of the tertiary amine, reducing the enzyme-inhibitor binding.

To study the structural requirements of the dibenzylamino moiety of the inhibitors, compounds 64–80 were synthesized and tested. First we found that compound 80 where the two phenyl rings are bound together, forming a comparatively flat tricyclic structure, had only 20% of the activity of the parent compound 54, showing that higher activity is obtained when the benzyl groups are free.

In compounds 64–79 we have introduced a variety of substituents in one or both of the aromatic rings. Generally speaking, the results of these experiments show that the introduction of a hydrophilic substituent with hydrogen bonding properties decreases the activity compared to that of 54. Lipophilic substituents, on the other hand, increase the activity of the inhibitor. Some of the compounds with lipophilic substituents, notably 74–79, are among the most active fibrinoligase inhibitors described.

Experimental Section

General Comments. Melting points were determined using calibrated Anschütz thermometers in an electrically heated metal block. IR spectra were run on a Perkin-Elmer 237 spectrophotometer and nmr spectra were recorded with a Varian A-60 instrument using CDCl₃ solutions and with TMS as internal standard. New compounds, which were analyzed for C, H, and N, gave values within $\pm 0.4\%$ of the theoretical ones.

4-Chlorodibenzylamine¹¹ (1), 4-dimethylaminodibenzylamine¹¹ (2), 4-nitrodibenzylamine¹¹ (3), 4-methoxydibenzylamine¹¹ (4), 4-methyldibenzylamine¹¹ (5), 3,4-dichlorodibenzylamine¹² (6), 4,4'-dichlorodibenzylamine¹³ (7), and benzylnaphthylmethylamine¹⁴ (8) were prepared as described in the literature.

4-Acetamidodibenzylamine (9). A mixture of 4-acetamidobenzaldehyde (3.7 g, 0.02 mol) and benzylamine (2.2 g, 0.02 mol) was refluxed for 2 hr in 250 ml of ethanol. The solution was cooled and hydrogenated over platinum. The solvent was evaporated and the product (4 g, 80%) crystallized upon standing; mp 74–75° (from ethanol-ether). *Anal.* (C₁₆H₁₈N₂O) C, H, N.

4-Cyanodibenzylamine (10). A mixture of 4-cyanobenzylbromide (20 g, 0.1 mol) and benzylamine (21.8 g, 0.2 mol) in 300 ml of absolute ethanol was refluxed for 1 hr. Water (400 ml) was then added and the product was extracted with chloroform (3 $\times$ 300 ml). The extract was dried (Na₂SO₄), the solvent evaporated, and the residual yellow oil distilled. This yielded 14.5 g (65%) of the title compound; bp 154° (0.01 mm); the hydrochloride had mp 253–254° (from ethanol). *Anal.* (C₁₅H₁₄N₂ · HCl) C, H, N.

4-Fluorodibenzylamine (11) was prepared as described for 10 in 53% yield; bp 105–108° (0.1 mm); mp (hydrochloride) 270–271° (from ethanol). *Anal.* (C₁₄H₁₄FN · HCl) C, H, N.

4-Bromodibenzylamine (12) was prepared as described for 10 in 55% yield; bp 147–150° (0.1 mm); mp (hydrochloride) 260–261° (from ethanol). *Anal.* (C₁₄H₁₄BrN · HCl) C, H, N.

6,7-Dihydro-5H-dibenz[c,e]azepine (13). This compound was obtained by LiAlH₄ reduction of diphenimide.¹⁵ It had the same physical constants as previously reported for the compound,¹⁶ obtained via a different route.

Synthesis of the Inhibitors. Method A. This procedure is illustrated by the synthesis of 5-(4-methyldibenzylamino)pentylamine (73).

5-(4-Methyldibenzylamino)valeronitrile (26). 4-Methyldibenzylamine (5 21.1 g, 0.1 mol) and 5-chlorovaleronitrile (5.8 g, 0.05 mol) were dissolved in dry xylene (300 ml), 0.1 g of NaI was added, and the mixture was stirred and refluxed overnight. The amine hydrochloride formed was filtered off, and the filtrate was washed with saturated NaHCO₃ solution and extracted with 2 N HCl (500 ml). The acidic aqueous solution was made alkaline and the base extracted with ether (3 $\times$ 100 ml). The extract was dried (Na₂SO₄) and the solvent evaporated *in vacuo*. The residual oil was purified by column chromatography on silica gel eluted with ether-petroleum ether mixtures of increasing polarity. The title compound was obtained as an oil (6.6 g, 45%) and precipitated as oxalate; mp 126–128° (from 2-propanol). *Anal.* (C₂₀H₂₄N₂ · C₂H₂O₄) C, H, N.

5-(4-Methyldibenzylamino)pentylamine (73). To a suspension of LiAlH₄ (1.2 g, 32 mmol) in anhydrous ether (100 ml) was slowly added a solution of 26 (in the basic form) (6 g, 0.02 mmol) in anhydrous ether, whereupon the mixture was stirred at room temperature for 3 hr. After addition of water the amine was extracted with 2 N HCl (200 ml), the aqueous phase realkalized, and the amine extracted into ether. The extract was dried (Na₂SO₄) and the solvent evaporated yielding 4.7 g (77%) of a yellow oil which was converted to its hydrochloride; mp 175–177° (from 2-propanol-ether). *Anal.* (C₂₀H₂₈H₂ · 2HCl) C, H, N.

Method B is illustrated by the synthesis of 5-(4-bromodibenzylamino)pentylamine (76).

1-(4-Bromodibenzylamino)-5-phthalimidopentane (35). A solution of 4-bromodibenzylamine (12, 7.8 g, 28.2 mmol), *N*-(5-bromopentyl)phthalimide (8.3 g, 28.2 mmol), and triethylamine (2.85 g, 28.2 mmol) in dry DMF (200 ml) was heated to 150° for 3 hr. The solution was cooled and diluted with 500 ml of water and the product extracted with chloroform (3 $\times$ 100 ml). The extract was dried (Na₂SO₄) and the solvent evaporated *in vacuo*. The brown oil was purified by column chromatography on silica gel eluted with ether-petroleum ether mixtures of increasing polarity. This yielded the title compound as an oil which solidified upon standing; yield 7.5 g (55%); mp 54–55° (from petroleum ether). *Anal.* (C₂₇H₂₇BrN₂O₂) C, H, N.

5-(4-Bromodibenzylamino)pentylamine (76). Compound 35

(6.3 g, 13 mmol) and hydrazine hydrate (0.64 g, 13 mmol) in absolute ethanol (100 ml) were refluxed for 3 hr. The reaction mixture was then cooled to 5°, 10 ml of concentrated HCl was added, and the suspension was stirred at 5° for 1 hr. The precipitate was filtered and washed with 100 ml of absolute ethanol. The filtrate and washings were pooled and concentrated *in vacuo* and alkalized and the product was extracted with ether. The ether extract was dried (Na_2SO_4) and evaporated yielding 4 g (87%) of an oil which was converted to its hydrochloride: mp 184–186° (from ether–ethanol). *Anal.* ($\text{C}_{19}\text{H}_{25}\text{BrN}_2 \cdot 2\text{HCl}$) C, H, N.

Method C is illustrated by the synthesis of 44.

5-Benzylaminopentylamine (44). A mixture of benzyl bromide (3.4 g, 0.02 mol) and cadaverine (6.1 g, 0.06 mol) in chloroform (100 ml) was stirred at room temperature overnight. Aqueous NaOH (50 ml, 1 N) was added, the mixture was stirred for 5 min, and the layers were separated. The chloroform layer was dried (Na_2SO_4) and evaporated and the residue distilled. This yielded 2.5 g (66%) of the title compound, bp 110–112° (0.4 mm). *Anal.* ($\text{C}_{12}\text{H}_{20}\text{N}_2$) C, H, N.

Method D is illustrated by the synthesis of 49.

5-(1-Phenylethylamino)pentylamine (49). A mixture of acetophenone (2.4 g, 0.02 mol) and cadaverine (3 g, 0.03 mol) in ethanol (50 ml) was heated to reflux for 1 hr. Platinum oxide was then added and the cool solution was hydrogenated at atmospheric pressure and room temperature until the theoretical amount of hydrogen had been consumed. After filtration, the solvent was removed *in vacuo* and the residue distilled: bp 110–112° (0.7 mm); yield 1.3 g (33%). *Anal.* ($\text{C}_{13}\text{H}_{22}\text{N}_2$) C, H, N.

N-But-3-ynylphthalimide (38). Phthalic acid anhydride (11.8 g, 0.08 mol) was suspended in glacial acetic acid (80 ml) whereupon 4-amino-1-butyne¹⁷ (5.5 g, 0.08 mol) was added dropwise. The mixture was stirred and refluxed until a clear solution was obtained and then for another 45 min. Upon cooling, the product separated as white needles: mp 134–135° (from glacial acetic acid); yield 14 g (88%); ir (KBr) 3250 ($\equiv\text{CH}$), 1760 and 1710 cm^{-1} (imide); nmr δ 1.9 (t, 1 H, $J = 2.6$ Hz, $\equiv\text{CH}$), 2.55 (triplet of doublets, 2 H, $J = 7.0$ and 2.6 Hz, respectively, $\text{NCH}_2\text{CH}_2\text{C}\equiv\text{CH}$), 4.75 (t, 2 H, $J = 7.0$ Hz, $\text{NCH}_2\text{CH}_2\text{C}\equiv\text{CH}$), and 7.6 ppm (m, 4 H, ArH). *Anal.* ($\text{C}_{12}\text{H}_9\text{NO}_2$) C, H, N.

1-Dibenzylamino-5-phthalimido-2-pentyne (39). Compound 38 (10 g, 0.05 mol), paraformaldehyde (2.0 g, 0.067 mol), and cuprous chloride (0.1 g, catalyst) were suspended in a mixture of glacial acetic acid (7 ml) and dry peroxide-free dioxane (20 ml). The suspension was stirred for 15 min whereupon dibenzylamine (11.0 g, 0.06 mol) was added dropwise. The mixture was then heated at 80° for 45 min. To the cool reaction mixture was then added water (100 ml) and sulfuric acid (1 M, 30 ml). The acidic solution was washed with two 100-ml portions of ether and the washings were discarded. The aqueous phase was then alkalized using K_2CO_3 and 25% ammonia (10 ml) was added to dissolve precipitated copper salts. The alkaline aqueous mixture was then extracted with chloroform (3 $\times$ 100 ml) and the extracts were dried (Na_2SO_4). After evaporation of the solvent, 10.2 g (50%) of the title compound was obtained: mp 85–86° (from 70% ethanol); ir (KBr) 1770 and 1720 cm^{-1} (imide); nmr δ 2.63 (triplet of triplets, 2 H, $J = 7.0$ and 2.0 Hz, respectively, $\text{NCH}_2\text{CH}_2\text{C}\equiv\text{CCH}_2\text{N}$), 3.10 (t, 2 H, $J = 2.0$ Hz, $\text{NCH}_2\text{CH}_2\text{C}\equiv\text{CCH}_2\text{N}$), 3.45 (s, 4 H, ArCH_2), 3.82 (t, 2 H, $J = 7.0$ Hz, NCH_2CH_2), 7.10 (s, 10 H, ArH in benzylic groups), and 7.59 ppm (m, 4 H, ArH in phthalimido group). *Anal.* ($\text{C}_{27}\text{H}_{24}\text{N}_2\text{O}_2$) C, H, N.

5-Phthalimidopentanoic Acid. Phthalic acid anhydride (14.8 g, 0.1 mol) and 5-aminopentanoic acid (11.7 g, 0.1 mol) were refluxed in 100 ml of glacial acetic acid for 1 hr. The solvent was evaporated leaving the crude product as a dark mass, which was macerated with 600 ml of ether. A small amount of tarry material remained undissolved and was discarded. The ether was evaporated and the product recrystallized three times from ethyl acetate: yield 12.0 g (48%); mp 118.5–119.5°. *Anal.* ($\text{C}_{13}\text{H}_{13}\text{NO}_4$) C, H, N.

N-(4-Dibenzylcarbamoylbutyl)phthalimide (41). 5-Phthalimidopentanoic acid (6.0 g, 0.024 mol) was suspended in dry benzene. Triethylamine (2.4 g, 0.024 mol) was added. The acid dissolved rapidly. To the clear solution was added ethyl chloroformate (2.6 g, 0.024 mol). Almost immediately a finely crystalline precipitate started to form. The mixture was stirred for 1 hr. Dibenzylamine (4.7 g, 0.024 mol) was added and the mixture was stirred for another hour. The precipitate was filtered off and the solvent was evaporated leaving 9 g of crude product as a yellowish, soft solid. Recrystallization from 70% EtOH yielded 7.0 g (70%) of white crystals, mp 85–86°. *Anal.* ($\text{C}_{27}\text{H}_{26}\text{N}_2\text{O}_3$) C, H, N.

2-Amino-2'-dibenzylaminodiethyl Sulfide (57). Sodium (2.6 g, 0.113 g-atom) was dissolved in absolute ethanol (150 ml) and 2-mercaptoethylamine (4.3 g, 0.056 mol) was slowly added under nitrogen atmosphere. 2-Chloroethyldibenzylamine hydrochloride (15 g, 0.051 mol) was then added and the mixture was refluxed under nitrogen for 4 hr. After filtration, the solvent was removed *in vacuo*, and the residual oil was dissolved in ether, washed twice with saturated NaHCO_3 solution (200 ml) and with water (200 ml), and dried (Na_2SO_4). The product was precipitated as its fumarate: mp 140–142° (from ethanol–ether); yield 18.3 g (85%); nmr δ 1.20 (s, 2 H, $-\text{NH}_2$), 2.10–2.80 (m, 8 H, $-\text{CH}_2-$), 3.49 (s, 4 H, ArCH_2-), and 7.18 ppm (m, 10 H, ArH). *Anal.* ($\text{C}_{18}\text{H}_{24}\text{N}_2\text{S} \cdot 0.5\text{C}_4\text{H}_4\text{O}_4$) C, H, N.

N,N-Dibenzyl-5-amino-3-thiapentanoic Acid Amide (59). Sodium (0.7 g, 0.03 g-atom) was dissolved in absolute ethanol (80 ml) and 2-mercaptoethylamine (2.5 g, 0.03 mol) was added under nitrogen atmosphere, followed by N,N-dibenzylchloroacetamide¹⁸ (8.2 g, 0.03 mol). A precipitate was formed and the mixture was stirred and refluxed under nitrogen for 2.5 hr. After filtration, the solution was neutralized using acetic acid and the solvent was evaporated *in vacuo*. The residue was dissolved in chloroform (100 ml), washed twice with saturated NaHCO_3 solution (100 ml) and once with water (100 ml), and dried (Na_2SO_4). The product was then precipitated as the fumarate: mp 139–140° (from ethanol–ether); yield 11.6 g (90%); ir (film) of the base, 3480 and 3410 (NH_2), 1630 cm^{-1} (CO); nmr δ 1.40 (s, 2 H, $-\text{NH}_2$), 2.6–3.0 (m, 4 H, $-\text{SCH}_2\text{CH}_2\text{N}$), 4.45 (s, 4 H, ArCH_2-), and 7.10 ppm (m, 10 H, ArH). *Anal.* ($\text{C}_{18}\text{H}_{22}\text{N}_2\text{OS} \cdot \text{C}_4\text{H}_4\text{O}_4$) C, H, N.

4-(Dibenzylcarbamoyl)butylamine (60). N-(4-Dibenzylcarbamoylbutyl)phthalimide (5.0 g, 0.012 mol) was treated with hydrazine hydrate in refluxing ethanol according to the method described by Ing and Manske.¹⁹ The crude product was placed on an alumina column and eluted with ether. The purified product was isolated as a viscous, colorless oil, which gave only one spot, R_f 0.7, on tlc (silica gel, eluent 75% Et_2O + 20% MeOH + 5% NH_4OH): mass spectrum parent ion m/e 296 (5%); mol wt ($\text{C}_{19}\text{H}_{24}\text{N}_2\text{O}$) 296.42.

5-(4-Carboxydibenzylamino)pentylamine (67). Compound 66 (1 g, 3.2 mmol) was dissolved in 50 ml of 80% ethanol containing 10% KOH. The solution was refluxed overnight, cooled, and neutralized with 2 M HCl. Inorganic salts were precipitated by addition of ether (200 ml) and removed by filtration. The filtrate was evaporated *in vacuo* and the remaining water was removed by azeotropic distillation with benzene. This yielded 0.4 g (37%) of 67 as white crystals, mp 202–205° (from ethanol–water). *Anal.* ($\text{C}_{20}\text{H}_{26}\text{N}_2\text{O}_2$) C, H, N.

5-(4-Ethylaminodibenzylamino)pentylamine (69). Compound 68 (1 g, 2.9 mmol) was stirred with LiAlH_4 (0.3 g, 8 mmol) in dry THF at 70° for 24 hr. After addition of water, the amine was extracted with 50 ml of 2 M HCl, the aqueous layer alkalized, and the product extracted with ether (2 $\times$ 50 ml). The extract was dried (Na_2SO_4), the solvent evaporated, and the title compound was obtained as an oil (0.5 g, 51%). The compound could not be obtained in crystalline form as the base or as salts. A small amount was purified by preparative tlc for elementary analysis: ir (film) 3450–3200 (NH_2), 1600 and 1500 cm^{-1} (phenyl). *Anal.* ($\text{C}_{21}\text{H}_{31}\text{N}_3$) C, H, N.

5-(4-Aminomethyldibenzylamino)pentylamine (72). This compound was prepared from 66 by reduction with LiAlH_4 as described for 69: yield 92%; mp 149–150° (fumarate, from ethanol). *Anal.* ($\text{C}_{20}\text{H}_{26}\text{N}_3 \cdot \text{C}_4\text{H}_4\text{O}_4$) C, H, N.

N,N-Dibenzyl-N',N'-dimethyl-1,5-diaminopentane (81). N,N-Dibenzyl-1,5-diaminopentane (28.2 g, 0.1 mol) and 98% formic acid (42.0 g, 0.9 mol) were mixed and cooled in an ice bath. Ice-cold 37% formaldehyde solution (18.0 g, 0.22 mol) was added and the mixture was heated on a steam bath. A vivid evolution of carbon dioxide indicated that the reaction had started. The heating was continued for 45 min whereupon 100 ml of ice-water was added to the reaction mixture, which was then made alkaline by the addition of solid K_2CO_3 . The product separated as an oil and was taken up in 200 ml of ether. The ether extract was dried (K_2CO_3) and the solvent was evaporated leaving the crude product as a faintly yellow oil. Distillation yielded 27.3 g (88%) of pure product, bp 163–165° (0.05 mm). The ir spectrum of the base showed no NH band. The dihydrochloride had mp 214–216°. *Anal.* ($\text{C}_{21}\text{H}_{30}\text{N}_2 \cdot 2\text{HCl}$) C, N, H.

5-(Dibenzylmethyllumonio)pentylamine Bromide (82). Compound 54 (0.9 g) was dissolved in dry acetone (50 ml) and the solution was left at room temperature overnight. Methyl bromide (1 g) was then added and the solution left at room temperature for

another 3 days. An oil separated, which could not be crystallized: ir (film) a band at 1690 cm^{-1} indicated the presence of a $\text{-N=C(CH}_3)_2\text{-}$ structure. The oil was dissolved in water (25 ml) and the solution left at room temperature overnight. Evaporation yielded 0.5 g of an oil which could not be crystallized: ir (film) $3200\text{--}3300\text{ cm}^{-1}$ (NH_2), no band at 1690 cm^{-1} . A small amount was purified for analysis by preparative tlc developed in ethanol. *Anal.* ($\text{C}_{20}\text{H}_{29}\text{BrN}_2$) C, H, N.

Acknowledgment. We wish to thank Professor Laszlo Lorand for valuable discussion. This work has been supported by the Swedish Medical Research Council Project No. 03X-3770.

References

- (1) J. L. G. Nilsson, P. Stenberg, Ch. Ljunggren, K.-J. Hoffmann, R. Lundén, O. Eriksson, and L. Lorand, *Ann. N. Y. Acad. Sci.*, **202**, 286 (1972).
- (2) L. Lorand and J. L. G. Nilsson, "Drug Design," Vol. 3, Academic Press, New York, N.Y., 1972, p 415.
- (3) L. Lorand, *Thromb. Diath. Haemorrh., Suppl.*, **39**, 75 (1970).
- (4) L. Lorand, C. H. Chou, and I. Simpson, *Proc. Nat. Acad. Sci. U. S.*, **69**, 2645 (1972).
- (5) Ch. Ljunggren, K.-J. Hoffmann, P. Stenberg, and J. L. G. Nilsson, *J. Med. Chem.*, **16**, 1186 (1973).
- (6) P. Stenberg, Ch. Ljunggren, J. L. G. Nilsson, R. Lundén, and O. Eriksson, *J. Med. Chem.*, **15**, 674 (1972).
- (7) J. L. G. Nilsson, P. Stenberg, O. Eriksson, and R. Lundén, *Acta Pharm. Suecica*, **7**, 441 (1970).
- (8) J. H. Brewster and E. L. Eliel, *Org. React.*, **7**, 99 (1953).
- (9) J. L. G. Nilsson, P. Stenberg, Ch. Ljunggren, O. Eriksson, and R. Lundén, *Acta Pharm. Suecica*, **8**, 497 (1971).
- (10) Ch. Ljunggren, K.-J. Hoffman, P. Stenberg, U. Svensson, J. L. G. Nilsson, A. Hartkoorn, and R. Lundén, *J. Med. Chem.*, **17**, 649 (1974).
- (11) R. L. Hinmann and K. L. Hamm, *J. Org. Chem.*, **23**, 529 (1958).
- (12) W. J. Houlihan, German Patent 1231690; *Chem. Abstr.*, **66**, 55248e (1967).
- (13) C. Winans, *J. Amer. Chem. Soc.*, **61**, 3564 (1939).
- (14) H. Dahn, U. Solms, and P. Zoller, *Helv. Chim. Acta*, **35**, 2117 (1952).
- (15) H. W. Underwood, Jr., and L. A. Clough, *J. Amer. Chem. Soc.*, **51**, 583 (1929).
- (16) J. O. Hawthorne, E. L. Mihelic, M. S. Morgan, and M. H. Witt, *J. Org. Chem.*, **28**, 2831 (1963).
- (17) J. L. Dumont, J. Tohier, and P. Cadiot, *C. R. Acad. Sci.*, **256**, 3146 (1963).
- (18) A. J. Speziale and P. C. Hamm, *J. Amer. Chem. Soc.*, **78**, 2556 (1956).
- (19) H. R. Ing and R. F. Manske, *J. Chem. Soc.*, 2348 (1926).

[1- β -Mercapto- β,β -pentamethylenepropionic acid]oxytocin, a Potent Inhibitor of Oxytocin[†]

John J. Nestor, Jr., Martha F. Ferger, and Vincent du Vigneaud*

Department of Chemistry, Cornell University, Ithaca, New York 14850. Received October 15, 1974

[1- β -Mercapto- β,β -pentamethylenepropionic acid]oxytocin was prepared from β -Mpa(β -(CH_2)₅)(Bzl)-Tyr(Bzl)-Ile-Gln-Asn-Cys(Bzl)-Pro-Leu-Gly-NH₂ by removal of the Bzl-protecting groups with Na-NH₃ followed by cyclization of the resulting disulfhydryl compound with K₃Fe(CN)₆. The analog was purified by desalting on Sephadex G-15 in 50% HOAc and gel filtration on Sephadex G-25 and LH-20. The protected intermediate above was synthesized from Z-Cys(Bzl)-Pro-Leu-Gly-NH₂ by the stepwise *p*-nitrophenyl ester method using N^α-Boc protection at the penta-, hexa-, and octapeptide stages. The analog was found to be a potent inhibitor of the oxytocic and avian vasodepressor effects of oxytocin (pA_2 values of 7.43 and 8.30, respectively) but was only a weak inhibitor of the rat pressor effect of 8-lysine-vasopressin. The rat antipressor potency of [1-deaminopenicillamine]oxytocin was also determined in this study: $\text{pA}_2 = 6.27$. Of the alkyl-substituted 1-position analogs of oxytocin studied so far, [1- β -mercapto- β,β -pentamethylenepropionic acid]oxytocin is the most potent antioxytocic agent.

When [1-L-penicillamine]oxytocin, [1-D-penicillamine]oxytocin, and [1-deaminopenicillamine]oxytocin ([1- β -mercapto- β,β -dimethylpropionic acid]oxytocin) were found to be potent inhibitors² of the oxytocic and avian vasodepressor (AVD) activities of oxytocin (Figure 1), studies were undertaken of related modifications³ in the 1 position of the highly potent deaminooxytocin.⁴ [1- β -Mercapto- β,β -diethylpropionic acid]oxytocin³ ([1- β -Mpa(β -Et₂)]oxytocin) was found to be the most potent inhibitor of the compounds that had been prepared, having pA_2 values of 7.24 in the antioxytocic and 8.11 in the anti-AVD assays in comparison with corresponding pA_2 values for [1-deaminopenicillamine]oxytocin of 6.94 and 7.88, respectively.

As a further variation, the incorporation of a β -mercapto- β,β -pentamethylenepropionic acid residue [β -Mpa(β -(CH_2)₅)] into the 1 position of oxytocin is reported here. This residue is structurally similar to β -Mpa(β -Et₂) and

can be looked upon as having the β -geminal ethyl groups connected by a methylene bridge. The possible conformations of the β -alkyl substituents are thus limited and the overall lipophilicity of the residue is increased.

[1- β -Mpa(β -(CH_2)₅)]oxytocin was prepared from the protected polypeptide β -Mpa(β -(CH_2)₅)(Bzl)-Tyr(Bzl)-Ile-Gln-Asn-Cys(Bzl)-Pro-Leu-Gly-NH₂ by removal of the benzyl-protecting groups with Na-NH₃⁵ followed by cyclization with potassium ferricyanide⁶ solution. The hormone analog was purified by desalting in 50% HOAc on Sephadex G-15 by the method of Manning⁷ followed by gel filtration in 0.2 N HOAc on Sephadex G-15 and in DMF on Sephadex LH-20.

N^α-tert-Butyloxycarbonyl (Boc) protection was used on the penta-, hexa-, and octapeptide intermediates. Boc removal was accomplished by treatment with trifluoroacetic acid (TFA) at room temperature. In the case of the Boc-Tyr(Bzl)-terminal octapeptide it is important that the deprotection period be no longer than 15 min. Longer periods lead to removal of the benzyl ether protection. *p*-Nitrophenyl esters⁸ were used for coupling reactions throughout. β -(S-Benzylmercapto)- β,β -pentamethylenepropionic acid was prepared from ethyl cyclohexylideneacetate⁹ by means

[†] This work was supported in part by Grant HL-11680 from the National Heart and Lung Institute, U. S. Public Health Service. All optically active amino acids are of the L configuration. The symbols for the amino acids follow the recommendations of the IUPAC-IUB Commission on Biochemical Nomenclature.¹ Where analyses are indicated only by symbols for the elements, analytical results obtained for the elements were within $\pm 0.4\%$ of the theoretical values.