

## Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/uopp20>

### A FACILE METHOD WITH IMPROVED YIELDS IN THE SYNTHESIS OF 6-ARYLPYRIDO[2',3':4,5]PYRIMIDO[1,6-a]BENZIMIDAZOLES

G. Suma<sup>a</sup>, R. H. Bahekar<sup>a</sup> & A. R. R. Rao<sup>a</sup>

<sup>a</sup> Medicinal Chemistry Division, University College of Pharmaceutical Sciences, Kakatiya University, Warangal, AP, 506 009, INDIA

Published online: 18 Feb 2009.

To cite this article: G. Suma, R. H. Bahekar & A. R. R. Rao (2000) A FACILE METHOD WITH IMPROVED YIELDS IN THE SYNTHESIS OF 6-ARYLPYRIDO[2',3':4,5]PYRIMIDO[1,6-a]BENZIMIDAZOLES, Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 32:1, 99-101, DOI: [10.1080/00304940009356756](https://doi.org/10.1080/00304940009356756)

To link to this article: <http://dx.doi.org/10.1080/00304940009356756>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms &



8. Y. Ikeda, Y. Ueki, T. Nishihara and Y. Kamikawa, US 5607948; *Chem. Abstr.*, **122**, 266008 (1997).
9. Vendor: Bachem Bioscience, King of Prussia, PA, USA.
10. S. Nahm and S. M. Weinreb, *Tetrahedron Lett.*, **22**, 3815 (1981).
11. M. Bodanszky and A. Bodanszky, "*The Practice of Peptide Synthesis*", p. 109, Springer-Verlag, New York, NY, **1984**.
12. Vendor: Aldrich Chemical, Milwaukee, WI, USA.

\*\*\*\*\*


### A FACILE METHOD WITH IMPROVED YIELDS IN THE SYNTHESIS OF 6-ARYLPYRIDO[2',3':4,5]PYRIMIDO[1,6-a]BENZIMIDAZOLES<sup>†</sup>

Submitted by  
(03/02/99)

G. Suma<sup>†</sup>, R. H. Bahekar and A. R. R. Rao\*

*Medicinal Chemistry Division  
University College of Pharmaceutical Sciences  
Kakatiya University, Warangal (A. P.) 506 009, INDIA*

Methylxanthines are a major class of bronchodilators employed in the treatment of asthma despite a narrow therapeutic index. New heterocyclic compounds designed on the basis of the xanthine skeleton are being investigated as possible bronchodilators with a wider margin of safety.<sup>1</sup> In continuation of our work on the synthesis and bronchodilator activity of some new fused quinazolines,<sup>2</sup> we carried out the preparation of some fused pyridopyrimidines as isosteres of the quinazoline moiety. A literature survey revealed that the title compounds (**4**) have been synthesized with overall yields of about 34%.<sup>3,4</sup> We now report the preparation of the above compounds by a route analogous to the synthesis of benzimidazoquinazolines,<sup>2</sup> involving mild reaction conditions and easy work-up procedures with improved overall yields (46-70%).



## EXPERIMENTAL SECTION

Melting points were determined in open capillaries on a Thermonik Precision Melting Point cum Boiling Point Apparatus-Model C-PMB-2 and are uncorrected. Column chromatography and tlc were performed on silica gel (230-400 mesh) and on precoated plates (E. Merck Kieselgel 60 F<sub>254</sub>) respectively. IR spectra were recorded using KBr pellets on Perkin-Elmer Infrared 337 spectrophotometer and <sup>1</sup>H NMR spectra were recorded on a Varian EM 390 90 MHz spectrometer using TMS as internal standard and the chemical shift values expressed in  $\delta$  ppm. Mass spectra (EIMS) were recorded on Jeol D-300 spectrometer at 70 eV. Elemental analyses was carried out using Heraeus Carlo Erba 1108 CHN analyser. The starting material, 2-aminonicotinic acid was procured from Janssen Chimica, Belgium.

**2-Arylpyrido[2,3-b]1,3-oxazin-8-one (2).**- The respective arylcarboxylic acid chloride (10 mmol) was added dropwise to a solution of 2-aminonicotinic acid (1.38 g, 10 mmol) in dry pyridine (20 mL) with continuous stirring for 4 hrs under ice-cold conditions. The reaction mixture was poured onto ice-cold water (150 mL) and the crystalline product was collected and recrystallized from methanol.

**6-Aryl-(2-aminophenyl)pyrido[2',3'-b]-7H-pyrimidin-8-ones (3).**- Compound 2 (2.0 mmol) was reacted with *o*-phenylenediamine (0.324 g, 3.0 mmol) in dry pyridine by refluxing for 15 hrs. Excess solvent was removed under reduced pressure and the residue obtained was dissolved in crushed ice. The pH was adjusted to 7 by neutralization with glacial acetic acid and the solid product obtained was collected and recrystallized from methanol, except **3b** which was carried out in DMSO.

**6-Arylpyrido[2',3':4,5]pyrimido[1,6-a]benzimidazole (4).**- Compound 3 (1.0 mmol) was heated above its melting point under anhydrous conditions for 15 min. The solid residue obtained on cooling to room temperature was triturated with petroleum ether and recrystallized from methanol.

TABLE 1. Yields, mps. and Elemental Analyses of Compounds 2-4

Cmpd	Yield (%)	mp. <sup>a</sup> (°C)	Elemental Analyses (Found)		
			C	H	N
<b>2a</b>	70	141	69.64 (69.51)	3.57 (3.62)	12.50 (12.61)
<b>2b</b>	90	231	57.99 (57.72)	2.60 (2.51)	15.61 (15.74)
<b>2c</b>	86	201	60.35 (60.51)	2.71 (2.62)	10.83 (10.65)
<b>2d</b>	84	187	70.59 (70.68)	4.20 (4.35)	11.76 (11.82)
<b>3a</b>	74	203	72.61 (72.46)	4.46 (4.52)	17.83 (17.99)
<b>3b</b>	85	283 <sup>b</sup>	63.51 (63.26)	3.62 (3.48)	19.50 (19.41)
<b>3c</b>	78	241	65.42 (65.26)	3.73 (3.89)	16.06 (16.21)
<b>3d</b>	73	234	73.17 (73.29)	4.88 (4.95)	17.07 (17.16)
<b>4a</b>	89	241 <sup>c</sup>	77.03 (77.26)	4.05 (4.19)	18.92 (18.78)
<b>4b</b>	92	>300 <sup>c</sup>	66.86 (66.61)	3.22 (3.39)	20.53 (20.65)
<b>4c</b>	88	242 <sup>c</sup>	68.99 (68.82)	3.33 (3.11)	16.94 (16.79)
<b>4d</b>	84	261 <sup>c</sup>	77.42 (77.39)	4.52 (4.61)	18.06 (18.25)

a) Recrystallized from methanol except where noted. b) Recrystallized from DMSO. c) Lit.<sup>3</sup> mp. 4a-259°, 4b-284°, 4c-278°, 4d-236°.

**TABLE 2.** Mass Spectral Data of Compounds **4<sup>a</sup>**

Cmpd	Mass Spectrum [m/e (ion %)]
<b>4a</b>	297 ( $M^+ + 1$ , 20.9), 296 ( $M^+$ , 100), 295 ( $M^+ - H$ , 78.1), 268 (295-HCN, 2.3), 193 ( $M^+ - C_6H_5 - CN$ , 13.7), 166 (193-HCN, 9.8), 103 ( $PhCN^+$ , 14), 77 (Ph, 12).
<b>4b</b>	342 ( $M^+ + 1$ , 18.5), 341 ( $M^+$ , 95.4), 340 ( $M^+ - H$ , 14.1), 311 ( $M^+ - NO$ , 30.3), 310 (340-NO, 34.9), 296 (342- $NO_2$ , 27.5), 295 ( $M^+ - NO_2$ , 39.6), 294 ( $M^+ - NO_2$ , 52.1), 283 (311-CO, 6.1), 282 (310-CO, 8.4), 194 (342- $NO_2 - C_6H_4 - CN$ , 10), 193 ( $M^+ - NO_2 - C_6H_4 - CN$ , 23.6), 166 (193-HCN, 13), 148 ( $NO_2 - C_6H_4 - CN$ , 6.2), 104 (Ph-CN, 18), 77 (Ph, 19).
<b>4c</b>	333 ( $M^+ + 3$ , isotopic, 7), 332 ( $M^+ + 2$ , isotopic, 32.8), 331 ( $M^+ + 1$ , 42.6), 330 ( $M^+$ , 100), 329 ( $M^+ - H$ , 73.6), 193 ( $M^+ - Cl - C_6H_5 - CN$ , 66), 166 (193-HCN, 19), 104 (Ph-CN, 18), 77 (Ph, 15).
<b>4d</b>	310 ( $M^+$ , 2.1), 209 ( $M^+ - 101$ , 28), 208 ( $M^+ - 102$ , 100), 193 ( $M^+ - H_3C - C_6H_4CN$ , 3.7), 166 (193-HCN, 0.8), 117 ( $H_3C - C_6H_4 - CN$ , 2.7), 104 ( $PhCN^+$ , 10.7), 91 ( $C_7H_7$ , 11), 77 (Ph, 4.3).

a) Spectral Data for **4b**: IR (KBr): 3054 (Ar-H), 1625, 1594 (C=N)  $cm^{-1}$ ;  $^1H$  NMR ( $CDCl_3$ ):  $\delta$  7.5-8.0 (m, 8H, Aromatic), 6.9 (d, 1H,  $C_2-H$ ), 9.0 (s, 2H,  $C_1$  and  $C_3-H$ ).

**Acknowledgement.**- The authors thank All India Council for Technical Education (AICTE), New Delhi for financial support, Regional Sophisticated Instrumentation Centre of Central Drug Research Institute, Lucknow for spectral and analytical data and Dr. K. Mogilaiah for useful suggestions. One of them (GS) is thankful to University Grants Commission, New Delhi for a fellowship (JRF).

## REFERENCES

- † G. Suma, Part of the Master of Pharmacy Dissertation, Kakatiya University, India (1998).
1. F. Suzuki, T. Kuroda, Y. Nakasato, H. Manabe, K. Ohmori, S. Kitamura, S. Itchikawa and T. Ohno, *J. Med. Chem.*, **35**, 4045 (1992).
2. A. Raghu Ram Rao and Rajesh H. Bahekar, *Indian J. Chem. Sect. B*, **38B**, 434 (1999).
3. K. Vijayender Reddy, K. Mogilaiah and B. Sreenivasulu, *ibid.*, **23B**, 1106 (1984).
4. K. Vijayender Reddy, K. Mogilaiah and B. Sreenivasulu, *ibid.*, **23B**, 866 (1984).