


Nitrosocarbonyl Compounds as Intermediates in the Oxidative Cleavage of Hydroxamic Acids

By GORDON W. KIRBY* and JAMES G. SWEENEY

(Department of Chemistry, University of Glasgow, Glasgow G12 8QQ)

Summary Periodate oxidation of hydroxamic acids in the presence of conjugated dienes gives, in good yield, *N*-acyl-3,6-dihydro-2*H*-1,2-oxazines derived, presumably, from nitrosocarbonyl intermediates.

NITROSOCARBONYL-ALKANES and -ARENES, RCONO (1), have been proposed as transient intermediates in the oxidative cleavage of hydroxamic acids,^{1,2} RCONHOH (2), and the pyrolysis of alkyl nitrites in the presence of aldehydes.² No direct evidence for the existence of the species (1) appears to be available; characteristic decomposition products are acyl derivatives formed, apparently, from attack by nucleophiles at the carbonyl group. The powerful dienophilic character of nitrosyl cyanide³ suggested that *C*-nitrosocarbonyl compounds, though short-lived, might be trapped efficiently by conjugated dienes. We find that this is so.



Addition of benzohydroxamic acid (**2**; R=Ph) (2.3 mol. equiv.) to a stirred mixture of thebaine (**3**) (1.0 mol. equiv.) in ethyl acetate and tetraethylammonium periodate⁴ (1.45 mol. equiv.) in aqueous acetic acid-sodium acetate (pH *ca.* 6) at 0 °C gave the adduct (**4**; R = Ph), isolated as the hydriodide (97% yield). Hydrolysis with refluxing aqueous methanolic hydrogen chloride gave 14-hydroxy-aminocodeinone (**5**; R = H), obtained earlier⁵ from the reaction of thebaine hydrochloride with 1-chloronitroso-cyclohexane in aqueous ethanol. Similarly, thebaine was converted, with acetohydroxamic acid (**2**; R = Me) and periodate, into the adduct (**4**; R = Me) (96%). Acid-catalysed hydrolysis at 0 °C gave the enone (**5**; R = Ac) which, upon treatment with sodium methoxide in methanol, cyclised to the phenol (**6**; R = H). Acetylation then gave the acetate (**6**; R = Ac) obtainable⁵ similarly from (**5**; R = H) by cyclisation and acetylation. Catalytic (PtO₂) hydrogenation of the adduct (**4**; R = Me) gave the corres-

ponding 7,8-dihydro-derivative. Treatment of butadiene with benzohydroxamic acid in nitromethane, or acetohydroxamic acid in dichloromethane, and tetraethylammonium periodate yielded the known⁶ *N*-acyloxazines (**7**; R = Ph or Me).

Compelling evidence for the free existence of nitroso-carbonylmethane and nitrosocarbonylbenzene was obtained as follows. The crystalline adducts (**8**; R = Me or Ph) were prepared (*ca.* 50% yield) from 9,10-dimethylantracene in the usual way. Both were stable at ambient temperatures. However, treatment of thebaine (**3**) with each adduct (**8**) in hot benzene caused intermolecular transfer of RCONO to yield the corresponding thebaine adducts (**4**) and 9,10-dimethylantracene.

We thank the S.R.C. for financial support and Dr. D. G. Morris for helpful discussions.

(Received, 27th July 1973; Com. 1087.)

¹ B. Sklarz and A. F. Al-Sayyab, *J. Chem. Soc.*, 1964, 1318; J. E. Rowe and A. D. Ward, *Austral J. Chem.*, 1968, **21**, 2761; U. Lerch and J. G. Moffatt, *J. Org. Chem.*, 1971, **36**, 3391; and references cited therein.

² A. L. J. Beckwith and G. W. Evans, *J. Chem. Soc.*, 1962, 130.

³ P. Horsewood and G. W. Kirby, *Chem. Comm.*, 1971, 1139; R. Dickinson, G. W. Kirby, J. G. Sweeny, and J. K. Tyler, *J.C.S. Chem. Comm.*, 1973, 241.

⁴ A. K. Qureshi and B. Sklarz, *J. Chem. Soc. (C)*, 1966, 412.

⁵ P. Horsewood, Ph.D. Thesis, University of Loughborough, 1972; *cf.* K. W. Bentley, P. Horsewood, G. W. Kirby, and Serjinder Singh, *Chem. Comm.*, 1969, 1411.

⁶ O. Wichterle and J. Novak, *Coll. Czech. Chem. Comm.*, 1950, **15**, 309.