

CATALYTIC REDUCTION OF HYDRAZINE TO AMMONIA BY THE REDUCED
SPECIES OF $[\text{Mo}_2\text{Fe}_6\text{S}_8\text{L}_9]^{3-}$ AND $[\text{Fe}_4\text{S}_4\text{L}_4]^{2-}$ (L = SPh, $\text{SCH}_2\text{CH}_2\text{OH}$)

Yoshiyuki HOZUMI, Yoshinobu IMASAKA, Koji TANAKA,
and Toshio TANAKA*

*Department of Applied Chemistry, Faculty of Engineering,
Osaka University, Suita, Osaka 565*

Hydrazine can be reduced to NH_3 catalytically in the presence of $[\text{Mo}_2\text{Fe}_6\text{S}_8\text{L}_9]^{3-}$ or $[\text{Fe}_4\text{S}_4\text{L}_4]^{2-}$ (L = SPh, $\text{SCH}_2\text{CH}_2\text{OH}$) in H_2O or MeOH/THF by the controlled potential electrolysis. The maximum current efficiency in the electrochemical reduction of N_2H_4 attains 97% and the turnover number of NH_3 produced per 1 h is 23 in MeOH/THF.

The reduction of N_2 to NH_3 by nitrogenase has been suggested to proceed *via* enzyme-bound N_2H_2 and N_2H_4 as intermediates.¹⁾ In fact, the formation of N_2H_4 was confirmed in the reduction of N_2 by *Klebsiella pneumonia*,²⁾ and some dinitrogen-molybdenum and -tungsten complexes reacted with mineral acids to afford N_2H_4 as well as NH_3 .³⁾ The reduction of N_2H_4 to NH_3 also was successfully conducted by the catalyst composed of sodium molybdate, L-cystein, and NaBH_4 with the turnover number of 4.2 NH_3 mol/(Mo-cystein complex mol)h.⁴⁾ The amount of H_2 which would evolve concomitantly, however, has not been determined, though the determination of H_2 is essentially important for examining the selectivity of such a catalytic system. Recently, we have found that N_2 can be reduced to NH_3 by the electrochemically reduced species of $[\text{Mo}_2\text{Fe}_6\text{S}_8(\text{SPh})_9]^{3-}$ and $[\text{Fe}_4\text{S}_4(\text{SPh})_4]^{2-}$.⁵⁾ This result has driven us to study the reduction of N_2H_4 by the same catalysts. This letter reports the catalytic reduction of N_2H_4 to NH_3 .

The reduction of N_2H_4 was conducted on an Hg working electrode in an H_2O or MeOH/THF solution containing $[\text{R}_4\text{N}]_3[\text{Mo}_2\text{Fe}_6\text{S}_8\text{L}_9]$ (R = *n*-Bu for L = SPh,^{6a)} R = Et for L = $\text{SCH}_2\text{CH}_2\text{OH}$ ^{6b)}) or $[\text{R}_4\text{N}]_2[\text{Fe}_4\text{S}_4\text{L}_4]$ (R = *n*-Bu for L = SPh,^{7a)} R = Me for L = SCH_2CH_2 -




Fig. 1. The reduction of N_2H_4 by the reduced species of $[\text{Mo}_2\text{Fe}_6\text{S}_8(\text{SPh})_9]^{3-}$ (-1.30 V vs. SCE) suspended in H_2O at pH 12.0.

$\text{OH}^{7b)}$ under the controlled potential electrolysis conditions, as described elsewhere.⁸⁾ Hydrazine was reduced to NH_3 with concomitant evolution of H_2 arising from the reduction of hydrogen ions under the controlled potential electrolysis (-1.30 V vs. SCE) of $[\text{Mo}_2\text{Fe}_6\text{S}_8(\text{SPh})_9]^{3-9)}$ ($16 \mu\text{mol}$) suspended¹⁰⁾ in an aqueous solution (20 cm^3) of N_2H_4 (1.4 mmol) at pH 12.0 buffered with $\text{H}_3\text{PO}_4\text{-NaOH}$. The plots of the amount of NH_3 or H_2 produced divided by the amount of the catalyst vs. the reaction time were linear after the initial induction period of about 1 h, as shown in Fig. 1. The amounts of NH_3 and H_2 produced attain 26 and 19 times of that of the catalyst in 4 h, respectively, indicating that the reduction of N_2H_4 and hydrogen ions proceeds catalytically. Table 1 summarizes the results of the electrochemical reduction of N_2H_4 by four catalysts¹¹⁾ in H_2O at pH 7.0 and 12.0 or in MeOH/THF . The sum of current efficiencies calculated from Eq. 1 for the


$$\text{Current efficiency } (\eta) = \frac{\text{Amount of } \text{NH}_3 \text{ or } 2\text{H}_2 \text{ produced}}{\text{Quantity of electricity consumed}/96480 \text{ C mol}^{-1}} \times 100 \quad (1)$$

Table 1. Electrochemical Reduction of N_2H_4 (1.4 mmol) in the Presence of the Cluster Compounds (16 μ mol) for 4 h at 25°C

Cluster	Medium	Applied potential V vs. SCE	Charge C	Amount of product		η_{NH_3} %	η_{H_2} %
				NH_3	H_2		
				μ mol			
$[Mo_2Fe_6S_8(SPh)_9]^{3-}$	H_2O , pH 7.0 ^a	-1.30	175	283	724	16	80
	H_2O , pH 12.0 ^a	-1.30	110	413	311	36	56
	MeOH/THF	-1.25	145	1450	24	97	3.2
$[Fe_4S_4(SPh)_4]^{2-}$	H_2O , pH 7.0 ^a	-1.30	268	138	1370	5.0	99
	H_2O , pH 12.0 ^a	-1.30	49	97	200	19	79
	MeOH/THF	-1.25	171	790	494	44	56
$[Mo_2Fe_6S_8(SCH_2CH_2OH)_9]^{3-}$	H_2O , pH 7.0 ^b	-1.25	316	556	1370	17	83
	H_2O , pH 12.0	-1.25	34	199	62	57	35
$[Fe_4S_4(SCH_2CH_2OH)_4]^{2-}$	H_2O , pH 7.0 ^b	-1.25 ^c	194	109	936	5.4	93
	H_2O , pH 12.0	-1.25	85	110	372	12	84
None	H_2O , pH 7.0	-1.30	0	0	0	—	—

^a Suspension. ^b $HSCH_2CH_2OH$ (1.0×10^{-2} cm³) was added in order to stabilize the cluster in H_2O (20 cm³) at pH 7.0. ^c For 1 h at 25°C.

formation of NH_3 (η_{NH_3}) and H_2 (η_{H_2}) is almost 100% within the experimental errors in most cases; the evolution of N_2 resulting from the decomposition of N_2H_4 has not been observed. Therefore, the reaction involved only the reductions of N_2H_4 (Eq. 2) and hydrogen ions (Eq. 3) taking place competitively.



The catalytic activity of $[Mo_2Fe_6S_8L_9]^{3-}$ was apparently superior to $[Fe_4S_4L_4]^{2-}$ with respect to the formation of NH_3 . All the catalysts gave larger η_{NH_3} values at pH 12.0 than at pH 7.0, suggesting that smaller hydrogen ion concentrations favour the formation of NH_3 rather than H_2 . In fact, the η_{NH_3} value in an MeOH/THF

(1:1 v/v) solution of $[\text{Mo}_2\text{Fe}_6\text{S}_8(\text{SPh})_9]^{3-}$ or $[\text{Fe}_4\text{S}_4(\text{SPh})_4]^{2-}$ containing LiCl as supporting electrolyte was several times larger than that in H_2O at pH 12.0, as expected from the weak acidity of MeOH; in particular an η_{NH_3} value of 97% was obtained with $[\text{Mo}_2\text{Fe}_6\text{S}_8(\text{SPh})_9]^{3-}$ in MeOH/THF (Table 1).

References

- 1) H. Dalton and L. E. Mortenson, *Bacterial. Rev.*, **36**, 231 (1972).
- 2) R. N. F. Thormoly, A. J. Eady, and D. J. Lowe, *Nature*, **272**, 557 (1978).
- 3) J. Chatt, A. J. Pearman, and R. L. Richards, *J. Chem. Soc., Dalton Trans.*, **1977**, 1852. T. Takahashi, Y. Minobe, M. Sato, Y. Uchida, and M. Hidai, *J. Am. Chem. Soc.*, **102**, 7461 (1980).
- 4) G. N. Schrauzer, P. R. Robinson, E. L. Moorehead, and T. M. Vickrey, *J. Am. Chem. Soc.*, **98**, 2816 (1976).
- 5) K. Tanaka, Y. Hozumi, and T. Tanaka, *Chem. Lett.*, **1982**, 1203.
- 6) (a) G. Christou, C. D. Garner, and F. E. Mabbs, *J. Chem. Soc., Chem. Commun.*, **1978**, 740; (b) G. Christou, C. D. Garner, and F. E. Mabbs, *ibid.*, **1979**, 91.
- 7) (a) B. A. Averill, T. Herskovitz, R. H. Holm, and J. A. Ibers, *J. Am. Chem. Soc.*, **95**, 3523 (1973); (b) C. L. Hill, J. Renaud, R. H. Holm, and L. E. Mortenson, *ibid.*, **99**, 2549 (1977).
- 8) K. Tanaka, M. Tanaka, and T. Tanaka, *Chem. Lett.*, **1981**, 895. K. Tanaka, Y. Imasaka, M. Tanaka, M. Honjo, and T. Tanaka, *J. Am. Chem. Soc.*, **104**, 4258 (1982).
- 9) The cyclic voltammetric anode peak potentials of $[\text{Mo}_2\text{Fe}_6\text{S}_8(\text{SPh})_9]^{3-}$ in MeOH/THF were -1.08 and -1.25 V vs. SCE for the 3-/4- and 4-/5- processes, respectively.
- 10) A dimethylsulfoxide solution (0.5 cm^3) of $[\text{Mo}_2\text{Fe}_6\text{S}_8(\text{SPh})_9]^{3-}$ (16 μmol) and Triton X-100 ($1.5 \times 10^{-2} \text{ cm}^3$) was injected into water (20 cm^3) to disperse the cluster.
- 11) The anode peak potential of $[\text{Fe}_4\text{S}_4(\text{SPh})_4]^{2-}$ (2-/3-) in MeOH/THF was -1.25 V vs. SCE and those of $[\text{Mo}_2\text{Fe}_6\text{S}_8(\text{SCH}_2\text{CH}_2\text{OH})_9]^{3-}$ (3-/4- and 4-/5-) and $[\text{Fe}_4\text{S}_4(\text{SCH}_2\text{CH}_2\text{OH})_4]^{2-}$ (2-/3-) in water at pH 7.0 were -0.58, -0.76, and -0.75 V vs. SCE, respectively.

(Received March 26, 1983)