

Article

Visible Light-mediated Photochemical Reactions of 2-(2'-Alkenyloxy)cycloalk-2-enones

Raphaella Graßl, Christian Jandl, and Thorsten Bach

J. Org. Chem., **Just Accepted Manuscript** • DOI: 10.1021/acs.joc.0c01501 • Publication Date (Web): 27 Jul 2020

Downloaded from pubs.acs.org on July 31, 2020

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.

Visible Light-mediated Photochemical Reactions of 2-(2'-Alkenyloxy)cycloalk-2-enones


Raphaela Graßl, Christian Jandl, Thorsten Bach*

Department Chemie and Catalysis Research Center (CRC), Technische Universität München,
Lichtenbergstr. 4, 85747 Garching, Germany

Corresponding Author

*Email: thorsten.bach@ch.tum.de

ABSTRACT



The title compounds were prepared and their reactivity was studied upon sensitized irradiation at $\lambda = 420$ nm. Thioxanthen-9-one was employed as the sensitizer at a loading of 10 mol% in small scale reactions and of 2.5 mol% on larger scale. Cyclohex-2-enones substituted by a 2'-propenyloxy, 2'-butenyloxy, 2'-pentyloxy, or 2'-methyl-2'-propenyloxy group in 2-position gave the products of an intramolecular [2+2] photocycloaddition. The reaction proceeded with high regioselectivity (crossed product) and perfect diastereoselectivity (nine examples, 34%-99%


yield). If the olefin in the tether was trisubstituted (3'-methyl-2'-butenyloxy) no cycloaddition was observed. Rather, a cyclization with subsequent hydrogen abstraction occurred (three examples, 65-86% yield). The results are consistent with a reaction course via a triplet enone intermediate and the formation of a 1,4-diradical by an initial cyclization. The analogous cyclopent-2-enones were less prone to an intramolecular reaction. Instead, decomposition or intermolecular [2+2] photocycloaddition reactions prevailed. In the latter event, two main products were identified (three examples, 30-43% yield) resulting either from a head-to-head [2+2]-photodimerization or from a two-fold [2+2] photocycloaddition of the enone to the olefin. The latter reaction sequence generated pentacyclic products with a central [1,5]dioxocane ring. The structure assignment of the two product types was corroborated by single crystal X-ray analysis.

INTRODUCTION

Like most intramolecular cycloadditions, the [2+2] photocycloaddition¹ benefits from an improved regioselectivity if the two reaction partners are linked by a suitable tether. Cyclic α,β -unsaturated ketones represent the most frequently used chromophores in these reactions² and the preferred position for attachment of a tether is the α - or the β -position.³ For an α -substituted enone **I** (Scheme 1) the number m of atoms in the tether governs the regioselectivity. A preference towards a parallel approach of the two olefins is observed if three or four atoms link the two olefinic groups ($m = 3, 4$) and the photocycloaddition products **II** are frequently referred to as straight products. If the tether is short ($m = 2$) photocycloaddition products **III** prevail in which the two olefins approach each other in a crossed fashion. The regioselectivity can be understood by the mechanistic course of the reaction⁴ which involves an initial attack of the olefin on the photoexcited enone to a 1,4-diradical intermediate (vide infra). The first step of the

cycloaddition is consequently a cyclization in which the formation of five-membered rings⁵ is preferred.

SCHEME 1. Regioselectivity in the Intramolecular [2+2] Photocycloaddition of α -Substituted Enones **I** and Possible Reaction Pathways of α -Allyloxy-substituted Enones **IV**



2-(2'-Alkenyloxy)cycloalk-2-enones with the general structure **IV** represent a relatively unexplored compound class in [2+2] photocycloaddition chemistry. Ikeda et al. performed experiments with three 2-allyloxycyclohex-2-enones as substrates employing a high-pressure mercury lamp as the light source and acetone as the solvent.⁶ Crossed products (general structure **V**) were obtained in 53-63% yield and subsequent transformations of these products were intensively studied. We have now looked at an expanded set of substrates with two main objectives: (a) Instead of UV light it was attempted to perform the photochemical reactions with visible light. To this end, a suitable sensitizer was required which absorbs in the visible range and which does not lead to any side reactions. While visible-light mediated, sensitized reactions of styrenes and related compounds have recently received broad attention,⁷ studies with cyclic enones have remained scarce.⁸ (b) It was expected that other reaction pathways might be feasible depending on the structure of the chromophore and the substitution pattern of the olefin. Most

importantly, [2+2] photodimerization and hydrogen abstraction⁹ were foreseen as competing reactions which would in turn lead to intriguing new structures. The results of our experimental work are summarized in this account.

RESULTS AND DISCUSSION

Figure 1 provides an overview about the compounds we employed in the present study. With regard to the olefin component, three 2'-alkenyloxy groups were employed, i.e. a 2'-propenyloxy (allyloxy) group (substrates **1**), a 2'-methyl-2'-propenyloxy (methallyloxy) group (substrates **2**), and a 3'-methyl-2'-butenyloxy (prenyloxy) group (substrates **3**). With regard to the enone component, some substituted cyclic enones (substrates **b**, **c**, **d**, **f**) were probed as starting materials apart from the unsubstituted cyclohex-2-enones (substrates **a**) and cyclopent-2-enones (substrates **e**). Cyclohex-2-enones with (*Z*)- and (*E*)-substituted 2'-alkenyloxy groups were also prepared but will be discussed in a later section.




FIGURE 1. Overview about the 2-(2'-alkenyloxy)cycloalk-2-enones **1-3** employed in this study

Since the synthesis of 2-(2'-alkenyloxy)-4,4-dimethylcyclohex-2-enones was low yielding only substrates **1d** and **2d** were synthesized but not **3d**. In general, three approaches towards the [2+2] photocycloaddition precursors were taken. Starting from cyclic 1,2-diketones the alkenyloxy

group was introduced either by condensation with the respective allylic alcohol under acidic conditions (method A: TsOH in cyclohexane, reflux)^{6,10} or by nucleophilic substitution of an allyl bromide via the enolate (method B: K₂CO₃ in DMF, room temperature).¹¹ Along these lines, 1,2-diketones **4a**, **4b**, **4e**, and **4f** (Figure 2) served as precursors for the respective 2-(2'-alkenyloxy)cycloalk-2-enones **1a-3a**, **1b-3b**, **1e-3e**, and **1f-3f**. For the synthesis of substituted cyclohexenones **1c-3c** and **1d**, **2d** epoxides **4c** and **4d** served as starting materials, which underwent nucleophilic substitution in position C2 when treated with an excess of allylic alcohol under basic conditions (method C: NaH or KOH as base).¹² As mentioned above the reaction gave low yields (6-12%) when applied to epoxide **4d** which limited the supply of starting material. Reasons for the inefficiency of the procedure were (a) difficulties in removing excess allylic alcohol and (b) formation of a side product (see the SI for further details) by dimerization of the epoxide. Since the emphasis of the study was on the [2+2] photocycloaddition chemistry alternative routes to the substituted enones were not pursued. With the exception of compounds **1a**, **2a**, and **1c**, 2-(2'-alkenyloxy)cycloalk-2-enones have not been previously employed in photochemical reactions.





FIGURE 2. Structure of starting materials **4** for the synthesis of 2-(2'-alkenyloxy)cycloalk-2-enones and structure of thioxanthene-9-one (**5**)

Preliminary photochemical studies were performed with substrate **1a**. Its triplet energy was determined from phosphorescence experiments (77 K, CH₂Cl₂) as $E_T = 235 \text{ kJ mol}^{-1}$. Since a wavelength of $\lambda = 400 \text{ nm}$ corresponds to an energy of ca. 300 kJ Es^{-1} ($1 \text{ Es} = 6.022 \times 10^{23}$ photons) it seemed feasible to promote the [2+2] photocycloaddition with visible light upon judicious choice of a sensitizer. In this regard, parent thioxanthen-9-one (**5**) and its derivatives represent an excellent option and have been nicely exploited by Booker-Milburn and co-workers in [2+2] photocycloaddition reactions on large scale.^{8b} In recent work on visible light-mediated reactions, we have also seen that catalyst loadings can be as low as 1 mol% for a thioxanthone sensitizer.¹³ Experiments with a selection of potential sensitizers (see the SI for details) confirmed the suitability of thioxanthen-9-one (**5**) as the catalyst in the planned reaction ($E_T = 268 \text{ kJ mol}^{-1}$).¹⁴ Dichloromethane was established as the preferred solvent and there was no improvement in selectivity if the reaction was performed at low temperature. The reactions were run on small scale (50-200 μmol substrate, $c = 10 \text{ mM}$) at a wavelength of $\lambda = 420 \text{ nm}$ (16 fluorescent lamps with an emission maximum at $\lambda = 420 \text{ nm}$, for the emission spectrum, see the SI) and yields of product **8a** (Scheme 2) were consistently around 70% irrespective of whether 2.5, 5, or 10 mol% of **5** were used. Indeed, the low molecular weight (212 Da) of thioxanthen-9-one (**5**) made it relatively difficult to measure the exact quantity. Iridium catalysts (2.5 mol%) with a much higher molecular weight performed similar to **5** and there was a correlation between the reaction time and the triplet energy of the catalyst. Sensitizers with a triplet energy $E_T > 250 \text{ kJ mol}^{-1}$ led to a complete conversion after 2.5 hours. The reaction time increased with a decrease in triplet energy. The reaction with Ir(ppy)₃ ($E_T = 231 \text{ kJ mol}^{-1}$)¹⁵ for example required five hours before no substrate was detected by TLC. The ruthenium complex Ru(bpy)₃(PF₆)₂ ($E_T = 193 \text{ kJ mol}^{-1}$)¹⁶ did not promote the reaction and there was also no reaction in the absence of a sensitizer.


SCHEME 2. Mechanistic Proposal for the Course of the Sensitized Intermolecular [2+2]

Photocycloaddition of Cyclohex-2-enone **1a**

The formation of crossed product **8a** can be easily understood by assuming an energy transfer from the triplet state T_1 of thioxanthone **5** to substrate **1a**. Intermediate **5** (T_1) is populated with high efficiency by direct excitation and rapid¹⁷ intersystem crossing (ISC). An exothermic triplet energy transfer occurs by a spin-allowed, mutual electron-electron exchange process, often referred to as sensitization or triplet sensitization.¹⁸ Upon energy transfer, the reactive triplet state **1a** (T_1) is generated which is $\pi\pi^*$ in character and which is prone to undergo double bond addition.⁴ Subsequently, 1,4-diradicals **6** and **7** are formed by five-membered ring closure⁵ and both represent potential precursors to product **8a**. The low reduction potentials of α,β -unsaturated enones ($E_{1/2}^{\text{red}} \leq -2.0 \text{ V}$)¹⁹ render any single electron transfer (photoredox) processes not viable.

Several allyloxy- and methallyloxy-substituted cycloalk-2-enones reacted similar to substrate **1a** and the results are summarized in Scheme 3. In order to secure a reproducible reaction course, 10 mol% of the sensitizer were used for small scale reactions. Some of the reactions were also performed on larger scale (1 mmol) and at a higher concentration ($c = 25 \text{ mM}$). In this case, only 2.5 mol% of the sensitizer were employed. Despite a longer reaction time, the chemoselectivity did not suffer and yields were as good as on small scale or even higher.

SCHEME 3. Intramolecular [2+2] Photocycloaddition of Various 2-(2'-Alkenyloxy)cycloalk-2-enones upon Irradiation with Visible Light




^a Yield if the reaction was performed on 1 mmol scale.

Crossed photocycloaddition products **8a-8c** and **9a-9c** were isolated cleanly without any indication for the formation of other side products. Among the 4,4-dimethylcyclohexenones **1d** and **2d**, only the latter substrate showed a photochemical reaction. The former substrate did not react and the starting material was recovered. Since quantities were limited, the low yield of product **9d** might be associated with loss of product during chromatographic purification. Both compounds **9d** and **9e** were difficult to detect on TLC which further impaired a quantitative isolation. Cyclopentenone **1e** was converted upon sensitized irradiation (cf. Scheme 3) but no defined products could be isolated. The same observation was made when the compound was irradiated directly at $\lambda = 350$ nm ($c = 10$ mM, CH_2Cl_2).

The suspected intermediacy of 1,4-diradicals in the course of the [2+2] photocycloaddition implied the possibility to perform stereoconvergent reactions²⁰ with 1,2-disubstituted olefins in

the tether. Substrates *E*-**10**/*Z*-**10** (Scheme 4) were obtained from an *E/Z*-mixture (d.r. = 50/50) of crotyl alcohol (2-butenol) by condensation with 1,2-diketone **4a** (method A). Sensitized irradiation delivered photocycloaddition product **11** as a single diastereoisomer accompanied by an olefinic impurity which was removed by treatment with 3,6-bis(methoxycarbonyl)-1,2,4,5-tetrazine.²¹ The yield of clean isolated product was 78% and its relative configuration was assigned based on NOESY spectra (see the SI for further details).

SCHEME 4. Stereoconvergent Formation of [2+2] Photocycloaddition Products **11** and **13**




The ethyl-substituted substrates **12** were obtained in diastereomerically pure form from (*E*)- and (*Z*)-3-penten-1-ol and diketone **4a**. The compounds were individually subjected to the irradiation conditions and delivered the same product **13**. In this case, the product was not completely homogenous but was contaminated by compound **14** (yield from *E*-**12**: quant., **13/14** = 82/18; yield from *Z*-**12**: 90%, **13/14** = 83/17). By-product formation is likely to occur from the intermediate 1,4-diradical by intramolecular hydrogen abstraction (vide infra). It was possible to remove the undesired compound by oxidation and the desired product **13** was obtained in 69% yield. The outcome of the reactions suggests that 1,4-biradicals related to **6** are the major intermediates in the [2+2] photocycloaddition to products **11** and **13**. Not only do they explain the

observed stereoconvergence of the reaction but they also account for the formation of the olefinic side products.

When cyclohexenones **3a-3c** with a trisubstituted tethered olefin were employed as starting materials, a cyclization/reduction pathway became dominant and [2+2] photocycloaddition products were not detected. The high yields achieved for products **16a-16c** (Scheme 5) render some synthetic utility to this method, which represents formally a conjugate addition to the β -carbon atom of the α,β -unsaturated enone. The assignment of the relative product configuration was suggested by $^1\text{H-NMR}$ coupling constants and NOESY experiments (see the SI for detail) and the selectivity of the process is remarkable as it allows to construct three contiguous stereogenic centers with a defined configuration.

SCHEME 5. Formation of Cyclization Products **16** from 2-Dimethylallyloxy-substituted Enones **3** via 1,4-Diradicals **15**



Mechanistically, the first step of the sequence occurs presumably in analogy to the C-C bond forming event previously mentioned (**1a** \rightarrow **6**), i.e. the triplet of **3** undergoes intramolecular addition to the tethered olefin by a five-membered ring cyclization. However, due to steric hindrance of the substituents (R^3 and CMe_2) bond formation occurs in a *trans* fashion. In the


intermediate bicyclic 1,4-diradical **15** cyclization to a cyclobutane by C-C bond formation is not feasible but an intramolecular hydrogen abstraction⁹ can readily occur that generates products **16**.

Like cyclopentenone **1e** in the [2+2] photocycloaddition study, the cyclopentenone with a prenyloxy substituent (**3e**) showed only decomposition under the conditions successfully employed for substrate **3a**. Also the cyclopentenones **1f-3f** with an additional methyl group in β -position displayed a reactivity that did not parallel the reactivity pattern of the comparable six-membered cyclic enones **1b-3b**. In this instance, however, defined products could be isolated albeit in only low to moderate yields. The preferred reaction pathway of the cyclopentenones **1f-3f** was a [2+2] photodimerization with the interesting twist that there are two double bonds available in each component. Along these lines, 2-(2'-butenyloxy)-3-methylcycloalk-2-enone (**1f**) produced a set of [2+2] photocycloaddition products from which two products **17** and **18** could be isolated (Scheme 6). Product **17** is the [2+2] photodimerization product resulting from addition of the two enone double bonds in what is commonly referred to as a *cis-anti-cis* addition.^{1,2} The *cis* fusion between the five-membered rings and the four-membered ring is to be expected due to the rigidity of the skeleton while the *anti* (*trans*) configuration of the cyclopentanone rings minimizes their steric repulsion. Based on its NMR data, the second product **18** was also symmetric and – as confirmed by its crystal structure – it displays an inversion center (S_2 symmetry).²² The crystal structure revealed that the bicyclo[3.2.0]heptane core is *cis* connected and all substituents within the cyclobutane are positioned in a *cis* fashion. The two constitutionally identical bicyclo[3.2.0]heptane fragments possess opposite absolute configuration. Given the low combined yield for compounds **17** and **18**, it is likely that other dimerization products were formed but could not be isolated in pure form. [2+2]

Photodimerization in a head-to-head fashion is common for cyclopentenones²³ while the

regioselectivity of the intermolecular head-to-head double bond addition of a monosubstituted olefin to an enone has less precedence.^{9d,23d,24}


SCHEME 6. [2+2] Photodimerization of Cyclopent-2-enone **1f** and Crystal Structure^a of Product



^a Ellipsoids in the ORTEP structure are displayed at the 50% probability level.

The overall product yield achieved in the [2+2] photodimerization of compound **2f** was slightly higher (44%) than observed for **1f**. Again two regioisomers were isolated one of which was the enone dimerization product **19** and the other the product of an enone/olefin addition. The configuration assignment for the latter product **20** was based on analogy to product **18**.

SCHEME 7. [2+2] Photodimerization of Cyclopent-2-enone **2f** to Products **19** and **20**



Remarkably, even the prenyloxy-substituted enone **3f** did not display the cyclization tendency observed with cyclohexenones **3a-3c**. Instead, the only product isolated from its sensitized irradiation was the head-to-head regioisomer **21** of a [2+2] photodimerization that was obtained in 34% yield. The C_2 -symmetric compound displayed the same relative configuration as the previously discussed enone dimers **17** and **19** and the structure assignment could in this specific case be secured by single crystal structure analysis (Figure 3).




FIGURE 3. Structure of the [2+2] photodimerization product **21** (ellipsoids in the ORTEP structure are displayed at the 50% probability level) obtained by sensitized irradiation of substrate **3f** (34% yield)

The difference between the cyclopentenone and the cyclohexenone chromophore in the photochemical reactivity of their 2-(2'-alkenyloxy)-substituted derivatives is remarkable. Only in a single case (product **9e**) an intramolecular attack of the tethered olefin occurred while with all other substrates there was no indication for such an attack, not even for a cyclization to a 1,4-diradical (in analogy to the formation of intermediate **15**). The cyclization seems retarded and intermolecular [2+2] photocycloaddition reactions become competitive. The observation is important because it allows to access cyclopentenone dimers without protecting olefinic side chains.

In summary, an array of structurally diverse products has been obtained by irradiation of the title compounds in the presence of a sensitizer. The photochemical reactions of all cyclohexenone derivatives proceeded in moderate to excellent yields and with a high degree of diastereoselectivity. Intramolecular reactions were observed exclusively either as [2+2] photocycloaddition or as a cyclization/hydrogen abstraction cascade. Up to four stereogenic centers can be established in a single reaction and the products hold promise as scaffolds which enable a further functionalization at several positions. The cyclopentenone derivatives were less prone to intramolecular reactions but underwent preferentially an intermolecular reaction leading to [2+2] photodimerization products.

EXPERIMENTAL SECTION

General Methods. All air and moisture sensitive reactions were carried out in flame-dried glassware under an argon atmosphere using standard *Schlenk* techniques. For moisture sensitive reactions, tetrahydrofuran (THF), diethylether (Et₂O) and dichloromethane (CH₂Cl₂) were dried using a solvent purification system. Cyclohexane was dried over neutral aluminum oxide and stored over 4 Å molecular sieves. Dry *N,N*-dimethylformamide (DMF) was obtained in the highest available purity stored over molecular sieves and used without further purification. For photochemical reactions, dry dichloromethane was degassed by three freeze-pump-thaw cycles. Cooling baths used were ice/water (0 °C) and dry ice/ethanol (−78 °C). Technical solvents [1,2-dichloroethane (DCE), pentane (P), Et₂O, CH₂Cl₂, methanol (MeOH), ethyl acetate (EtOAc), cyclohexane] were distilled prior to use. Flash column chromatography was performed on silica 60 (230-400 mesh) and thin layer chromatography (TLC) was performed on silica coated glass plates (silica 60 F₂₅₄) with detection by UV-light ($\lambda = 254$ nm) and/or by staining with KMnO₄ solution followed by heat treatment. Infrared spectra (IR) were recorded by attenuated total

reflection (ATR) technique and are reported as wave numbers $\tilde{\nu}$ (cm^{-1}). The following abbreviations for intensities were used: vs (very strong), s (strong), m (medium), w (weak). NMR spectra were recorded at room temperature on either a 300, 400 or 500 MHz nuclear magnetic resonance spectrometer. The ^1H NMR spectra were referenced to the residual solvent peak of either chloroform (7.26 ppm) or C_6D_6 (7.16 ppm), and the $^{13}\text{C}\{^1\text{H}\}$ NMR spectra were referenced either against the central peak of CDCl_3 (77.16 ppm) or the residual solvent peak of C_6D_6 (128.06 ppm). ^1H NMR spectra were reported as follows: chemical shift in parts per million (ppm), peak shape (s - singlet, d - doublet, t - triplet, q - quartet, quint. - quintet, sept. - septet, m = multiplet), coupling constant in Hertz (Hz), and integration. Apparent multiplets which occur as a result of coupling constant equality between magnetically non-equivalent protons are marked as virtual (*virt.*). Mass spectra were measured with a mass selective quadrupole detector (EI, 70 eV). HRMS data were determined at a double-focussing magnetic sector instrument (EI, 70 eV). UV/Vis Spectra were recorded using a precision cell made of quartz with a pathway of 1 mm.

General procedure 1 (Method A): To a solution of the respective 1,2-cyclohexadione (1.0 equiv.) in dry cyclohexane (40 mL/g starting material) was added *p*-TsOH \cdot H $_2$ O (3.0 mol%) and the corresponding allylic alcohol (4.0 equiv.). The reaction mixture was heated at reflux in a Dean-Stark apparatus for 15 – 17 h. The solution was allowed to cool to room temperature, washed with brine (2×50 mL/g), dried over Na_2SO_4 , and filtrated. The solvent was removed under reduced pressure and the crude product was purified using flash column chromatography (P/Et $_2$ O).

General procedure 2 (Method B): To a solution of the respective 1,2-cyclohexadione (1.0 equiv.) in dry DMF (40 mL/g starting material) was added K_2CO_3 (1.2 equiv.) and the

1
2
3 respective allylbromide (1.2 equiv.). The reaction mixture was stirred at room temperature for
4
5 4.5 – 21 h. The reaction was quenched by the addition of H₂O (20 mL/g). Et₂O (40 mL/g) was
6
7 added and the layers were separated. The organic layer was washed with H₂O (4 × 40 mL/g) and
8
9 brine (1 × 60 mL/g), dried over Na₂SO₄, and filtrated. The solvent was removed under reduced
10
11 pressure and the crude product was purified using flash column chromatography (P/Et₂O).
12
13
14
15
16

17 **General procedure 3 (Method C):** A round bottom flask charged with NaH (60 wt%,
18
19 1.2 equiv.) was cooled to 0 °C and the respective allylic alcohol (30 equiv.) was added dropwise.
20
21 The reaction mixture was stirred for 10 min and subsequently the corresponding epoxide
22
23 (1.0 equiv.) was added dropwise. The reaction mixture was allowed to warm to room temperature
24
25 and stirred for 20 – 24 h. The mixture was diluted with Et₂O (75 mL/g) and quenched by the
26
27 addition of H₂O (35 mL/g). The layers were separated and the organic layer was washed with
28
29 H₂O (1 × 35 mL/g) and brine (1 × 75 mL/g), dried over Na₂SO₄, and filtrated. The solvent was
30
31 removed under reduced pressure and the crude product was purified by flash column
32
33 chromatography (P/Et₂O).
34
35
36
37
38
39

40 **General procedure 4 (Method B):** To a solution of the respective 1,2-cyclopentadione
41
42 (1.0 equiv.) in dry DMF (40 mL/g starting material) was added K₂CO₃ (1.2 equiv.) and the
43
44 respective allylbromide (1.2 equiv.). The reaction mixture was stirred at room temperature for 17
45
46 – 22 h. The reaction was quenched by the addition of H₂O (20 mL/g). Et₂O (40 mL/g) was added
47
48 and the layers were separated. The organic layer was washed with H₂O (4 × 40 mL/g) and brine
49
50 (1 × 60 mL/g), dried over Na₂SO₄, and filtrated. The solvent was removed under reduced
51
52 pressure and the crude product was purified using flash column chromatography (P/Et₂O).
53
54
55
56
57
58
59
60

2-(Allyloxy)cyclohex-2-en-1-one (1a)

According to general procedure 1, compound **1a** was synthesized starting from **4a** (1.00 g, 8.92 mmol, 1.0 equiv.), allylic alcohol (2.43 mL, 2.07 g, 35.7 mmol, 4.0 equiv.) and *p*-TsOH·H₂O (51.3 mg, 270 μmol, 3.0 mol%). Purification by flash column chromatography (SiO₂, P/Et₂O = 5/1, UV) afforded the product (667 mg, 4.38 mmol, 49%) as a colorless oil. TLC (P/Et₂O = 3/1): *R_f* = 0.18 [UV, KMnO₄]. ¹H NMR (400 MHz, CDCl₃, 300 K): δ = 1.97 (*virt.* quint. ³*J* ≅ ³*J* = 6.2 Hz, 2H), 2.41 (td, ³*J* = 6.0, 4.7 Hz, 2H), 2.48 – 2.53 (m, 2H), 4.31 (*virt.* dt ³*J* = 5.5 Hz, ⁴*J* ≅ ⁴*J* = 1.5 Hz, 2H), 5.25 (ddt, ²*J* = 1.5 Hz, ³*J* = 10.6 Hz, ⁴*J* = 1.5 Hz, 1H), 5.33 (ddt, ²*J* = 1.5 Hz, ³*J* = 17.4 Hz, ⁴*J* = 1.5 Hz, 1H), 5.89 (t, ³*J* = 4.7 Hz, 1H), 5.98 (ddt, ³*J* = 17.4, 10.6, 5.5 Hz, 1H). ¹³C{¹H} NMR (101 MHz, CDCl₃, 300 K): δ = 23.1 (t), 24.7 (t), 39.0 (t), 68.8 (t), 118.1 (t), 118.7 (d), 133.1 (d), 150.5 (s), 194.5 (s). UV/vis (CH₂Cl₂, *c* = 0.5 mM): λ = 259 nm (ε = 3736 cm⁻¹ M⁻¹). Data of this compound were in accordance with the literature.^{6b}

2-(Allyloxy)-3-methylcyclohex-2-en-1-one (1b)

According to general procedure 2, compound **1b**^{12a,25} was synthesized starting from **4a** (500 mg, 3.96 mmol, 1.0 equiv.), allyl bromide (410 μL, 580 mg, 4.76 mmol, 1.2 equiv.) and K₂CO₃ (670 mg, 4.85 mmol, 1.2 equiv.). Purification by flash column chromatography (SiO₂, P/Et₂O = 9/1, UV) afforded the product (259 mg, 1.56 mmol, 39%) as a colorless oil. TLC (P/Et₂O = 3/1): *R_f* = 0.30 [UV, KMnO₄]. ¹H NMR (500 MHz, CDCl₃, 300 K): δ = 1.92 (s, 3H), 1.92 – 1.97 (m, 2H), 2.38 (t, ³*J* = 6.1 Hz, 2H), 2.41 – 2.47 (m, 2H), 4.34 (*virt.* dt, ³*J* = 6.0 Hz, ²*J* ≅ ⁴*J* = 1.4 Hz, 2H), 5.18 (*virt.* dt, ³*J* = 10.3 Hz, ²*J* ≅ ⁴*J* = 1.4 Hz, 1H), 5.28 (*virt.* dtd, ³*J* = 17.2 Hz, ²*J* ≅ ⁴*J* = 1.6 Hz, ⁴*J* = 1.5 Hz, 1H), 5.99 (ddt, ³*J* = 17.0, 10.3, 6.0 Hz, 1H). ¹³C{¹H} NMR (101 MHz, CDCl₃, 300 K): δ = 18.1 (q), 22.3 (t), 31.7 (t), 38.9 (t), 73.0 (t), 117.8 (t), 134.5 (d), 146.4 (s), 147.9 (s), 194.9 (s). IR (ATR): $\tilde{\nu}$ = 2928 (w), 1672 (vs), 1631 (m), 1431

(w), 1379 (w), 1304 (w), 1193 (s), 1153 (s), 985 (s), 926 (s). MS (EI, 70 eV): m/z (%) = 166 (80), 151 (100), 137 (13), 125 (17), 113 (38), 110 (93), 95 (82), 82 (78). HRMS (EI) m/z : $[M]^+$ Calcd. for $C_{10}H_{14}O_2$: 166.0982; Found: 166.0988. UV/vis (CH_2Cl_2 , $c = 0.5$ mM): $\lambda = 247$ nm ($\epsilon = 8212$ cm⁻¹ M⁻¹).

2-(Allyloxy)-5,5-dimethylcyclohex-2-en-1-one (1c)

According to general procedure 3, compound **1c**^{12a} was synthesized starting from epoxide **4c** (200 mg, 1.43 mmol, 1.0 equiv.), allylic alcohol (3.00 mL, 2.55 g, 43.9 mmol, 31 equiv.) and NaH (60 wt%, 72.0 mg, 1.80 mmol, 1.3 equiv.). Purification by flash column chromatography (SiO_2 , P/Et₂O = 9/1 → 6/1, UV) afforded the product (115 mg, 640 μ mol, 45%) as a colorless oil. TLC (P/Et₂O = 3/1): R_f = 0.31 [UV, $KMnO_4$]. ¹H NMR (500 MHz, $CDCl_3$, 300 K): δ = 1.06 (s, 6H), 2.31 (d, ³ J = 4.6 Hz, 2H), 2.37 (s, 2H), 4.32 (*virt.* dt, ³ J = 5.8 Hz, ² J \cong ⁴ J = 1.4 Hz, 2H), 5.25 (*virt.* dq, ³ J = 10.6 Hz, ² J \cong ⁴ J = 1.3 Hz, 1H), 5.33 (*virt.* dq, ³ J = 17.3 Hz, ² J \cong ⁴ J = 1.5 Hz, 1H), 5.73 (t, ³ J = 4.6 Hz, 1H), 5.98 (ddt, ³ J = 17.2, 10.8, 5.5 Hz, 1H). ¹³C{¹H} NMR (126 MHz, $CDCl_3$, 300 K): δ = 28.3 (q), 34.1 (s), 38.6 (t), 52.4 (t), 68.8 (t), 116.1 (d), 118.2 (t), 133.0 (d), 149.8 (s), 194.5 (s). IR (ATR): $\tilde{\nu}$ = 2959 (m), 1690 (vs), 1629 (m), 1467 (w), 1368 (w), 1165 (m), 999 (m). MS (EI, 70 eV): m/z (%) = 180 (100), 165 (10), 151 (10), 127(10), 124 (51), 109 (49), 95 (52), 83 (19). HRMS (EI) m/z : $[M]^+$ Calcd. for $C_{11}H_{16}O_2$: 180.1145; Found: 180.1145. UV/vis (CH_2Cl_2 , $c = 0.5$ mM): $\lambda = 261$ nm ($\epsilon = 5484$ cm⁻¹ M⁻¹).

2-(Allyloxy)-4,4-dimethylcyclohex-2-en-1-one (1d)

KOH (85 wt%, 94.2 mg, 1.43 mmol, 1.0 equiv.) was dissolved in allylic alcohol (1.00 mL, 854 mg, 14.7 mmol, 10 equiv.). A solution of epoxide **4d** (200 mg, 1.43 mmol, 1.0 equiv.) in allylic alcohol (1.5 mL, 1.28 g, 22.1 mmol, 15 equiv.) was then added dropwise and the reaction mixture was stirred at room temperature over night. Subsequently the reaction mixture was

diluted with 15 mL Et₂O and quenched by the addition of 15 mL H₂O. The layers were separated and the aqueous layer was extracted with Et₂O (2 × 15 mL). The combined organic layers were washed with brine (1 × 50 mL), dried over Na₂SO₄, filtrated and the solvent as well as the remaining allylic alcohol were removed under reduced pressure. After purification by flash column chromatography (SiO₂, P/Et₂O = 20/1 → 6/1, UV) the product **1d** (30.5 mg, 169 μmol, 12%) was obtained as a colorless oil. TLC (P/Et₂O = 3/1): *R_f* = 0.28 [UV, KMnO₄]. ¹H NMR (500 MHz, CDCl₃, 300 K): δ = 1.18 (s, 6H), 1.82 (t, ³*J* = 6.5 Hz, 2H), 2.55 (t, ³*J* = 6.5 Hz, 2H), 4.27 (*virt.* dt, ³*J* = 5.6 Hz, ⁴*J* ≅ ⁴*J* = 1.5 Hz, 2H), 5.24 (*virt.* dq, ³*J* = 10.4 Hz, ²*J* ≅ ⁴*J* = 1.4 Hz, 1H), 5.32 (*virt.* dq, ³*J* = 17.2 Hz, ²*J* ≅ ⁴*J* = 1.6 Hz, 1H), 5.59 (s, 1H), 5.97 (ddt, ³*J* = 17.3, 10.8, 5.6 Hz, 1H). ¹³C{¹H} NMR (126 MHz, CDCl₃, 300 K): δ = 29.2 (q), 33.0 (s), 35.0 (t), 36.0 (t), 68.7 (t), 118.2 (t), 128.7 (d), 132.9 (d), 148.1 (s), 194.1 (s). IR (ATR): $\tilde{\nu}$ = 2958 (m), 1690 (vs), 1619 (s), 1458 (w), 1362 (m), 1261 (m), 1200 (s), 1130 (s), 1107 (s), 1014 (m), 925 (w). MS (EI, 70 eV): *m/z* (%) = 180 (24), 165 (24), 137 (4), 124 (100), 109 (23), 96 (15), 83 (11). HRMS (EI) *m/z*: [M]⁺ Calcd. for C₁₁H₁₆O₂: 180.1145; Found: 180.1144. UV/vis (CH₂Cl₂, *c* = 0.5 mM): λ = 260 nm (ε = 7584 cm⁻¹ M⁻¹).

2-(Allyloxy)cyclopent-2-en-1-one (**1e**)

To a solution of 1,2-cyclopentadione (**4e**) (95.0 mg, 968 μmol, 1.0 equiv.) in 5.0 mL dry cyclohexane was added *p*-TsOH·H₂O (6.10 mg, 32.1 μmol, 3.0 mol%) and allylic alcohol (210 μL, 178 mg, 3.06 mmol, 3.2 equiv.). The reaction mixture was heated at reflux using a Dean-Stark apparatus for 23 h. The solution was allowed to cool to room temperature and the solvent was removed under reduced pressure. After purification by flash column chromatography (SiO₂, P/Et₂O = 3/1, UV) the product **1e**²⁵ (100 mg, 720 μmol, 74%) was obtained as a colorless oil. TLC (P/Et₂O = 3/1): *R_f* = 0.14 [UV, KMnO₄]. ¹H NMR (500 MHz, CDCl₃, 300 K): δ = 2.38 –

2.48 (m, 2H), 2.52 (ddd, $^3J = 6.4, 4.2, 2.9$ Hz, 2H), 4.43 (*virt.* dt, $^3J = 5.6$ Hz, $^4J \cong ^4J = 1.5$ Hz, 2H), 5.28 (*virt.* dq, $^3J = 10.6$ Hz, $^2J \cong ^4J = 1.4$ Hz, 1H), 5.36 (*virt.* dq, $^3J = 17.3$ Hz, $^2J \cong ^4J = 1.6$ Hz, 1H), 5.98 (ddt, $^3J = 17.3, 10.9, 5.6$ Hz, 1H), 6.39 (t, $^3J = 2.9$ Hz, 1H). $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz, CDCl_3 , 300 K): $\delta = 22.1$ (t), 33.2 (t), 70.8 (t), 118.7 (t), 128.3 (d), 132.4 (d), 156.4 (s), 202.7 (s). IR (ATR): $\tilde{\nu} = 2925$ (w), 1715 (vs), 1624 (m), 1337 (w), 1276 (w), 1119 (m), 1027 (w), 934 (w). MS (EI, 70 eV): m/z (%) = 138 (100), 109 (15), 95 (11), 82 (52). HRMS (EI) m/z : $[\text{M}]^+$ Calcd. for $\text{C}_8\text{H}_{10}\text{O}_2$: 138.0675; Found: 138.0677, $[\text{M}]^+$ Calcd. for $\text{C}_7^{13}\text{C}_1\text{H}_{10}\text{O}_2$: 139.0709; Found: 139.0712. UV/vis (CH_2Cl_2 , $c = 0.5$ mM): $\lambda = 248$ nm ($\epsilon = 5180 \text{ cm}^{-1} \text{ M}^{-1}$).

2-(Allyloxy)-3-methylcyclopent-2-en-1-on (1f)

According to general procedure 4, compound **1f**^{12a} was synthesized starting from **4f** (500 mg, 4.46 mmol, 1.0 equiv.), allyl bromide (460 μL , 650 mg, 5.35 mmol, 1.2 equiv.) and K_2CO_3 (740 mg, 5.35 mmol, 1.2 equiv.). Purification by flash column chromatography (SiO_2 , $\text{P/Et}_2\text{O} = 3/1$, UV) afforded the product (387 mg, 2.54 mmol, 55%) as a colorless oil. TLC ($\text{P/Et}_2\text{O} = 3/1$): $R_f = 0.24$ [UV, KMnO_4]. ^1H NMR (400 MHz, CDCl_3 , 300 K): $\delta = 1.98$ (s, 3H), 2.30 – 2.39 (m, 2H), 2.43 (*virt.* dtd, $^2J = 7.1$ Hz, $^3J \cong ^3J = 2.5$, $^3J = 1.2$ Hz, 2H), 4.67 (*virt.* dt, $^3J = 5.9$ Hz, $^2J \cong ^2J = 1.4$ Hz, 2H), 5.18 (*virt.* dq, $^3J = 10.4$ Hz, $^2J \cong ^4J = 1.3$ Hz, 1H), 5.29 (*virt.* dq, $^3J = 17.2$ Hz, $^2J \cong ^4J = 1.6$ Hz, 1H), 5.94 (ddt, $^3J = 17.2, 10.4, 5.8$ Hz, 1H). $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3 , 300 K): $\delta = 15.1$ (q), 27.6 (t), 33.1 (t), 70.9 (t), 117.9 (t), 134.3 (d), 151.7 (s), 155.6 (s), 203.3 (s). IR (ATR): $\tilde{\nu} = 2917$ (w), 1699 (vs), 1643 (m), 1442 (w), 1389 (w), 1333 (m), 1204 (m), 1094 (s), 984 (m), 928 (m). MS (EI, 70 eV): m/z (%) = 152 (49), 137 (76), 123 (19), 112 (9), 96 (22), 84 (16), 81 (8), 69 (26), 67 (15), 55 (19), 41 (100). HRMS (EI) m/z : $[\text{M}]^+$ Calcd.

for $C_9H_{12}O_2$: 152.0827; Found: 152.0832. UV/vis (CH_2Cl_2 , $c = 0.5$ mM): $\lambda = 246$ nm ($\epsilon = 10132$ cm^{-1} M^{-1}).

2-[(2-Methylallyl)oxy]cyclohex-2-en-1-one (2a)

According to general procedure 1, compound **2a** was synthesized starting from **4a** (1.00 g, 8.92 mmol, 1.0 equiv.), methallylic alcohol (3.00 mL, 2.57 g, 35.7 mmol, 4.0 equiv.) and *p*-TsOH·H₂O (52.3 mg, 275 μ mol, 3.0 mol%). Purification by flash column chromatography (SiO₂, P/Et₂O = 9/1 \rightarrow 4/1, UV) afforded the product (682 mg, 4.10 mmol, 46%) as a colorless oil. TLC (P/Et₂O = 3/1): $R_f = 0.19$ [UV, KMnO₄]. ¹H NMR (500 MHz, C₆D₆, 300 K): $\delta = 1.36$ (*virt.* quint. $^3J \cong ^3J = 6.1$ Hz, 2H), 1.66 (s, 3H), 1.74 (td, $^3J = 6.0$, 4.6 Hz, 2H), 2.05 – 2.21 (m, 2H), 3.90 (s, 2H), 4.86 (s, 1H), 5.11 (s, 1H), 5.30 (t, $^3J = 4.6$ Hz, 1H). ¹³C{¹H} NMR (75 MHz, C₆D₆, 300 K): $\delta = 19.8$ (q), 23.5 (t), 24.8 (t), 39.6 (t), 71.9 (t), 113.0 (t), 118.1 (d), 141.6 (s), 151.5 (s), 190.4 (s). UV/vis (CH_2Cl_2 , $c = 0.5$ mM): $\lambda = 259$ ($\epsilon = 4518$ cm^{-1} M^{-1}). Data of this compound were in accordance with the literature.^{6b}

3-Methyl-2-[(2-methylallyl)oxy]cyclohex-2-en-1-one (2b)

According to general procedure 2, compound **2b** was synthesized starting from **4b** (500 mg, 3.96 mmol, 1.0 equiv.), methallyl bromide (480 μ L, 640 mg, 4.76 mmol, 1.2 equiv.) and K₂CO₃ (0.660 mg, 4.76 mmol, 1.2 equiv.). Purification by flash column chromatography (SiO₂, P/Et₂O = 9/1, UV) afforded the product (414 mg, 2.29 mmol, 58%) as a colorless oil. TLC (P/Et₂O = 3/1): $R_f = 0.29$ [UV, KMnO₄]. ¹H NMR (500 MHz, CDCl₃, 300 K): $\delta = 1.82$ (s, 3H), 1.90 – 1.98 (m, 5H), 2.38 (t, $^3J = 6.1$ Hz, 2H), 2.44 (t, $^3J = 6.7$ Hz, 2H), 4.22 (s, 2H), 4.90 (*virt.* t, $^2J \cong ^4J = 2.0$ Hz, 1H), 5.02 (s, 1H). ¹³C{¹H} NMR (126 MHz, CDCl₃, 300 K): $\delta = 18.0$ (q), 19.9 (q), 22.3 (t), 31.7 (t), 38.9 (t), 75.7 (t), 112.8 (t), 142.0 (s), 146.1 (s), 148.2 (s), 194.9 (s). IR

(ATR): $\tilde{\nu}$ = 2920 (w), 1673 (vs), 1632 (m), 1433 (w), 1378 (m), 1304 (m), 1193 (s), 1151 (vs), 993 (m), 895 (m). MS (EI, 70 eV): m/z (%) = 180 (100), 165 (37), 137 (11), 123 (21), 110 (94), 95 (74). HRMS (EI) m/z : $[M]^+$ Calcd. for $C_{11}H_{16}O_2$: 180.1145; Found: 180.1145, $[M]^+$ Calcd. for $C_{10}^{13}C_1H_{16}O_2$: 181.1185; Found: 181.1178. UV/vis (CH_2Cl_2 , c = 0.5 mM): λ = 248 nm (ϵ = 7196 $cm^{-1} M^{-1}$).

5,5-Dimethyl-2-[(2-methylallyl)oxy]cyclohex-2-en-1-one (2c)

According to general procedure 3, compound **2c** was synthesized starting from epoxide **4c** (200 mg, 1.43 mmol, 1.0 equiv.), methallylic alcohol (2.40 mL, 2.06 g, 28.6 mmol, 20 equiv.) and NaH (60 wt%, 70.0 mg, 1.75 mmol, 1.2 equiv.). Purification by flash column chromatography (SiO_2 , P/Et₂O = 9/1, UV) afforded the product (102 mg, 527 μ mol, 37%) as a colorless oil. TLC (P/Et₂O = 3/1): R_f = 0.40 [UV, $KMnO_4$]. 1H NMR (500 MHz, $CDCl_3$, 300 K): δ = 1.06 (s, 6H), 1.76 (s, 3H), 2.29 (d, 3J = 4.6 Hz, 2H), 2.36 (s, 2H), 4.25 (s, 2H), 4.95 (s, 1H), 5.00 (s, 1H), 5.73 (t, 3J = 4.6 Hz, 1H). $^{13}C\{^1H\}$ NMR (126 MHz, $CDCl_3$, 300 K): δ = 19.4 (q), 28.3 (q), 34.2 (s), 38.5 (t), 52.4 (t), 71.6 (t), 112.9 (t), 116.3 (d), 140.5 (s), 149.9 (s), 194.3 (s). IR (ATR): $\tilde{\nu}$ = 2958 (m), 1691 (vs), 1629 (m), 1455 (w), 1368 (w), 1163 (m), 902 (w). MS (EI, 70 eV): m/z (%) = 194 (57), 179 (8), 151 (9), 138 (22), 127 (14), 109 (100), 95 (9), 83 (14). HRMS (EI) m/z : $[M]^+$ Calcd. for $C_{12}H_{18}O_2$: 194.1301; Found: 194.1302. UV/vis (CH_2Cl_2 , c = 0.5 mM): λ = 261 nm (ϵ = 4374 $cm^{-1} M^{-1}$).

4,4-Dimethyl-2-[(2-methylallyl)oxy]cyclohex-2-en-1-one (2d)

KOH (85 wt%, 59.9 mg, 908 μ mol, 1.0 equiv.) was dissolved in methallylic alcohol (1.00 mL, 851 mg, 11.8 mmol, 13 equiv.). A solution of epoxide **4d** (127 mg, 908 μ mol, 1.0 equiv.) in methallylic alcohol (1.0 mL, 0.851 mg, 11.8 mmol, 13 equiv.) was then added dropwise and the

reaction mixture was stirred at room temperature over night. Subsequently the reaction mixture was diluted with 15 mL Et₂O and quenched by the addition of 15 mL H₂O. The layers were separated and the aqueous layer was extracted with Et₂O (2 × 15 mL). The combined organic layers were washed with brine (1 × 50 mL), dried over Na₂SO₄, filtrated and the solvent as well as the remaining volatiles were removed under reduced pressure. After purification by flash column chromatography (SiO₂, P/Et₂O = 6/1) the product **2d** (11.0 mg, 56.6 μmol, 6%) was obtained as a colorless oil. TLC (P/Et₂O = 3/1): *R_f* = 0.37 [UV, KMnO₄]. ¹H NMR (500 MHz, CDCl₃, 300 K): δ = 1.17 (s, 6H), 1.76 (t, ⁴*J* = 1.2 Hz, 3H), 1.80 – 1.83 (m, 2H), 2.52 – 2.60 (m, 2H), 4.18 (s, 2H), 4.94 (*virt.* quint., ²*J* ≅ ⁴*J* = 1.3 Hz, 1H), 5.00 (*virt.* dq, ²*J* = 2.1 Hz, ⁴*J* ≅ ⁴*J* = 1.2 Hz, 1H), 5.60 (s, 1H). ¹³C{¹H} NMR (126 MHz, CDCl₃, 300 K): δ = 19.5 (q), 29.2 (q), 33.0 (s), 35.1 (t), 36.0 (t), 71.5 (t), 113.1 (t), 128.9 (d), 140.3 (s), 148.2 (s), 194.1 (s). IR (ATR): $\tilde{\nu}$ = 2958 (m), 1692 (vs), 1619 (s), 1455 (w), 1363 (m), 1201 (s), 1130 (s), 1109 (s), 1014 (m), 901 (w). MS (EI, 70 eV): *m/z* (%) = 194 (44), 179 (31), 151 (14), 138 (100), 125 (20), 109 (37), 97 (22), 83 (10). HRMS (EI) *m/z*: [M]⁺ Calcd. for C₁₂H₁₈O₂: 194.1301; Found: 194.1301, [M]⁺ Calcd. for C₁₁¹³C₁H₁₈O₂: 195.1335; Found: 195.1350. UV/vis (CH₂Cl₂, *c* = 0.5 mM): λ = 260 nm (ε = 6300 cm⁻¹ M⁻¹), 338 nm (ε = 912 cm⁻¹ M⁻¹).

2-[(2-Methylallyl)oxy]cyclopent-2-en-1-one (**2e**)

In analogy to general procedure 4, compound **2e**²³ was synthesized starting from **4e** (100 mg, 1.02 mmol, 1.0 equiv.), methallyl bromide (150 μL, 207 mg, 1.53 mmol, 1.5 equiv.) and K₂CO₃ (211 mg, 1.53 mmol, 1.5 equiv.) in 4.0 mL dry DMF. The reaction was quenched by the addition of 5 mL H₂O and 10 mL Et₂O. The organic layer was washed with H₂O (4 × 10 mL) and brine (1 × 20 mL), dried over Na₂SO₄, filtrated and the solvent was removed under reduced pressure. Purification by flash column chromatography (SiO₂, P/Et₂O = 3/1, UV) afforded the product

(55.1 mg, 360 μ mol, 35%) as a colorless oil. TLC (P/Et₂O = 1/1): R_f = 0.30 [UV, KMnO₄]. ¹H NMR (500 MHz, CDCl₃, 300 K): δ = 1.78 (s, 3H), 2.40 – 2.44 (m, 2H), 2.50 (dt, ³ J = 6.1, 2.8 Hz, 2H), 4.34 (s, 2H), 4.98 (s, 1H), 5.03 (s, 1H), 6.40 (t, ³ J = 2.8 Hz, 1H). ¹³C{¹H} NMR (101 MHz, CDCl₃, 300 K): δ = 19.4 (q), 22.1 (t), 33.2 (t), 73.8 (t), 113.7 (t), 128.5 (d, C-3), 140.0 (s), 156.5 (s), 202.6 (s). IR (ATR): $\tilde{\nu}$ = 2922 (w), 1708 (vs), 1623 (s), 1407 (w), 1336 (w), 1273 (m), 1104 (vs), 1025 (m), 902 (m), 782 (m). MS (EI, 70 eV): m/z (%) = 152 (100), 137 (16), 124 (10), 109 (21), 96 (90), 82 (19). HRMS (EI) m/z : [M]⁺ Calcd. for C₉H₁₂O₂: 152.0832; Found: 152.0831. UV/vis (CH₂Cl₂, c = 0.5 mM): λ = 248 nm (ϵ = 6794 cm⁻¹ M⁻¹).

3-Methyl-2-[(2-methylallyl)oxy]cyclopent-2-en-1-one (2f)

According to general procedure 4, compound **2f** was synthesized starting from **4f** (500 mg, 4.46 mmol, 1.0 equiv.), methallyl bromide (550 μ L, 736 mg, 5.46 mmol, 1.2 equiv.) and K₂CO₃ (740 mg, 5.35 mmol, 1.2 equiv.). Purification by flash column chromatography (SiO₂, P/Et₂O = 9/1, UV) afforded the product (621 mg, 3.74 mmol, 84%) as a colorless oil. TLC (P/Et₂O = 3/1): R_f = 0.28 [UV, KMnO₄]. ¹H NMR (300 MHz, C₆D₆, 300 K): δ = 1.57 – 1.59 (m, 3H), 1.56 – 1.67 (m, 2H), 1.64 – 1.71 (m, 3H), 1.85 – 1.90 (m, 2H), 4.79 – 4.86 (m, 2H), 4.86 (virt. ddq, ² J = 2.3 Hz, ⁴ J = 1.6, ⁴ J \cong ⁴ J = 0.8 Hz, 1H), 5.11 (virt. dq, ² J = 2.3 Hz, ⁴ J \cong ⁴ J = 1.1 Hz, 1H). ¹³C{¹H} NMR (75 MHz, C₆D₆, 300 K): δ = 14.4 (q), 19.4 (q), 26.8 (t), 33.1 (t), 73.0 (t), 112.7 (t), 142.5 (s), 151.8 (s), 152.1 (s), 201.4 (s). IR (ATR): $\tilde{\nu}$ = 2917 (w), 1700 (vs), 1646 (m), 1448 (w), 1389 (m), 1335 (m), 1204 (m), 1096 (s), 990 (m), 903 (m). MS (EI, 70 eV): m/z (%) = 166 (56), 151 (31), 137 (10), 110 (9), 96 (12), 84 (15), 69 (16), 55 (100), 41 (24). HRMS (EI) m/z : [M]⁺ Calcd. for C₁₀H₁₄O₂: 166.0988; Found: 166.0985, [M]⁺ Calcd. for C₉¹³C₁H₁₄O₂: 167.1022; Found: 167.1026. UV/vis (CH₂Cl₂, c = 0.5 mM): λ = 248 nm (ϵ = 9284 cm⁻¹ M⁻¹).

2-[(3-Methylbut-2-en-1-yl)oxy]cyclohex-2-en-1-one (3a)

According to general procedure 2, compound **3a** was synthesized starting from **4a** (1.00 g, 8.92 mmol, 1.0 equiv.), 3,3-dimethylallyl bromide (1.24 mL, 1.60 g, 10.7 mmol, 1.2 equiv.) and K_2CO_3 (1.48 g, 10.7 mmol, 1.2 equiv.). Purification by flash column chromatography (SiO_2 , $P/Et_2O = 9/1 \rightarrow 4/1 \rightarrow 3/1$, UV) afforded the product (227 mg, 1.26 mmol, 14%) as a colorless oil. TLC ($P/Et_2O = 3/1$): $R_f = 0.15$ [UV, $KMnO_4$]. 1H NMR (400 MHz, $CDCl_3$, 300 K): $\delta = 1.64$ (s, 3H), 1.71 (s, 3H), 1.86 – 1.99 (m, 2H), 2.39 (td, $^3J = 6.1$, 4.7 Hz, 2H), 2.47 (dd, $^3J = 7.4$, 6.0 Hz, 2H), 4.23 (d, $^3J = 5.9$ Hz, 2H), 5.38 (*virt.* ddq, $^3J = 8.2$, 5.9 Hz, $^4J \cong ^4J = 1.5$ Hz, 1H), 5.85 (t, $^3J = 4.7$ Hz, 1H). $^{13}C\{^1H\}$ NMR (101 MHz, $CDCl_3$, 300 K): $\delta = 18.2$ (q), 23.0 (t), 24.6 (t), 25.8 (q), 38.9 (t), 64.6 (t), 117.9 (d), 119.6 (d), 137.6 (s), 150.7 (s), 194.5 (s). IR (ATR): $\tilde{\nu} = 2926$ (w), 1686 (vs), 1622 (m), 1444 (w), 1375 (w), 1259 (m), 1184 (s), 1148 (vs), 1003 (m), 874 (w). MS (EI, 70 eV): m/z (%) = 180 (27), 147 (11), 134 (8), 118 (57), 112 (100), 99 (12), 95 (7), 84 (27). HRMS (EI) m/z : $[M]^+$ Calcd. for $C_{11}H_{16}O_2$: 180.1148; Found: 180.1145, $[M]^+$ Calcd. for $C_{10}^{13}C_1H_{16}O_2$: 181.1185; Found: 181.1178. UV/vis (CH_2Cl_2 , $c = 0.5$ mM): $\lambda = 261$ nm ($\epsilon = 4126$ cm $^{-1}$ M $^{-1}$).

3-Methyl-2-[(3-methylbut-2-en-1-yl)oxy]cyclohex-2-en-1-one (3b)

According to general procedure 2, compound **3b**²⁶ was synthesized starting from **4b** (500 mg, 3.96 mmol, 1.0 equiv.), 3,3-dimethylallyl bromide (550 μ L, 710 mg, 4.76 mmol, 1.2 equiv.) and K_2CO_3 (0.660 mg, 4.76 mmol, 1.2 equiv.). Purification by flash column chromatography (SiO_2 , $P/Et_2O = 9/1 \rightarrow 3/1$, UV) afforded the product (197 mg, 1.01 mmol, 26%) as a colorless oil. TLC ($P/Et_2O = 3/1$): $R_f = 0.24$ [UV, $KMnO_4$]. 1H NMR (500 MHz, $CDCl_3$, 300 K): $\delta = 1.68$ (s, 3H), 1.75 (s, 3H), 1.86 – 2.03 (m, 5H), 2.38 (t, $^3J = 6.1$ Hz, 2H), 2.41 – 2.45 (m, 2H), 4.32 (d, $^3J = 7.4$ Hz, 2H), 5.44 (*virt.* ddq, $^3J = 7.4$, 5.9 Hz, $^4J \cong ^4J = 1.4$ Hz, 1H). $^{13}C\{^1H\}$ NMR (126

MHz, CDCl₃, 300 K): δ = 18.1 (q), 22.3 (t), 26.0 (q), 31.6 (t), 38.9 (t), 68.3 (t), 120.7 (d), 138.2 (s), 146.5 (s), 148.0 (s), 195.1 (s). IR (ATR): $\tilde{\nu}$ = 2917 (w), 1673 (vs), 1630 (m), 1433 (w), 1381 (m), 1301 (m), 1191 (s), 1148 (s), 970 (m). MS (EI, 70 eV): m/z (%) = 126 (100), 98 (11), 84 (23). HRMS (EI) m/z : [M]⁺ Calcd. for C₁₂H₁₈O₂: 194.1300; Found: 194.1301. UV/vis (CH₂Cl₂, c = 0.5 mM): λ = 248 nm (ϵ = 6640 cm⁻¹ M⁻¹).

5,5-Dimethyl-2-[(3-methylbut-2-en-1-yl)oxy]cyclohex-2-en-1-one (3c)

According to general procedure 3, compound **3c** was synthesized starting from epoxide **4c** (200 mg, 1.43 mmol, 1.0 equiv.), 3,3-dimethylallylic alcohol (4.50 mL, 3.83 g, 44.4 mmol, 31 equiv.) and NaH (60 wt%, 69.0 mg, 1.73 mmol, 1.2 equiv.). Purification by flash column chromatography (SiO₂, P/Et₂O = 9/1, P/Et₂O = 9/1 → 6/1, UV) afforded the product (45.2 mg, 217 μ mol, 15%) as a colorless oil. TLC (P/Et₂O = 3/1): R_f = 0.28 [UV, KMnO₄]. ¹H NMR (500 MHz, CDCl₃, 300 K): δ = 1.06 (s, 6H), 1.68 (s, 3H), 1.75 (s, 3H), 2.31 (d, ³ J = 4.6 Hz, 2H), 2.36 (s, 2H), 4.28 (d, ³ J = 7.3 Hz, 2H), 5.42 (ddd, ³ J = 7.3 Hz, ⁴ J = 4.1, 1.5 Hz, 1H), 5.71 (t, ³ J = 4.6 Hz, 1H). ¹³C{¹H} NMR (126 MHz, CDCl₃, 300 K): δ = 18.3 (q), 25.9 (q), 28.3 (q), 34.1 (s), 38.6 (t), 52.4 (t), 64.6 (t), 115.3 (d), 119.6 (d), 137.7 (s), 150.1 (s), 194.5 (s). IR (ATR): $\tilde{\nu}$ = 2958 (m), 1691 (vs), 1627 (m), 1452 (w), 1368 (w), 1162 (s), 1108 (m), 1003 (w). MS (EI, 70 eV): m/z (%) = 208 (28), 175 (12), 146 (30), 140 (100), 125 (49), 112 (10), 98 (71), 84 (57). HRMS (EI) m/z : [M]⁺ Calcd. for C₁₃H₂₀O₂: 208.1458; Found: 208.1445, [M]⁺ Calcd. for C₁₂¹³C₁H₂₀O₂: 209.1491; Found: 209.1482. UV/vis (CH₂Cl₂, c = 0.5 mM): λ = 263 nm (ϵ = 4500 cm⁻¹ M⁻¹).

2-[(3-Methylbut-2-en-1-yl)oxy]cyclopent-2-en-1-one (3e)

In analogy to general procedure 4, compound **3e** was synthesized starting from **4e** (170 mg, 1.73 mmol, 1.0 equiv.), 3,3-dimethylallyl bromide (310 μ L, 340 mg, 2.68 mmol, 1.5 equiv.) and K_2CO_3 (377 mg, 2.73 mmol, 1.6 equiv.) in 7.0 mL dry DMF. The reaction was quenched by the addition of 7 mL H_2O and 15 mL Et_2O . The organic layer was washed with H_2O (4×15 mL) and brine (1×30 mL), dried over Na_2SO_4 , filtrated and the solvent was removed under reduced pressure. Purification by flash column chromatography (SiO_2 , $P/Et_2O = 3/1$, UV) afforded the product (60.0 mg, 360 μ mol, 21%) as a colorless solid. Mp 42 – 43 $^{\circ}C$. TLC ($P/Et_2O = 1/1$): $R_f = 0.35$ [UV, $KMnO_4$]. 1H NMR (500 MHz, $CDCl_3$, 300 K): $\delta = 1.70$ (s, 3H), 1.76 (s, 3H), 2.41 – 2.44 (m, 2H), 2.48 – 2.60 (m, 2H), 4.40 (d, $^3J = 6.8$ Hz, 2H), 5.43 (t, $^3J = 6.8$ Hz, 1H), 6.36 (virt. q, $^3J \cong ^4J = 2.5$ Hz, 1H). $^{13}C\{^1H\}$ NMR (126 MHz, $CDCl_3$, 300 K): $\delta = 18.3$ (q), 22.2 (t), 25.9 (q), 33.2 (t), 66.7 (t), 119.1 (d), 127.6 (d), 138.7 (s), 156.6 (s), 202.9 (s). IR (ATR): $\tilde{\nu} = 2924$ (w), 1715 (vs), 1623 (s), 1449 (w), 1342 (w), 1275 (w), 1110 (s), 1026 (w), 964 (w), 782 (m). MS (EI, 70 eV): m/z (%) = 166 (59), 148 (21), 120 (11), 110 (23), 106 (28), 99 (100). HRMS (EI) m/z : $[M]^+$ Calcd. for $C_{10}H_{14}O_2$: 166.0988; Found: 166.0983, $[M]^+$ Calcd. for $C_9^{13}C_1H_{14}O_2$: 167.1022; Found: 167.1021. UV/vis (CH_2Cl_2 , $c = 0.5$ mM): $\lambda = 250$ nm ($\epsilon = 5976$ $cm^{-1} M^{-1}$).

3-Methyl-2-[(3-methylbut-2-en-1-yl)oxy]cyclopent-2-en-1-one (3f)

According to general procedure 4, compound **3f**²⁶ was synthesized starting from **4f** (500 mg, 4.46 mmol, 1.0 equiv.), 3,3-dimethylallyl bromide (620 μ L, 800 mg, 5.35 mmol, 1.2 equiv.) and K_2CO_3 (750 mg, 5.42 mmol, 1.2 equiv.). Purification by flash column chromatography (SiO_2 , $P/Et_2O = 9/1$, UV) afforded the product (558 mg, 3.09 mmol, 69%) as a colorless oil. TLC ($P/Et_2O = 3/1$): $R_f = 0.26$ [UV, $KMnO_4$]. 1H NMR (300 MHz, C_6D_6 , 300 K): $\delta = 1.55$ (dd, $^4J = 1.4, 0.8$ Hz, 6H), 1.60 – 1.62 (m, 3H), 1.63 – 1.67 (m, 2H), 1.82 – 1.99 (m, 2H), 4.97 (virt.

ddquint., $^3J = 7.1$ Hz, $^4J \cong ^4J = 0.9$ Hz, 2H), 5.52 (*virt.* ddquint., $^3J = 8.4$, 5.6 Hz, $^4J \cong ^4J = 1.4$ Hz, 1H). $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, C_6D_6 , 300 K): $\delta = 14.5$ (q), 18.0 (q), 25.8 (q), 33.1 (t), 66.4 (t), 121.9 (d), 137.4 (s), 152.2 (s), 152.7 (s), 201.8 (s). IR (ATR): $\tilde{\nu} = 2915$ (w), 1700 (vs), 1645 (m), 1443 (w), 1390 (m), 1338 (m), 1205 (m), 1092 (s), 969 (m), 905 (m). MS (EI, 70 eV): m/z (%) = 112 (90), 84 (32), 69 (100), 53 (9), 41 (74). HRMS (EI) m/z : $[\text{M}]^+$ Calcd. for $\text{C}_{11}\text{H}_{16}\text{O}_2$: 180.1145; Found: 180.1143. UV/vis (CH_2Cl_2 , $c = 0.5$ mM): $\lambda = 247$ nm ($\epsilon = 9532$ $\text{cm}^{-1} \text{M}^{-1}$).

3-Ethoxy-5,5-dimethylcyclohex-2-en-1-one

According to a modified literature procedure:²¹ To a solution of dimedone (2.10 g, 15.0 mmol, 1.0 equiv.) in 46 mL dry toluene were added EtOH (3.56 mL, 2.80 g, 61.0 mmol, 4.0 equiv.) and *p*-TsOH·H₂O (139 mg, 0.73 mmol, 5.0 mol%). The reaction mixture was heated at reflux using a water separator for 90 min at which point EtOH (3.00 mL, 2.37 g, 50.3 mmol, 3.4 equiv.) was added. Subsequently, the solution was refluxed for additional 14 h. It was allowed to cool to room temperature and the solvent was removed under reduced pressure. The crude product was then filtered over a short plug of Al_2O_3 (Et_2O , UV) and the product (2.52 g, 15.0 mmol, 100%) was obtained as a pale yellow oil. TLC ($\text{P}/\text{Et}_2\text{O} = 3/1$): $R_f = 0.17$ [UV, KMnO_4]. ^1H NMR (300 MHz, CDCl_3 , 300 K): $\delta = 1.07$ (s, 6H), 1.36 (t, $^3J = 7.0$ Hz, 3H), 2.21 (s, 2H), 2.27 (s, 2H), 3.90 (q, $^3J = 7.0$ Hz, 2H), 5.35 (s, 1H). $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, CDCl_3 , 300 K): $\delta = 14.3$ (q), 28.4 (q), 32.6 (t), 43.1 (t), 50.8 (t), 64.4 (t), 101.6 (d), 176.5 (s), 199.8 (s). Data of this compound were in accordance to the literature.²¹

5,5-Dimethylcyclohex-2-en-1-one

According to a modified literature procedure:²¹ To an ice-cooled suspension of LiAlH_4 (138 mg, 3.63 mmol, 0.4 equiv.) in 4.0 mL dry THF a solution of **S1** (1.50 g, 8.92 mmol, 1.0 equiv.) in

7.5 mL dry THF was added dropwise. Subsequently, the reaction mixture was allowed to warm to room temperature and stirred for 3 h. The reaction mixture was then cooled to 0 °C and quenched by the addition of MeOH (approx. 10 mL) until no further gas evolution was observed. After addition of 14 mL 1 M HCl the reaction mixture was stirred for 30 min, the layers were separated and the aqueous layer was extracted with Et₂O (3 × 30 mL). The combined organic layers were washed with brine (2 × 60 mL), dried over Na₂SO₄, filtrated and the solvent was removed under reduced pressure. After purification by flash column chromatography (SiO₂, P/Et₂O = 8/1, UV) the product (663 mg, 5.34 mmol, 60%) was obtained as a colorless oil. TLC (P/Et₂O = 1/1): *R_f* = 0.29 [UV, KMnO₄]. ¹H NMR (300 MHz, CDCl₃, 300 K): δ = 1.05 (s, 6H, CH₃), 2.24 (dd, ³*J* = 4.1 Hz, ⁴*J* = 2.1 Hz, 2H, H-4), 2.27 (s, 2H, H-6), 6.02 (dt, ³*J* = 10.1 Hz, ⁴*J* = 2.1 Hz, 1H, H-2), 6.86 (dt, ³*J* = 10.1, 4.1 Hz, 1H, H-3). ¹³C{¹H} NMR (75 MHz, CDCl₃, 300 K): δ = 28.5 (q), 34.0 (s), 40.0 (t), 51.9 (t), 129.1 (d), 148.5 (d), 200.1 (s). Data of this compound were in accordance to the literature.²¹

(1*SR*,6*SR*)-4,4-Dimethyl-7-oxabicyclo[4.1.0]heptan-2-one (4c)

According to a modified literature procedure:²⁷ To an ice-cooled solution of 5,5-dimethylcyclohex-2-en-1-one²¹ (990 mg, 7.97 mmol, 1.0 equiv.) in 16 mL MeOH were dropwise added H₂O₂ (30 wt%, 3.30 mL, 1.10 g, 32.3 mmol, 4.1 equiv.) and 6M NaOH (0.64 mL, 154 mg, 3.84 mmol, 0.5 equiv.). The reaction mixture was allowed to warm to room temperature and stirred for 2 h. Subsequently, the reaction mixture was diluted with 32 mL CH₂Cl₂ and H₂O was added until proper phase separation was achieved. The aqueous layer was extracted with CH₂Cl₂ (3 × 30 mL). The combined organic layers were washed with brine (2 × 80 mL), dried over Na₂SO₄, filtrated and the solvent was removed under reduced pressure. After purification by flash column chromatography (SiO₂, P/Et₂O = 2/1, KMnO₄) the product (901 mg, 6.43 mmol, 80%)

was obtained as a colorless oil. TLC (P/Et₂O = 2/1): R_f = 0.56 [KMnO₄]. ¹H NMR (400 MHz, CDCl₃, 300 K): δ = 0.92 (s, 3H), 1.06 (s, 3H), 1.74 – 1.89 (m, 2H), 2.04 (d, ² J = 14.3 Hz, 1H), 2.66 (d, ² J = 14.3 Hz, 1H), 3.21 (*virt. dt* ³ J = 3.8 Hz, ³ J \cong ³ J = 1.0 Hz, 1H), 3.51 (dd, ³ J = 3.8, 4.7 Hz, 1H). ¹³C{¹H} NMR (101 MHz, CDCl₃, 300 K): δ = 28.2 (q), 31.2 (q), 37.4 (t), 37.5 (t), 48.8 (s), 54.9 (d), 57.2 (d), 207.6 (s). Data of this compound were in accordance with the literature.²⁷

(1*SR*,6*SR*)-5,5-Dimethyl-7-oxabicyclo[4.1.0]heptan-2-one (4d)

According to a modified literature procedure:²⁷ To an ice-cooled solution of 4,4-dimethylcyclohex-2-en-1-one (2.00 g, 16.1 mmol, 1.0 equiv.) in 32 mL MeOH were dropwise added H₂O₂ (30 wt%, 8.23 mL, 80.5 mmol, 5.0 equiv.) and 6M NaOH (1.33 mL, 8.05 mmol, 0.5 equiv.). The reaction mixture was allowed to warm to room temperature and stirred for 3.5 h. Subsequently, the reaction mixture was diluted with 60 mL CH₂Cl₂ and H₂O was added until proper phase separation was achieved. The layers were separated and the aqueous layer was extracted with CH₂Cl₂ (3 \times 60 mL). The combined organic layers were washed with brine (1 \times 120 mL), dried over Na₂SO₄, filtrated and the solvent was removed under reduced pressure. After purification by flash column chromatography (SiO₂, P/Et₂O = 6/1, KMnO₄) the product (1.67 g, 11.9 mmol, 74%) was obtained as a colorless oil. TLC (P/Et₂O = 2/1): R_f = 0.68 [KMnO₄]. ¹H NMR (500 MHz, CDCl₃, 300 K): δ = 1.06 (s, 3H), 1.22 (s, 3H), 1.34 (dddd, ² J = 13.7 Hz, ³ J = 7.1, 3.0 Hz, ⁴ J = 1.3 Hz, 1H), 1.90 (ddd, ² J = 13.7 Hz, ³ J = 11.7, 6.4 Hz, 1H), 2.19 (ddd, ² J = 18.9 Hz, ³ J = 11.7, 7.0 Hz, 1H), 2.40 (ddd, ² J = 18.9 Hz, ³ J = 6.4, 3.0 Hz, 1H), 3.18 (dd, ³ J = 4.0 Hz, ⁴ J = 1.3 Hz, 1H), 3.23 (d, ³ J = 4.0 Hz, 1H). ¹³C{¹H} NMR (126 MHz, CDCl₃, 300 K): δ = 23.1 (q), 27.6 (q), 29.9 (t), 30.9 (s), 33.3 (t), 56.1 (d), 64.3 (d), 206.2 (s). Data of this compound were in accordance with the literature.²⁸

General procedure 5 (Irradiation): Thioxanthone **5** (10 mol%) was dissolved in 1.0 mL dry degassed CH₂Cl₂ and transferred to a flame dried Duran phototube. The respective 2-(2'-alkenyloxy)cycloalk-2-enone (1.0 equiv.) dissolved in 1.0 mL dry degassed CH₂Cl₂ was then added and the reaction mixture was diluted with dry degassed CH₂Cl₂ until a concentration of 10 mM (relative to the substrate) was reached. The reaction mixture was irradiated at room temperature in a previously described irradiation set-up²¹ ($\lambda = 420$ nm, for details about the light source, see the SI). Irradiation was continued until full conversion was reached and the solvent was removed under reduced pressure. The crude product was purified by flash column chromatography (P/Et₂O).

(3*RS*,3*aSR*,7*aSR*)-Hexahydro-7*H*-3,7*a*-methanobenzofuran-7-one (8a**)**

According to general procedure 5, compound **8a** was synthesized starting from **1a** (15.6 mg, 103 μ mol, 1.0 equiv.) and thioxanthone **5** (2.21 mg, 10.4 μ mol, 10 mol%) over 2.5 h. Purification by flash column chromatography (SiO₂, P/Et₂O = 1/2, KMnO₄) afforded the product (11.2 mg, 73.6 μ mol, 73%) as a colorless solid. The reaction on 1 mmol scale was performed with 152 mg **1a** (1.00 mmol, 1.0 equiv.) and 5.30 mg **5** (25.0 μ mol, 2.5 mol%) in 40 mL CH₂Cl₂ (25 mM, irradiation time: 5 h). The yield was 138 mg (0.91 mmol, 91%). Mp 59 – 65 °C. TLC (P/Et₂O = 1/2): R_f = 0.11 [KMnO₄]. ¹H NMR (500 MHz, CDCl₃, 300 K): δ = 1.64 – 1.78 (m, 1H), 1.81 – 1.96 (m, 2H), 2.03 – 2.18 (m, 2H), 2.30 (ddd, ³ J = 11.0, 7.8, 6.4 Hz, 1H), 2.35 – 2.44 (m, 2H), 2.84 (dd, ² J = 8.1 Hz, ³ J = 3.0 Hz, 1H), 2.88 (d, ³ J = 3.0 Hz, 1H), 3.85 (d, ³ J = 6.0 Hz, 1H), 3.94 (d, ³ J = 6.0 Hz, 1H). ¹³C{¹H} NMR (101 MHz, CDCl₃, 300 K): δ = 25.4 (t), 27.3 (t), 40.1 (t), 41.6 (t), 42.0 (d), 55.4 (d), 70.8 (t), 88.6 (s), 203.6 (s). Data of this compound were in accordance with the literature.^{6b}

(3*RS*,3*aSR*,7*aSR*)-3*a*-Methylhexahydro-7*H*-3,7*a*-methanobenzofuran-7-one (8*b*)

According to general procedure 5, compound **8b** was synthesized starting from **1b** (16.8 mg, 101 μ mol, 1.0 equiv.) and thioxanthone **5** (2.19 mg, 10.3 μ mol, 10 mol%) over 2 h. Purification by flash column chromatography (SiO₂, P/Et₂O = 1/2, KMnO₄) afforded the product (15.2 mg, 91.2 μ mol, 91%) as a colorless solid. Mp 38 – 47 °C. TLC (P/Et₂O = 1/2): *R_f* = 0.04 [KMnO₄]. ¹H NMR (400 MHz, CDCl₃, 300 K): δ = 0.99 (s, 3H), 1.76 (d, ²*J* = 8.2 Hz), 1.84 (ddt, ²*J* = 14.1 Hz, ³*J* = 4.2 Hz, ⁴*J* = 1.9 Hz, 1H), 2.00 (*virt.* qt, ²*J* \cong ³*J* = 13.9 Hz, ³*J* = 4.2 Hz, 1H), 2.08 – 2.18 (m, 1H), 2.28 (*virt.* td, ²*J* \cong ³*J* = 14.1 Hz, ³*J* = 4.3 Hz, 1H), 2.36 (ddt, ²*J* = 14.5 Hz, ³*J* = 4.1 Hz, ⁴*J* = 1.9 Hz, 1H), 2.49 (td, ²*J* = 14.5 Hz, ³*J* = 5.8 Hz, 1H), 2.72 (d, ³*J* = 3.0 Hz, 1H), 2.84 (dd, ²*J* = 8.2 Hz, ³*J* = 3.0 Hz, 1H), 3.86 (d, ²*J* = 6.7 Hz, 1H), 3.92 (d, ²*J* = 6.7 Hz, 1H). ¹³C{¹H} NMR (126 MHz, CDCl₃, 300 K): δ = 15.8 (q), 25.7 (t), 31.8 (t), 40.1 (t), 40.8 (t), 44.6 (d), 56.0 (s), 68.5 (t), 89.9 (s), 204.0 (s). IR (ATR): $\tilde{\nu}$ = 2925 (m), 1713 (s), 1458 (w), 1123 (w), 1043 (w), 945 (w), 862 (w). MS (EI, 70 eV): *m/z* (%) = 166 (100), 163 (21), 149 (39), 123 (35), 109 (62), 95 (44), 81 (100). HRMS (EI) *m/z*: [M]⁺ Calcd. for C₁₀H₁₄O₂: 166.0988; Found: 166.0982.

(3*RS*,3*aSR*,7*aSR*)-5,5-Dimethylhexahydro-7*H*-3,7*a*-methanobenzofuran-7-one (8*c*)

According to general procedure 5, compound **8c** was synthesized starting from **1c** (36.2 mg, 201 μ mol, 1.0 equiv.) and thioxanthone **5** (4.34 mg, 20.4 μ mol, 10 mol%) over 2.5 h. Purification by flash column chromatography (SiO₂, P/Et₂O = 3/1 \rightarrow 1/1, KMnO₄) afforded the product (36.2 mg, 201 μ mol, 100%) as a colorless solid. Mp 73 – 74 °C. TLC (P/Et₂O = 1/1): *R_f* = 0.14 [KMnO₄]. ¹H NMR (500 MHz, CDCl₃, 300 K): δ = 0.89 (s, 3H), 1.11 (s, 3H), 1.76 – 1.91 (m, 3H), 2.06 (dd, ²*J* = 14.2 Hz, ⁴*J* = 2.3 Hz, 1H), 2.34 – 2.45 (m, 2H), 2.81 (ddd, ²*J* = 8.2 Hz,

$^3J = 3.1$ Hz, $^4J = 1.1$ Hz, 1H), 2.84 (d, $^3J = 3.1$ Hz, 1H), 3.90 (d, $^2J = 6.0$ Hz, 1H), 3.97 (d, $^2J = 6.0$ Hz, 1H). $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz, CDCl_3 , 300 K): $\delta = 25.5$ (q), 31.4 (q), 38.2 (t), 39.9 (s), 41.0 (t), 41.3 (d), 52.8 (d), 53.1 (t), 71.6 (t), 88.5 (s), 203.4 (s). Data of this compound were in accordance with the literature.^{6b}

(3*RS*,3*aSR*,7*aSR*)-3-Methylhexahydro-7*H*-3,7*a*-methanobenzofuran-7-one (9a)

According to general procedure 5, compound **9a** was synthesized starting from **2a** (16.8 mg, 100 μmol , 1.0 equiv.) and thioxanthone **5** (2.24 mg, 10.6 μmol , 10 mol%) over 3 h. Purification by flash column chromatography (SiO_2 , $\text{P/Et}_2\text{O} = 1/2$, KMnO_4) afforded the product (15.3 mg, 92.0 μmol , 91%) as a colorless oil. TLC ($\text{P/Et}_2\text{O} = 1/2$): $R_f = 0.15$ [KMnO_4]. ^1H NMR (500 MHz, CDCl_3 , 300 K): δ [ppm] = 1.25 (s, 3H), 1.66 – 1.82 (m, 2H), 1.88 (*virt.* t, $^2J \cong ^4J = 7.9$ Hz, 1H), 1.95 – 2.04 (m, 1H), 2.14 (ddt, $^2J = 10.7$ Hz, $^3J = 6.0$, 2.9 Hz, 1H), 2.24 (*virt.* qd, $^3J \cong ^3J = 7.7$ Hz, $^4J = 3.7$ Hz, 1H), 2.31 – 2.44 (m, 2H), 2.59 (d, $^2J = 7.9$ Hz, 1H), 3.64 (dd, $^2J = 5.9$ Hz, $^4J = 1.2$ Hz, 1H), 3.74 (d, $^2J = 5.9$ Hz, 1H). $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3 , 300 K): δ [ppm] = 12.8 (q), 23.6 (t), 27.1 (t), 40.1 (t), 45.4 (t), 49.1 (s), 57.4 (d), 75.4 (t), 87.2 (s), 204.2 (s). Data of this compound were in accordance with the literature.^{6b}

(3*RS*,3*aSR*,7*aSR*)-3,3*a*-Dimethylhexahydro-7*H*-3,7*a*-methanobenzofuran-7-one (9b)

According to general procedure 5, compound **9b** was synthesized starting from **2b** (18.2 mg, 101 μmol , 1.0 equiv.) and thioxanthone **5** (2.14 mg, 10.1 μmol , 10 mol%) over 5 h. Purification by flash column chromatography (SiO_2 , $\text{P/Et}_2\text{O} = 1/2$, KMnO_4) afforded the product (15.9 mg, 88.3 μmol , 88%) as a colorless solid. The reaction on 1 mmol scale was performed with 180 mg **2b** (1.00 mmol, 1.0 equiv.) and 5.40 mg **5** (25.0 μmol , 2.5 mol%) in 40 mL CH_2Cl_2 (25 mM,

irradiation time: 9 h). The yield was 169 mg (0.94 mmol, 94%). Mp 67 – 70 °C. TLC (P/Et₂O = 1/2): R_f = 0.08 [KMnO₄]. ¹H NMR (500 MHz, CDCl₃, 300 K): δ = 0.90 (s, 3H), 1.14 (s, 3H), 1.65 (ddt, ² J = 13.5 Hz, ³ J = 3.9 Hz, ⁴ J = 1.8 Hz, 1H), 1.78 (d, ² J = 8.2 Hz, 1H), 1.92 – 2.06 (m, 1H), 2.07 – 2.20 (m, 2H), 2.28 – 2.36 (m, 1H), 2.38 – 2.51 (m, 1H), 2.62 (d, ² J = 8.2 Hz, 1H), 3.61 (d, ² J = 6.5 Hz, 1H), 3.66 (dd, ² J = 6.5 Hz, ⁴ J = 1.4 Hz, 1H). ¹³C{¹H} NMR (126 MHz, CDCl₃, 300 K): δ = 10.5 (q), 13.8 (q), 25.6 (t), 29.9 (t), 40.0 (t), 45.4 (t), 50.6 (s), 57.0 (s), 72.9 (t), 89.4 (s), 204.5 (s). IR (ATR): $\tilde{\nu}$ = 2922 (s), 1714 (vs), 1456 (w), 1385 (w), 1245 (w), 949 (w). MS (EI, 70 eV): m/z (%) = 180 (6), 165 (48), 135 (54), 125 (78), 110 (49), 97 (100), 83 (71). HRMS (EI) m/z : [M]⁺ Calcd. for C₁₁H₁₆O₂: 180.1145; Found: 180.1145.

(3*RS*,3*aSR*,7*aSR*)-3,5,5-Trimethylhexahydro-7*H*-3,7*a*-methanobenzofuran-7-one (9c)

According to general procedure 5, compound **9c** was synthesized starting from **2c** (38.9 mg, 200 μ mol, 1.0 equiv.) and thioxanthone **5** (4.36 mg, 20.5 μ mol, 10 mol%) over 2.5 h. Purification by flash column chromatography (SiO₂, P/Et₂O = 1/1, KMnO₄) afforded the product (35.1 mg, 181 μ mol, 90%) as a colorless solid. The reaction on 1 mmol scale was performed with 194 mg **2c** (1.00 mmol, 1.0 equiv.) and 5.30 mg **5** (25.0 μ mol, 2.5 mol%) in 40 mL CH₂Cl₂ (25 mM, irradiation time: 4.5 h). The yield was 178 mg (0.92 mmol, 92%). Mp 65 – 67 °C. TLC (P/Et₂O = 1/1): R_f = 0.19 [KMnO₄]. ¹H NMR (500 MHz, CDCl₃, 300 K): δ = 0.90 (s, 3H), 1.12 (s, 3H), 1.23 (s, 3H), 1.68 (ddd, ² J = 14.0 Hz, ³ J = 7.0 Hz, ⁴ J = 2.3 Hz, 1H), 1.75 (dd, ² J = 14.0 Hz, ³ J = 11.1 Hz, 1H), 1.88 (*virt.* t, ² J \cong ⁴ J = 7.9 Hz, 1H), 2.05 (dd, ² J = 14.2 Hz, ⁴ J = 2.3 Hz, 1H), 2.28 – 2.41 (m, 2H), 2.56 (dd, ² J = 8.0 Hz, ⁴ J = 1.2 Hz, 1H), 3.68 (dd, ² J = 5.9 Hz, ⁴ J = 1.2 Hz, 1H), 3.76 (d, ² J = 5.9 Hz, 1H). ¹³C{¹H} NMR (126 MHz, CDCl₃, 300 K): δ = 12.8 (q), 25.7 (q), 31.4 (q), 36.4 (t), 39.9 (s), 44.7 (t), 48.4 (s), 53.0 (t), 54.7 (d), 76.1 (t),

87.1 (s), 204.0 (s). IR (ATR): $\tilde{\nu}$ = 2954 (m), 1714 (vs), 1468 (w), 1245 (w), 1163 (w), 1082 (w), 964 (s). MS (EI, 70 eV): m/z (%) = 194 (25), 179 (9), 151 (16), 124 (10), 109 (100), 95 (16), 83 (15). HRMS (EI) m/z : $[M]^+$ Calcd. for $C_{12}H_{18}O_2$: 194.1301; Found: 194.1301.

(3*RS*,3*aSR*,7*aSR*)-3,4,4-Trimethylhexahydro-7*H*-3,7*a*-methanobenzofuran-7-one (9d)

According to general procedure 5, compound **9d** was synthesized starting from **2d** (11.0 mg, 56.6 μ mol, 1.0 equiv.) and thioxanthone **5** (1.20 mg, 5.66 μ mol, 10 mol%) over 7.5 h. Purification by flash column chromatography (SiO_2 , P/Et₂O = 3/1 \rightarrow 1/1, KMnO₄) afforded the product (4.20 mg, 21.6 μ mol, 38%) as a colorless oil. TLC (P/Et₂O = 1/1): R_f = 0.14 [KMnO₄]. ¹H NMR (500 MHz, CDCl₃, 300 K): δ = 1.17 (s, 3H), 1.32 – 1.41 (m, 6H), 1.66 – 1.75 (m, 2H), 1.88 (*virt.* t, $^2J \cong ^4J$ = 8.4 Hz, 1H), 2.01 (d, 4J = 8.5 Hz, 1H), 2.31 (dt, 2J = 16.2 Hz, 3J = 4.2 Hz, 1H), 2.50 (ddd, 2J = 16.2 Hz, 3J = 10.2, 7.4 Hz, 1H), 2.76 (dd, 2J = 8.1 Hz, 4J = 1.2 Hz, 1H), 3.61 (dd, 2J = 5.9 Hz, 4J = 1.2 Hz, 1H), 3.64 (d, 2J = 5.9 Hz, 1H). ¹³C{¹H} NMR (126 MHz, CDCl₃, 300 K): δ = 13.6 (q), 24.0 (q), 32.9 (s), 33.7 (q), 37.0 (t), 41.4 (t), 45.9 (t), 49.7 (s), 65.0 (d), 76.8 (t), 86.3 (s), 205.1 (s). IR (ATR): $\tilde{\nu}$ = 2954 (m), 1719 (vs), 1458 (w), 1062 (m), 948 (w). MS (EI, 70 eV): m/z (%) = 194 (8), 179 (33), 164 (19), 149 (34), 138 (100), 125 (38), 109 (36), 97 (34), 81 (43). HRMS (EI) m/z : $[M]^+$ Calcd. for $C_{12}H_{18}O_2$: 194.1301; Found: 194.1298.

(3*RS*,3*aSR*,6*aSR*)-3-Methyltetrahydro-3,6*a*-methanocyclopenta[*b*]furan-6(2*H*)-one (9e)

According to general procedure 5, compound **9e** was synthesized starting from **2e** (30.7 mg, 202 μ mol, 1.0 equiv.) and thioxanthone **5** (4.26 mg, 20.1 μ mol, 10 mol%) over 19 h. Purification by flash column chromatography (SiO_2 , P/Et₂O = 3/1, KMnO₄) afforded the product (10.4 mg, 68.3 μ mol, 34%) as a colorless oil. TLC (P/Et₂O = 3/1): R_f = 0.07 [KMnO₄]. ¹H NMR (500 MHz, CDCl₃, 300 K): δ = 1.29 (s, 3H), 1.95 (*virt.* t, $^2J \cong ^4J$ = 8.2 Hz, 1H), 2.02 (ddd, 3J = 9.2, 7.7,

5.8 Hz, 2H), 2.48 (*virt.* q, $^3J \cong ^3J = 8.3$ Hz, 1H), 2.57 (dt, $^2J = 19.2$ Hz, $^3J = 9.2$ Hz, 1H), 2.62 – 2.73 (m, 2H), 4.04 (dd, $^2J = 6.2$ Hz, $^4J = 1.3$ Hz, 1H), 4.08 (d, $^2J = 6.2$ Hz, 1H). $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz, CDCl_3 , 300 K): $\delta = 13.8$ (q), 16.9 (t), 41.3 (t), 44.2 (t), 49.0 (s), 51.3 (d), 83.1 (t), 93.0 (s), 206.3 (s). IR (ATR): $\tilde{\nu} = 2956$ (w), 1737 (vs), 1451 (w), 1238 (w), 1038 (m), 1006 (m), 975 (s), 919 (m), 894 (m). MS (EI, 70 eV): m/z (%) = 151 (6), 137 (36), 123 (13), 110 (100), 96 (70), 81 (92). HRMS (EI) m/z : $[\text{M}]^+$ Calcd. for $\text{C}_9\text{H}_{12}\text{O}_2$: 152.0832; Found: 152.0828.

(*E/Z*)-2-(But-2-en-1-yloxy)cyclohex-2-en-1-one (*E*-10/*Z*-10)

According to general procedure 1, compound *E*-10/*Z*-10²⁹ was synthesized starting from **4a** (1.00 g, 8.92 mmol, 1.0 equiv.), crotyl alcohol (*E/Z* = 1/1) (3.00 mL, 2.55 g, 35.4 mmol, 4.0 equiv.) and *p*-TsOH·H₂O (51.7 mg, 272 μmol , 3.0 mol%). Purification by flash column chromatography (SiO_2 , $\text{P/Et}_2\text{O} = 3/1$, UV) afforded the product (510 mg, 3.07 mmol, 34%) as a colorless oil. TLC ($\text{P/Et}_2\text{O} = 3/1$): $R_f = 0.14$ [UV, KMnO_4]. ^1H NMR (400 MHz, CDCl_3 , 300 K): $\delta = 1.72$ (dd, $^3J = 6.3$ Hz, $^4J = 1.1$ Hz, 3H), 1.96 (*virt.* quint. $^3J \cong ^3J = 6.2$ Hz, 2H), 2.41 (*virt.* td, $^3J \cong ^3J = 5.9$ Hz, $^3J = 4.6$ Hz, 2H), 2.50 (dd, $^3J = 7.4$, 6.0 Hz, 2H), 4.20 (d, $^3J = 6.3$ Hz, 2H), 5.62 – 5.71 (m, 1H), 5.71 – 5.83 (m, 1H), 5.88 (t, $^3J = 4.6$ Hz, 1H). $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3 , 300 K): $\delta = 17.9$ (q), 23.1 (t), 24.7 (t), 39.0 (t), 68.6 (t), 118.1 (d), 125.9 (d), 130.9 (d), 150.6 (s), 194.5 (s). IR (ATR): $\tilde{\nu} = 2939$ (w), 1686 (vs), 1622 (m), 1439 (w), 1365 (w), 1259 (m), 1185 (s), 1149 (vs), 1006 (m), 965 (s), 872 (w). MS (EI, 70 eV): m/z (%) = 166 (21), 112 (100), 95 (6), 84 (32). HRMS (EI) m/z : $[\text{M}]^+$ Calcd. for $\text{C}_{10}\text{H}_{14}\text{O}_2$: 166.0988; Found: 166.0988, $[\text{M}]^+$ Calcd. for $\text{C}_9^{13}\text{C}_1\text{H}_{14}\text{O}_2$: 167.1022; Found: 167.1025. UV/vis (CH_2Cl_2 , $c = 0.5$ mM): $\lambda = 260$ nm ($\epsilon = 4882 \text{ cm}^{-1} \text{ M}^{-1}$).

(3*RS*,3*aSR*,7*aSR*,8*SR*)-8-Methylhexahydro-7*H*-3,7*a*-methanobenzofuran-7-one (11)

In analogy to general procedure 5, compound **11** was synthesized starting from *E*-**10**/*Z*-**10** (16.6 mg, 100 μ mol, 1.0 equiv.) and thioxanthone **5** (2.15 mg, 10.1 μ mol, 10 mol%) over 4.5 h. The solvent was removed under reduced pressure and the residue redissolved in 1.0 mL CH_2Cl_2 and dimethyl 1,2,4,5-tetrazine-3,6-dicarboxylate (5.30 mg, 26.7 μ mol, 0.27 equiv.) was added.²¹ The reaction mixture was stirred at room temperature for 4.5 h and the solvent was removed under reduced pressure. Purification by flash column chromatography (SiO_2 , $\text{P/Et}_2\text{O}$ = 1/2, KMnO_4) afforded the product (13.0 mg, 78.2 μ mol, 78%) as a colorless solid. Mp 48 – 53 °C. TLC ($\text{P/Et}_2\text{O}$ = 1/2): R_f = 0.21 [KMnO_4]. ^1H NMR (500 MHz, CDCl_3 , 300 K): δ = 0.91 (d, 3J = 6.4 Hz, 3H), 1.62 – 1.76 (m, 1H), 1.91 (*virt.* tdd, $^2J \cong ^3J$ = 14.2 Hz, 3J = 11.0 Hz, 3J = 3.7 Hz, 1H), 2.00 – 2.07 (m, 1H), 2.11 (ddd, 2J = 13.4 Hz, 3J = 6.7, 3.6 Hz, 1H), 2.21 (dd, 3J = 11.0, 6.7 Hz, 1H), 2.28 – 2.41 (m, 2H), 2.65 (d, 3J = 2.9 Hz, 1H), 3.06 (qd, 3J = 6.4 Hz, 2.9 Hz, 1H), 3.76 (d, 2J = 6.5 Hz, 1H), 3.88 (d, 2J = 6.5 Hz, 1H). $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz, CDCl_3 , 300 K): δ = 9.7 (q), 24.4 (t), 27.1 (t), 40.1 (t), 44.4 (d), 45.0 (d), 52.7 (d), 67.5 (t), 89.1 (s), 203.8 (s). IR (ATR): $\tilde{\nu}$ = 2946 (m), 1710 (vs), 1451 (w), 1320 (w), 1094 (m), 994 (m), 930 (m), 848 (m). MS (EI, 70 eV): m/z (%) = 166 (20), 151 (4), 138 (5), 123 (5), 112 (100), 109 (15), 95 (11), 84 (41). HRMS (EI) m/z : [M]⁺ Calcd. for $\text{C}_{10}\text{H}_{14}\text{O}_2$: 166.0988; Found: 166.0982.

(*E*)-2-(Pent-2-en-1-yloxy)cyclohex-2-en-1-one (*E*-12)

According to general procedure 1, compound *E*-**12** was synthesized starting from **4a** (250 mg, 2.23 mmol, 1.0 equiv.), (*E*)-pent-2-en-1-ol (0.90 mL, 770 mg, 8.92 mmol, 4.0 equiv.) and *p*-TsOH· H_2O (12.7 mg, 70.0 μ mol, 3.0 mol%). Purification by flash column chromatography (SiO_2 , $\text{P/Et}_2\text{O}$ = 6/1 \rightarrow 3/1, $\text{P/Et}_2\text{O}$ = 6/1, UV) afforded the product (152 mg, 840 μ mol, 38%) as a colorless oil. TLC ($\text{P/Et}_2\text{O}$ = 3/1): R_f = 0.18 [UV, KMnO_4]. ^1H NMR (500 MHz, CDCl_3 , 300 K): δ = 1.00 (t, 3J = 7.5 Hz, 3H), 1.96 (dt, 2J = 12.4 Hz, 3J = 6.0 Hz, 2H), 2.02 – 2.13 (m,

2H), 2.41 (*virt.* td, $^3J \cong ^3J = 6.0$ Hz, $^3J = 4.6$ Hz, 2H), 2.50 (dd, $^3J = 7.4$, 6.0 Hz, 2H), 4.22 (dd, $^3J = 6.3$ Hz, $^4J = 1.4$ Hz, 2H), 5.64 (dt, $^3J = 15.5$, 6.3 Hz, $^4J = 1.6$ Hz, 1H), 5.75 – 5.85 (m, 1H), 5.88 (t, $^3J = 4.6$ Hz, 1H). $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz, CDCl_3 , 300 K): $\delta = 13.3$ (q), 23.1 (t), 24.7 (t), 25.5 (t), 39.0 (t), 68.7 (t), 118.2 (d), 123.5 (d), 137.6 (d), 150.5 (s), 194.6 (s). IR (ATR): $\tilde{\nu} = 2933$ (w), 1686 (vs), 1623 (m), 1458 (w), 1365 (w), 1259 (m), 1185 (s), 1149 (vs), 1123 (s), 1007 (m), 967 (s), 880 (w). MS (EI, 70 eV): m/z (%) = 180 (11), 145 (93), 129 (11), 113 (16), 99 (100), 83 (23). HRMS (EI) m/z : $[\text{M}]^+$ Calcd. for $\text{C}_{11}\text{H}_{16}\text{O}_2$: 180.1145; Found: 180.1143. UV/vis (CH_2Cl_2 , $c = 0.5$ mM): $\lambda = 260$ nm ($\epsilon = 4760$ $\text{cm}^{-1} \text{M}^{-1}$).

(Z)-1-Bromopent-2-ene³⁰

(Z)-Pent-2-en-1-ol (0.81 mL, 689 mg, 8.00 mmol, 1.0 equiv.) was dissolved in 16 mL dry Et₂O and cooled to 0 °C. PBr₃ (0.90 mL, 2.60 g, 9.60 mmol, 1.2 equiv.) was added dropwise, the reaction mixture was allowed to warm to room temperature and stirred for 4 h. The reaction was quenched by addition of 20 mL of H₂O and the layers were separated. The organic layer was washed with NaHCO₃ (1 × 20 mL), H₂O (1 × 20 mL) and brine (1 × 20 mL), dried over Na₂SO₄, filtrated and the solvent was removed under reduced pressure. After purification by flash column chromatography (SiO₂, P) the product (119 mg, 0.80 mmol, 10%) was obtained as a colorless oil. TLC (P): *R_f* = 0.67 [UV, KMnO₄]. ¹H NMR (500 MHz, CDCl₃, 300 K): δ = 1.03 (t, ³*J* = 7.5 Hz, 3H), 2.16 (*virt. quint.* d ³*J* ≅ ³*J* = 7.5 Hz, ⁴*J* = 1.5 Hz, 2H), 4.00 (dd, ³*J* = 8.3 Hz, ⁴*J* = 0.7 Hz, 2H), 5.60 (dt, ³*J* = 10.6, 7.4 Hz, 1H), 5.70 (m, 1H). ¹³C {¹H} NMR (75 MHz, CDCl₃, 300 K): δ = 13.9 (q), 20.4 (t), 27.4 (t), 124.8 (d), 137.7 (d). Data of this compound were in accordance to the literature.³⁰

(Z)-2-(Pent-2-en-1-yloxy)cyclohex-2-en-1-one (Z-12)

According to general procedure 2, compound Z-12 was synthesized starting from **4a** (520 mg, 4.64 mmol, 1.0 equiv.), (Z)-1-bromopent-2-ene (830 mg, 5.57 mmol, 1.2 equiv.) and K₂CO₃ (0.770 mg, 5.57 mmol, 1.2 equiv.). Purification by flash column chromatography (SiO₂, P/Et₂O = 3/1, H/EtOAc = 24/1 → 7/3 → 0/1 UV) afforded the product (133 mg, 737 μmol, 17%) as a colorless oil. TLC (P/Et₂O = 3/1): *R_f* = 0.19 [UV, KMnO₄]. ¹H NMR (500 MHz, CDCl₃, 300 K): δ = 0.99 (t, ³*J* = 7.5 Hz, 3H), 1.89 – 2.03 (m, 2H), 2.03 – 2.15 (m, 2H), 2.42 (*virt. td*, ³*J* ≅ ³*J* = 6.0 Hz, ³*J* = 4.6 Hz, 2H), 2.51 (dd, ³*J* = 7.5, 6.0 Hz, 2H), 4.36 (d, ³*J* = 5.1 Hz, 2H), 5.43 – 5.67 (m, 2H), 5.87 (t, ³*J* = 4.6 Hz, 1H). ¹³C {¹H} NMR (126 MHz, CDCl₃, 300 K): δ = 14.2 (q), 21.3 (t), 23.1 (t), 24.7 (t), 39.0 (t), 63.8 (t), 118.2 (d), 124.0 (d), 135.7 (d), 150.5 (s), 194.6

(s). IR (ATR): $\tilde{\nu}$ = 2933 (w), 1687 (vs), 1622 (m), 1457 (w), 1374 (w), 1259 (m), 1185 (m), 1149 (vs), 1123 (s), 1007 (m), 873 (w). MS (EI, 70 eV): m/z (%) = 180 (17), 145 (58), 129 (11), 113 (100), 99 (87), 84 (40). HRMS (EI) m/z : $[M]^+$ Calcd. for $C_{11}H_{16}O_2$: 180.1145; Found: 180.1145, $[M]^+$ Calcd. for $C_{10}^{13}C_1H_{16}O_2$: 181.1178; Found: 181.1183. UV/vis (CH_2Cl_2 , c = 0.5 mM): λ = 259 nm (ϵ = 3754 $cm^{-1} M^{-1}$).

(3*RS*,3*aSR*,7*aSR*,8*SR*)-8-Ethylhexahydro-7*H*-3,7*a*-methanobenzofuran-7-one (13)

According to general procedure 5, compound **13** was synthesized starting from *E*-**12** (18.1 mg, 100 μ mol, 1.0 equiv.) and thioxanthone **5** (2.14 mg, 10.2 μ mol, 10 mol%) over 2.5 h. Purification by flash column chromatography (SiO_2 , P/Et_2O = 2/1 \rightarrow 1/1, $KMnO_4$) afforded a mixture of product **13** and olefinic by-product **14** (18.1 mg, 100 μ mol, 100%, **13/14** = 82/18) as a colorless solid.

According to general procedure 5, compound **13** was synthesized starting from *Z*-**12** (18.1 mg, 100 μ mol, 1.0 equiv.) and thioxanthone **5** (2.17 mg, 10.1 μ mol, 10 mol%) over 2.5 h. Purification by flash column chromatography (SiO_2 , P/Et_2O = 1/1, $KMnO_4$) afforded a mixture of product **13** and olefinic by-product **14** (16.3 mg, 90.4 μ mol, 90%, **13/14** = 83/17) as a colorless solid.

The product mixture (14.2 mg, 78.8 μ mol) was dissolved in 500 μ L DCE and 400 μ L H_2O . Subsequently $RuCl_3 \cdot xH_2O$ (1 small crystal, calc. 0.14 mg, 0.7 μ mol, 3.5 mol% relative to by-product) and $NaIO_4$ (18.6 mg, 87.0 μ mol, 4.3 equiv.) were added and the reaction mixture was stirred at room temperature for 20 h. The reaction was quenched by addition of 2.0 mL H_2O , diluted with 2.0 mL Et_2O and the layers were separated. The aqueous layer was extracted with Et_2O (3 \times 5 mL). The combined organic layers were then washed with H_2O (1 \times 10 mL) and brine (1 \times 10 mL), dried over Na_2SO_4 , filtrated and the solvent was removed under reduced pressure. Purification by flash column chromatography (SiO_2 , P/Et_2O = 2/1, $KMnO_4$) afforded

the product (9.80 mg, 54.4 μmol , 69%) as a colorless solid. Mp 58 – 61 $^{\circ}\text{C}$. TLC (P/Et₂O = 1/2): R_f = 0.16 [KMnO₄]. ¹H NMR (500 MHz, CDCl₃, 300 K): δ = 0.79 (t, ³ J = 7.5 Hz, 3H), 1.21 – 1.33 (m, 1H), 1.37 – 1.51 (m, 1H), 1.63 – 1.78 (m, 1H), 1.93 (*virt.* tdd, ² J \cong ³ J = 14.2 Hz, ³ J = 11.0 Hz, ³ J = 3.8 Hz, 1H), 1.99 – 2.09 (m, 1H), 2.09 – 2.16 (m, 1H), 2.21 (dd, ³ J = 11.0, 6.7 Hz, 1H), 2.31 – 2.43 (m, 2H), 2.71 (d, ³ J = 2.9 Hz, 1H), 2.84 (ddd, ³ J = 8.6, 6.0, 2.9 Hz, 1H), 3.75 (d, ² J = 6.5 Hz, 1H), 3.85 (d, ² J = 6.5 Hz, 1H). ¹³C{¹H} NMR (126 MHz, CDCl₃, 300 K): δ = 11.4 (q), 17.9 (t), 24.4 (t), 27.3 (t), 40.2 (t), 43.4 (d), 52.4 (d), 52.8 (d), 67.8 (t), 88.9 (s), 204.2 (s). IR (ATR): $\tilde{\nu}$ = 2956 (m), 1713 (vs), 1462 (w), 1327 (w), 1095 (m), 1013 (w), 947 (m), 851 (m). MS (EI, 70 eV): m/z (%) = 180 (8), 151 (6), 123 (7), 113 (100), 109 (9), 95 (12), 84 (22). HRMS (EI) m/z : [M]⁺ Calcd. for C₁₀H₁₄O₂: 180.1145; Found: 180.1148.

(3*RS*,3*aRS*,7*aRS*)-3-(Prop-1-en-2-yl)hexahydrobenzofuran-7(4*H*)-one (16a)

According to general procedure 5, compound **16a** was synthesized starting from **3a** (18.4 mg, 102 μmol , 1.0 equiv.) and thioxanthone **5** (2.21 mg, 10.4 μmol , 10 mol%) over 4.5 h. Purification by flash column chromatography (SiO₂, P/Et₂O = 1/2, KMnO₄) afforded the product (11.9 mg, 66.0 μmol , 65%, d.r. > 95/5) as a colorless oil. TLC (P/Et₂O = 1/2): R_f = 0.05 [KMnO₄]. ¹H NMR (500 MHz, CDCl₃, 300 K): δ = 1.46 (*virt.* qd, ² J \cong ³ J = 12.4 Hz, ³ J = 3.8 Hz, 1H), 1.64 (*virt.* qdd, ² J \cong ³ J = 12.6 Hz, ³ J = 5.4, 3.8 Hz, 1H), 1.81 (s, 3H), 1.99 – 2.06 (m, 1H), 2.08 – 2.17 (m, 2H), 2.26 (*virt.* tdd, ² J \cong ³ J = 14.0 Hz, ³ J = 5.4 Hz, ⁴ J = 1.7 Hz, 1H), 2.36 (*virt.* ddt, ² J = 14.1 Hz, ³ J = 5.1 Hz, ³ J \cong ⁴ J = 1.8 Hz, 1H), 2.80 – 2.93 (m, 1H), 4.00 (dd, ² J = 9.1 Hz, ³ J = 1.7 Hz, 1H), 4.10 (dd, ² J = 9.1 Hz, ³ J = 6.3 Hz, 1H), 4.15 (dd, ³ J = 12.4 Hz, ⁴ J = 1.7 Hz, 1H), 4.79–4.81 (m, 1H), 5.03 (*virt.* quint., ² J \cong ⁴ J = 1.3 Hz, 1H). ¹³C{¹H} NMR (75 MHz, CDCl₃, 300 K): δ = 24.7 (q), 25.9 (t), 27.3 (t), 39.8 (t), 47.5 (d), 50.7 (d), 73.7 (t), 84.0 (d), 112.7 (t), 144.3 (s), 207.8 (s). IR (ATR): $\tilde{\nu}$ = 2929 (m), 1730 (vs), 1449 (w), 1105 (m), 1045 (m), 900 (m). MS (EI, 70 eV): m/z

(%) = 180 (73), 165 (8), 136 (21), 123 (17), 107 (100), 93 (46), 81 (29). HRMS (EI) m/z : $[M]^+$ Calcd. for $C_{11}H_{16}O_2$: 180.1145; Found: 180.1145, $[M]^+$ Calcd. for $C_{10}^{13}C_1H_{16}O_2$: 181.1178; Found: 181.1183.

(3*SR*,3*aRS*,7*aRS*)-3*a*-Methyl-3-(prop-1-en-2-yl)hexahydrobenzofuran-7(4*H*)-one (16*b*)

According to general procedure 5, compound **16b** was synthesized starting from **3b** (19.6 mg, 101 μ mol, 1.0 equiv.) and thioxanthone **5** (2.16 mg, 10.2 μ mol, 10 mol%) over 5 h. Purification by flash column chromatography (SiO_2 , P/Et₂O = 2/1, KMnO₄) afforded the product (15.4 mg, 79.1 μ mol, 79%, d.r. > 95/5) as a colorless solid. Mp 87 – 92 °C. TLC (P/Et₂O = 1/2): R_f = 0.08 [KMnO₄]. ¹H NMR (400 MHz, CDCl₃, 300 K): δ = 0.97 (s, 3H), 1.66 (*virt.* td, $^2J \cong ^3J$ = 13.0 Hz, 3J = 5.4 Hz, 1H), 1.73 – 1.79 (m, 1H), 1.81 (s, 3H), 1.87 – 1.98 (m, 1H), 1.98 – 2.06 (m, 1H), 2.20 – 2.34 (m, 2H), 2.52 (ddd, 3J = 7.0, 2.1 Hz, 4J = 0.9 Hz, 1H), 3.95 (dd, 2J = 9.6 Hz, 3J = 2.1 Hz, 1H), 4.19 – 4.26 (m, 2H), 4.79–4.81 (m, 1H), 5.07 (*virt.* quint., $^2J \cong ^4J$ = 1.4 Hz, 1H). ¹³C{¹H} NMR (126 MHz, CDCl₃, 300 K): δ = 20.2 (q), 23.4 (t), 25.5 (q), 30.8 (t), 38.8 (t), 50.6 (s), 55.3 (d), 72.6 (t), 85.3 (d), 112.3 (t), 145.5 (s), 207.8 (s). IR (ATR): $\tilde{\nu}$ = 2924 (s), 1732 (vs), 1455 (w), 1377 (w), 1260 (w), 1104 (w), 1054 (m), 907 (w). MS (EI, 70 eV): m/z (%) = 176 (6), 149 (7), 126 (97), 111 (100), 97 (67), 93 (45). HRMS (EI) m/z : $[M]^+$ Calcd. for $C_{12}H_{18}O_2$: 194.1301; Found: 194.1299.

(3*RS*,3*aRS*,7*aRS*)-5,5-Dimethyl-3-(prop-1-en-2-yl)hexahydrobenzofuran-7(4*H*)-one (16*c*)

According to general procedure 5, compound **16c** was synthesized starting from **3c** (24.8 mg, 120 μ mol, 1.0 equiv.) and thioxanthone **5** (2.57 mg, 12.1 μ mol, 10 mol%) over 1.5 h. Purification by flash column chromatography (SiO_2 , P/Et₂O = 3/1, KMnO₄) afforded the product (21.2 mg, 102 μ mol, 86%, d.r. > 95/5) as a colorless solid. Mp 95 – 101 °C. TLC (P/Et₂O = 3/1): R_f = 0.09

[KMnO₄]. ¹H NMR (500 MHz, CDCl₃, 300 K): δ = 0.99 (s, 3H), 1.13 (s, 3H), 1.46 (*virt.* t, ²J ≡ ³J = 13.0 Hz, 1H), 1.70 (ddd, ²J = 13.0 Hz, ³J = 3.2 Hz, ⁴J = 2.3 Hz, 1H), 1.81 (dd, ⁴J = 1.4, 0.8 Hz, 3H), 2.06 (dd, ²J = 13.5 Hz, ⁴J = 2.3 Hz, 1H), 2.25 (ddd, ²J = 13.5 Hz, ⁴J = 1.7, 0.9 Hz), 2.36 (*virt.* tdd, ³J ≡ ³J = 12.7 Hz, ³J = 6.8, 3.2 Hz, 1H), 2.84 (*virt.* t, ³J ≡ ³J = 6.6 Hz, 1H), 4.03 (dd, ²J = 9.2 Hz, ³J = 1.6 Hz, 1H), 4.10 – 4.21 (m, 2H), 4.79–4.81 (m, 1H), 5.06 (*virt.* quint., ²J ≡ ⁴J = 1.3 Hz, 1H). ¹³C{¹H} NMR (126 MHz, CDCl₃, 300 K): δ = 24.9 (q), 27.2 (q), 32.1 (q), 38.5 (s), 39.0 (t), 46.3 (d), 47.4 (d), 52.9 (t), 74.4 (t), 83.7 (d), 112.7 (t), 144.4 (s), 207.2 (s). IR (ATR): $\tilde{\nu}$ = 2957 (m), 1733 (vs), 1459 (w), 1370 (w), 1126 (w), 1060 (m), 919 (w). MS (EI, 70 eV): *m/z* (%) = 208 (100), 193 (8), 164 (28), 152 (6), 135 (44), 123 (87), 105 (26), 95 (56), 81 (10). HRMS (EI) *m/z*: [M]⁺ Calcd. for C₁₃H₂₀O₂: 208.1458; Found: 208.1458.

(3a*SR*,3b*SR*,6a*RS*,6b*RS*)-6a,6b-Bis(allyloxy)-3a,3b-dimethyloctahydrocyclo-buta[1,2:3,4]-di[5]annulene-1,6-dione (17)

According to general procedure 5, compound **17** was synthesized starting from **1f** (61.7 mg, 405 μmol, 1.0 equiv.) and thioxanthone **5** (8.50 mg, 40.0 μmol, 10 mol%) over 14 h. Purification by flash column chromatography (SiO₂, P/Et₂O = 3/1, KMnO₄) afforded the product (7.10 mg, 23.3 μmol, 12%) as a colorless solid. Mp > 220 °C. TLC (P/Et₂O = 3/1): *R_f* = 0.47 [KMnO₄]. ¹H NMR (400 MHz, CDCl₃, 300 K): δ = 1.25 (s, 6H), 1.70 (ddd, ²J = 14.9 Hz, ³J = 13.1, 8.8 Hz, 2H), 2.24 – 2.41 (m, 4H), 2.54 – 2.73 (m, 2H), 4.08 (*virt.* ddt, ²J = 12.8 Hz, ³J = 5.4 Hz, ⁴J ≡ ⁴J = 1.5 Hz, 2H), 4.18 (*virt.* ddt, ²J = 12.8 Hz, ³J = 5.0 Hz, ⁴J ≡ ⁴J = 1.6 Hz, 2H), 5.07 (*virt.* dq, ³J = 10.5 Hz, ²J ≡ ⁴J = 1.6 Hz, 2H), 5.18 (*virt.* dq, ³J = 17.2 Hz, ²J ≡ ⁴J = 1.6 Hz, 2H), 5.71 – 5.89 (m, 2H). ¹³C{¹H} NMR (101 MHz, CDCl₃, 300 K): δ = 19.1 (q), 32.7 (t), 37.2 (t), 44.2 (s), 68.4 (t), 86.1 (s), 116.1 (t), 134.8 (d), 214.2 (s). IR (ATR): $\tilde{\nu}$ = 2964 (w), 1748 (vs), 1451 (w), 1413 (w), 1186 (w), 1089 (m), 1060 (w), 988 (w), 923 (w). MS (EI, 70 eV): *m/z* (%) = 152 (66),

137 (100), 123 (31), 96 (25), 81 (19), 69 (31), 57 (34), 41 (95). HRMS (EI) m/z : $[M]^+$ Calcd. for $C_{18}H_{24}O_4$: 304.1669; Found: 304.1660.

(3aRS,4aSR,6aSR,9aSR,10aRS,12aRS)-3a,9a-Dimethyldodecahydro-1H,7H-

cyclopenta[1,4]cyclobuta[1,2-*b*]cyclopenta[1,4]cyclobuta[1,2-*f*][1,5]dioxocine-1,7-dione (18)

According to general procedure 5, compound **18** was synthesized starting from **1f** (61.7 mg, 405 μ mol, 1.0 equiv.) and thioxanthone **5** (8.50 mg, 40.0 μ mol, 10 mol%) over 14 h. Purification by flash column chromatography (SiO_2 , P/Et₂O = 3/1, KMnO₄) afforded the product (11.1 mg, 36.5 μ mol, 18%) as a colorless solid. Mp >220 °C. TLC (P/Et₂O = 3/1): R_f = 0.30 [KMnO₄]. ¹H NMR (400 MHz, CDCl₃, 300 K): δ = 1.15 (s, 6H), 1.31 (dd, ² J = 12.9 Hz, ³ J = 4.6 Hz, 2H), 1.58 – 1.70 (m, 2H), 1.72 – 1.89 (m, 4H), 2.27 – 2.47 (m, 4H), 2.67 (ddd, ² J = 17.8 Hz, ³ J = 13.1, 8.8 Hz, 2H), 3.74 (dd, ² J = 12.8 Hz, ³ J = 3.0 Hz, 2H), 4.46 – 4.53 (m, 2H). ¹³C{¹H} NMR (126 MHz, CDCl₃, 300 K): δ = 22.7 (q), 28.9 (t), 34.0 (t), 35.8 (t), 36.0 (d), 44.5 (s), 66.8 (t), 81.3 (s), 220.2 (s). IR (ATR): $\tilde{\nu}$ = 2927 (m), 1735 (vs), 1456 (w), 1266 (w), 1114 (w), 1068 (s), 802 (w). MS (EI, 70 eV): m/z (%) = 304 (33), 263 (25), 248 (53), 230 (20), 208 (49), 191 (15), 175 (10), 161 (13), 152 (72), 137 (100), 113 (54), 97 (74), 84 (67), 67 (63), 55 (85), 41 (95). HRMS (EI) m/z : $[M]^+$ Calcd. for $C_{18}H_{24}O_4$: 304.1669; Found: 304.1674, $[M]^+$ Calcd. for $C_{17}^{13}C_1H_{24}O_4$: 305.1703; Found: 305.1723.

(3aSR,3bSR,6aRS,6bRS)-3a,3b-Dimethyl-6a,6b-bis((2-methylallyl)oxy)octahydrocyclobuta[1,2:3,4]di[5]annulene-1,6-dione (19)

According to general procedure 5, compound **19** was synthesized starting from **2f** (66.9 mg, 402 μ mol, 1.0 equiv.) and thioxanthone **5** (8.50 mg, 40.0 μ mol, 10 mol%) over 3 h. Purification by flash column chromatography (SiO_2 , P/Et₂O = 9/1 \rightarrow 3/1 \rightarrow 1/1, KMnO₄) afforded the

product (11.3 mg, 34.0 μ mol, 17%) as a colorless solid. Mp 50 – 65 °C. TLC (P/Et₂O = 3/1): R_f = 0.44 [KMnO₄]. ¹H NMR (500 MHz, CDCl₃, 300 K): δ = 1.26 (s, 6H), 1.66 (s, 6H), 1.69 – 1.78 (m, 2H), 2.21 – 2.42 (m, 4H), 2.63 (ddd, ² J = 17.6 Hz, ³ J = 13.7, 9.4 Hz, 2H), 3.99 (d, ² J = 12.7 Hz, 2H), 4.10 (d, ² J = 12.7 Hz, 2H), 4.78 (*virt.* dt, ⁴ J = 3.0 Hz, ² J \cong ⁴ J = 1.6 Hz, 2H), 4.83 – 4.94 (m, 2H). ¹³C{¹H} NMR (126 MHz, CDCl₃, 300 K): δ = 19.1 (q), 19.6 (q), 32.7 (t), 37.2 (t), 44.4 (s), 70.3 (t), 85.8 (s), 110.5 (t), 142.3 (s), 214.0 (s). IR (ATR): $\tilde{\nu}$ = 2967 (m), 1749 (vs), 1450 (w), 1186 (w), 1097 (m), 1060 (w), 897 (w). MS (EI, 70 eV): m/z (%) = 213 (13), 182 (25), 166 (65), 151 (44), 123 (23), 110 (41), 97 (23), 84 (46), 69 (27), 55 (100). HRMS (EI) m/z : [M]⁺ Calcd. for C₂₀H₂₈O₄: 332.1982; Found: 332.1998.

(3a*RS*,4a*SR*,6a*RS*,9a*SR*,10a*RS*,12a*SR*)-3a,4a,9a,10a-Tetramethyldodecahydro-1*H*,7*H*-cyclopenta[1,4]cyclobuta[1,2-*b*]cyclopenta[1,4]cyclobuta[1,2-*f*][1,5]dioxocine-1,7-dione (20)

According to general procedure 5, compound **20** was synthesized starting from **2f** (66.9 mg, 402 μ mol, 1.0 equiv.) and thioxanthone **5** (8.50 mg, 40.0 μ mol, 10 mol%) over 3 h. Purification by flash column chromatography (SiO₂, P/Et₂O = 9/1 \rightarrow 3/1 \rightarrow 1/1, KMnO₄) afforded the product (17.3 mg, 52.0 μ mol, 26%) as a colorless solid. Mp 25 – 30 °C. TLC (P/Et₂O = 3/1): R_f = 0.12 [KMnO₄]. ¹H NMR (500 MHz, CDCl₃, 300 K): δ = 0.99 (s, 6H), 1.18 (s, 6H), 1.65 (dddd, ² J = 14.1 Hz, ³ J = 9.2, 4.0 Hz, ⁴ J = 0.9 Hz, 2H), 1.84 (d, ² J = 8.3 Hz, 2H), 2.27 (dt, ² J = 14.1 Hz, ³ J = 8.7 Hz, 2H), 2.49 – 2.68 (m, 6H), 3.95 (d, ² J = 6.7 Hz, 2H), 4.11 (dd, ² J = 6.7 Hz, ⁴ J = 1.7 Hz, 2H). ¹³C{¹H} NMR (126 MHz, CDCl₃, 300 K): δ = 11.6 (q), 15.0 (q), 25.0 (t), 39.6 (t), 44.7 (t), 51.1 (s), 52.2 (s), 81.0 (t), 95.4 (s), 205.5 (s). IR (ATR): $\tilde{\nu}$ = 2957 (m), 1742 (vs), 1450 (w), 1043 (w), 976 (w), 902 (w). MS (EI, 70 eV): m/z (%) = 151 (37), 137 (9), 123 (58), 110 (43), 95 (100), 81 (64), 67 (92), 55 (79), 41 (47). HRMS (EI) m/z : [M]⁺ Calcd. for C₂₀H₂₈O₄: 332.1982; Found: 332.1983.

(3a*SR*,3b*SR*,6a*RS*,6b*RS*)-3a,3b-Dimethyl-6a,6b-bis((3-methylbut-2-en-1-yl)oxy)octa-hydrocyclobuta[1,2:3,4]di[5]annulene-1,6-dione (21**)**

According to general procedure 5, compound **21** was synthesized starting from **3f** (72.3 mg, 401 μ mol, 1.0 equiv.) and thioxanthone **5** (8.50 mg, 40.0 μ mol, 10 mol%) over 4 h. Purification by flash column chromatography (SiO₂, P/Et₂O = 9/1 \rightarrow 3/1 \rightarrow 1/1, KMnO₄) afforded the product (24.3 mg, 67.4 μ mol, 34%) as a colorless solid. Mp 85 – 98 °C. TLC (P/Et₂O = 3/1): R_f = 0.44 [KMnO₄]. ¹H NMR (500 MHz, CDCl₃, 300 K): δ = 1.24 (s, 6H), 1.58 (s, 6H), 1.64 – 1.72 (m, 8H), 2.23 – 2.39 (m, 4H), 2.62 (ddd, ² J = 17.6 Hz, ³ J = 13.8, 9.3 Hz, 2H), 4.05 (dd, ² J = 11.6 Hz, ³ J = 6.9 Hz, 2H), 4.14 (dd, ² J = 11.6 Hz, ³ J = 6.5 Hz, 2H), 5.08 – 5.26 (m, 2H). ¹³C{¹H} NMR (75 MHz, CDCl₃, 300 K): δ = 18.3 (q), 19.3 (q), 25.9 (q), 32.8 (t), 37.1 (t), 44.0 (s), 64.7 (t), 86.3 (s), 121.6 (d), 135.7 (s), 214.4 (s). IR (ATR): $\tilde{\nu}$ = 2967 (m), 1748 (vs), 1448 (w), 1381 (w), 1185 (w), 1090 (m), 1060 (w), 982 (w). MS (EI, 70 eV): m/z (%) = 224 (13), 160 (14), 124 (11), 113 (43), 97 (21), 83 (29), 69 (100), 55 (39), 41 (64). HRMS (EI) m/z : [M]⁺ Calcd. for C₂₂H₃₂O₄: 360.2295; Found: 360.2296.

ASSOCIATED CONTENT

¹H and ¹³C{¹H} NMR spectra for all compounds, configuration assignment by NOESY studies, luminescence spectrum of compound **1a**, emission spectrum of the light source, UV/Vis spectra of all photochemical substrates, X-ray crystallographic data of compounds **18** and **21**. FAIR Data is available as Supporting Information for Publication and includes the primary NMR FID files for all compounds mentioned in the Experimental Section.

ACKNOWLEDGEMENT

Financial support by the European Research Council under the European Union's Horizon 2020 research and innovation programme (grant agreement No 665951 – ELICOS) is gratefully acknowledged. We thank Thomas Pickl for his help in the synthesis of starting materials and in the structure elucidation of a by-product.

REFERENCES

- ¹ General reviews: (a) Poplata, S.; Tröster, A.; Zou, Y.-Q.; Bach, T. Recent Advances in the Synthesis of Cyclobutanes by Olefin [2 + 2] Photocycloaddition Reactions. *Chem. Rev.* **2016**, *116*, 9748–9815. (b) Hoffmann, N. Photochemical Reactions as Key Steps in Organic Synthesis. *Chem. Rev.* **2008**, *108*, 1052-1103. (c) Fleming, S. A. Photocycloaddition of Alkenes to Excited Alkenes. In *Synthetic Organic Photochemistry, Molecular and Supramolecular Photochemistry, Vol. 12*; Griesbeck, A.G.; Mattay, J., Eds.; Dekker: New York, 2005; pp 141-160.
- ² Reviews with a focus on cyclic enones: (a) Hehn, J. P.; Müller, C.; Bach, T. Formation of a Four-Membered Ring: From a Carbonyl-Conjugated Alkene. In *Handbook of Synthetic Photochemistry*; Albini, A.; Fagnoni, M., Eds.; Wiley-VCH: Weinheim, 2010; pp 171-215. (b) Margaretha, P. Photocycloaddition of Cycloalk-2-enones to Alkenes. In *Synthetic Organic Photochemistry, Molecular and Supramolecular Photochemistry, Vol. 12*, Griesbeck, A. G.; Mattay, J. Eds.; Dekker: New York, 2005; pp 211–237. (c) Pete, J.-P. Asymmetric Photoreactions of Conjugated Enones and Esters. *Adv. Photochem.* **1996**, *21*, 135-216. (d) Crimmins, M. T.; Reinhold, T. L. Enone Olefin [2+2] Photochemical Cycloadditions. *Org. React.* **1993**, *44*, 297–588. (e) Baldwin, S. W. Synthetic Aspects of [2+2] Cycloadditions of α,β -Unsaturated Carbonyl Compounds. *Org. Photochem.* **1981**, *5*, 123-225.

³ Reviews with a focus on intramolecular reactions: (a) Crimmins, M. T. Synthetic Applications of Intramolecular Enone-Olefin Photocycloadditions. *Chem. Rev.* **1988**, *88*, 1453-1473. (b) Becker, D.; Haddad, N. Applications of Intramolecular (2+2)Photocycloadditions in Organic Synthesis. *Org. Photochem.* **1989**, *10*, 1-162.

⁴ (a) Schuster, D. I.; Lem, G.; Kaprinidis, N. A. New Insights into an Old Mechanism: [2 + 2] Photocycloaddition of Enones to Alkenes. *Chem. Rev.* **1993**, *93*, 3-22. (b) Schuster, D. I. Mechanistic Issues in [2+2]-Photocycloadditions of Cyclic Enones to Alkenes. In *CRC Handbook of Photochemistry and Photobiology*, 2nd ed.; Horspool, W. M., Lenci, F., Eds.; CRC Press: Boca Raton, 2004; pp 72/1-72/24.

⁵ (a) Srinivasan, R.; Carlough, K. H. Mercury (³P₁) Photosensitized Internal Cycloaddition Reactions in 1,4-, 1,5-, and 1,6-Dienes. *J. Am. Chem. Soc.* **1967**, *89*, 4932-4936. (b) Liu, R. S. H.; Hammond, G. S. Photosensitized Internal Addition of Dienes to Olefins. *J. Am. Chem. Soc.* **1967**, *89*, 4936-4944. (c) Maradyn, D. J.; Weedon, A. C. Trapping of Triplet 1,4-Biradicals with Hydrogen Selenide in the Intramolecular Photochemical Cycloaddition Reaction of 3-(4'-Pentenyl)cycloalk-2-enones: Verification of the Rule of Five. *J. Am. Chem. Soc.* **1995**, *117*, 5359-5360.

⁶ (a) Ikeda, M.; Takahashi, M.; Ohno, K.; Tamura, Y.; Kido, M. Intramolecular photo[2+2]cycloaddition of 2-allyloxy- and 2-allylaminocyclohex-2-enones: Formation of 2-oxa- and 2-aza-bicyclo[2.1.1]hexane systems. *Chem. Pharm. Bull.* **1982**, *30*, 2269-2271. (b) Ikeda, M.; Takahashi, M.; Uchino, T.; Ohno, K.; Tamura, Y.; Kido, M. Intramolecular [2 + 2] Photocycloaddition of 2-(Alkenyloxy)cyclohex-2-enones. *J. Org. Chem.* **1983**, *48*, 4241-4247.

⁷ For some early examples, see: (a) Lu, Z.; Yoon, T. P. Visible Light Photocatalysis of [2+2] Styrene Cycloadditions by Energy Transfer. *Angew. Chem. Int. Ed.* **2012**, *51*, 10329-10332. (b) Zou, Y.-Q.; Duan, S.-W.; Meng, X.-G.; Hu, X.-Q.; Gao, S.; Chen, J.-R.; Xiao, W.-J. Visible light

induced intermolecular [2+2]-cycloaddition reactions of 3-ylideneoxindoles through energy transfer pathway. *Tetrahedron* **2012**, *68*, 6914-6919. (c) Hurtley, A. E.; Lu, Z.; Yoon, T. P. [2+2] Cycloaddition of 1,3-Dienes by Visible Light Photocatalysis. *Angew. Chem., Int. Ed.* **2014**, *53*, 8991-8994. (d) Miller, Z. D.; Lee, B. J.; Yoon, T. P. Enantioselective Crossed Photocycloadditions of Styrenic Olefins by Lewis Acid Catalyzed Triplet Sensitization. *Angew. Chem., Int. Ed.* **2017**, *56*, 11891-11895. (e) Pagire, S. K.; Hossain, A.; Traub, L.; Kerres, S.; Reiser, O. Photosensitized regioselective [2+2]-cycloaddition of cinnamates and related alkenes. *Chem. Comm.* **2017**, *53*, 12072-12075.

⁸ (a) Clay, A.; Vallavoju, N.; Krishnan, R.; Ugrinov, A.; Sivaguru, J. Metal-Free Visible Light-Mediated Photocatalysis: Controlling Intramolecular [2 + 2] Photocycloaddition of Enones through Axial Chirality. *J. Org. Chem.* **2016**, *81*, 7191-7200. (b) Elliott, L. D.; Berry, M.; Harji, B.; Klauber, D.; Leonard, J.; Booker-Milburn, K. I. A Small-Footprint, High-Capacity Flow Reactor for UV Photochemical Synthesis on the Kilogram Scale. *Org. Process Res. Dev.* **2016**, *20*, 1806-1811. (c) Zhao, J.; Brosmer, J. L.; Tang, Q.; Yang, Z.; Houk, K. N.; Diaconescu, P. L.; Kwon, O. Intramolecular Crossed [2+2] Photocycloaddition through Visible Light-Induced Energy Transfer. *J. Am. Chem. Soc.* **2017**, *139*, 9807-9810. (d) Williams, J. D.; Nakano, M.; Gérardy, R.; Rincón, J. A.; de Frutos, Ó.; Mateos, C.; Monbaliu, J.-C. M.; Kappe, C. O. Finding the Perfect Match: A Combined Computational and Experimental Study toward Efficient and Scalable Photosensitized [2 + 2] Cycloadditions in Flow. *Org. Process Res. Dev.* **2019**, *23*, 78-87. (e) Strieth-Kalthoff, F.; Henkel, C.; Teders, M.; Kahnt, A.; Knolle, W.; Gómez-Suárez, A.; Dirian, K.; Alex, W.; Bergander, K.; Daniliuc, C. G.; Abel, B.; Guldi, D. M.; Glorius, F. Discovery of Unforeseen Energy-Transfer-Based Transformations Using a Combined Screening Approach. *Chem* **2019**, *5*, 2183-2194. (f) Pecho, F.; Zou, Y.-Q.; Gramüller, J.; Mori, T.; Huber, S. M.; Bauer, A.; Gschwind, R. M.; Bach, T. A Thioxanthone Sensitizer with a Chiral Phosphoric

1
2
3 Acid Binding Site: Properties and Applications in Visible Light-Mediated Cycloadditions. *Chem.*
4
5 *Eur.* **2020**, 26, 5190-5194.
6

7
8 ⁹ For some early examples for hydrogen abstraction reactions, see: (a) Corey, E. J.; Bass, J. D.;
9
10 LeMahieu, R.; Mitra, R. B. A Study of the Photochemical Reactions of 2-Cyclohexenones with
11
12 Substituted Olefins. *J. Am. Chem. Soc.* **1964**, 86, 5570-5583. (b) Hunter, N. R.; MacAlpine, G.
13
14 A.; Liu, H. J.; Valenta, Z. Cyclobutanes in organic synthesis. Part I. Fragmentation to 1,4-
15
16 dicarbonyl compounds. General considerations. *Can. J. Chem.* **1970**, 48, 1436-1445. (c)
17
18 Greenlee, M. L.; Fritzen Jr. E. L.; Swenton, J. S. A Unique Reversal of Regioselectivity in the
19
20 Photoaddition of 2-Fluorocyclohexenone to Isobutylene. *J. Org. Chem.* **1978**, 43, 4512-4515. (d)
21
22 Shih, C.; Fritzen, E. L.; Swenton, J. S. Photocycloaddition Chemistry of 2-
23
24 (Trimethylsilyl)cyclopentenone and 5-(Trimethylsilyl)uracil. The Utility of a Trimethylsilyl
25
26 Group as a Removable Directing Group in Photochemistry. *J. Org. Chem.* **1980**, 45, 4462-4471.
27
28 (e) Pirrung, M. C.; Webster, N. J. G. Cyclooctenone photocycloadditions. *Tetrahedron Lett.*
29
30 **1986**, 27, 3983-3986. (f) Pirrung, M. C.; Webster, N. J. G. Mechanism of Intramolecular
31
32 Photocycloadditions of Cyclooctenones. *J. Org. Chem.* **1987**, 52, 3603-3613.
33
34
35
36

37
38 ¹⁰ Rodríguez, J. R.; Castedo, L.; Mascareñas, J. L. Synthesis of Eight- and Nine-Membered
39
40 Carbocycles through a Ring-Closing Metathesis/Ring Fragmentation Strategy: A Rapid and
41
42 Versatile Approach to Bicyclo[6.4.0]- and Bicyclo[7.4.0]alkene Ring Systems. *Chem. Eur. J.*
43
44 **2002**, 8, 2923-2930.
45

46
47 ¹¹ (a) Dauben, W. G.; Ponaras, A. A.; Chollet, A. An Approach to Angularly Functionalized
48
49 Methylhydrindan Systems. *J. Org. Chem.* **1980**, 45, 4413-4417. (b) Schick, H.; Schwarz, H.;
50
51 Finger, A.; Schwarz, S. 2,2-Disubstituierte Cyclopentan-1,3-dione – 2: Untersuchung der
52
53 Regioselektivität von Alkylierungsreaktionen an 2-Methyl-cyclopentan-1,3-dion. *Tetrahedron*
54
55
56
57
58
59
60

1982, 38, 1279-1283. (c) Ponaras, A. A. Axial allylation on the diosphenol claisen rearrangement. *Tetrahedron Lett.* **1983**, 24, 3-6.

¹² (a) Ponaras, A. A. The diosphenol Claisen rearrangement. *Tetrahedron Lett.* **1980**, 21, 4803-4806. (b) Matoba, K.; Karibe, N.; Yamazaki, T. Chemistry of 2-Methoxy-2, 5-cyclohexadienones. I. Photochemistry of 2-Methoxy-4, 4-dimethyl-2, 5-cyclohexadienone. *Chem. Pharm. Bull.* **1982**, 30, 3906-3911. (c) Münster, N.; Parker, N. A.; van Dijk, L.; Paton, R. S.; Smith, M. D. Visible Light Photocatalysis of 6 π Heterocyclization. *Angew. Chem., Int. Ed.* **2017**, 56, 9468-9472. (d) Edtmüller, V.; Pöthig, A.; Bach, T. Enantioselective photocyclisation reactions of 2-aryloxy-cyclohex-2-enones mediated by a chiral copper-bisoxazoline complex. *Tetrahedron* **2017**, 73, 5038-5047.

¹³ Li, X.; Jandl, C.; Bach, T. Visible-Light-Mediated Enantioselective Photoreactions of 3-Alkylquinolones with 4-*O*-Tethered Alkenes and Allenes. *Org. Lett.* **2020**, 22, 3618-3622.

¹⁴ Iyer, A.; Clay, A.; Jockusch, S.; Sivaguru, J. Evaluating brominated thioxanones as organophotocatalysts. *J. Phys. Org. Chem.* **2017**, 30, e3738.

¹⁵ Singh, A.; Teegardin, K.; Kelly, M.; Prasad, K. S.; Krishnan, S.; Weaver, J. D. Facile Synthesis and Complete Characterization of Homoleptic and Heteroleptic Cyclometalated Iridium(III) Complexes for Photocatalysis. *J. Organomet. Chem.* **2015**, 776, 51-59.

¹⁶ (a) Juris, A.; Balzani, V.; Belser, P.; von Zelewsky, A. Characterization of the Excited State Properties of Some New Photosensitizers of the Ruthenium (Polypyridine) Family. *Helv. Chim. Acta* **1981**, 64, 2175-2182. (b) Kalyanasundaram, K. Photophysics, photochemistry and solar energy conversion with tris(bipyridyl)ruthenium(II) and its analogues. *Coord. Chem. Rev.* **1982**, 46, 159-244.

¹⁷ (a) Angulo, G.; Grilj, J.; Vauthey, E.; Serra-Andrés, L.; Rubio-Pons, Ò.; Jacques, P. Ultrafast Decay of the Excited Singlet States of Thioxanthone by Internal Conversion and Intersystem Crossing. *ChemPhysChem* **2010**, *11*, 480–488. (b) Mundt, R.; Villnow, T.; Ziegenbein, C. T.; Gilch, P.; Marian, C.; Rai-Constapel, V. Thioxanthone in apolar solvents: ultrafast internal conversion precedes fast intersystem crossing. *Phys. Chem. Chem. Phys.* **2016**, *18*, 6637–6647.

¹⁸ For recent reviews on sensitized photochemical reactions, see: (a) Zhou, Q.-Q.; Zou, Y.-Q.; Lu, L.-Q.; Xiao, W.-J. Visible-Light-Induced Organic Photochemical Reactions through Energy-Transfer Pathways. *Angew. Chem. Int. Ed.* **2019**, *58*, 1586-1604. (b) Strieth-Kalthoff, F.; James, M. J.; Teders, M.; Pitzer, L.; Glorius, F. Energy Transfer Catalysis Mediated by Visible Light: Principles, Applications, Directions. *Chem. Soc. Rev.* **2018**, *47*, 7190-7202.

¹⁹ (a) House, H. O.; Huber, L. E.; Umen, M. J. Empirical Rules for Estimating the Reduction Potential of α,β -Unsaturated Carbonyl Compounds. *J. Am. Chem. Soc.* **1972**, *94*, 8471-8475. (b) Mandell, L.; Heldrich, F. J.; Day, Jr., R. A. The Electrochemical Reductive Coupling of Mono-Enol Ethers of β -Diketones. *Synth. Commun.* **1981**, *11*, 55-59. (c) Harada, J.; Sakakibara, Y.; Kunai, A.; Sasaki, K. Electrochemical Carboxylation of α,β -Unsaturated Ketones with Carbon Dioxide. *Bull. Chem. Soc. Jpn.* **1984**, *57*, 611-612.

²⁰ (a) Dilling, W. L.; Tabor, T. E.; Boer, F. P.; North, P. P. Organic Photochemistry. IX. The Photocycloaddition of 2-Cyclopentenone to *cis*- and *trans*-Dichloroethylene. Evidence for Initial Attack at Carbon-3 and Rotational Equilibration of the Diradical Intermediates. *J. Am. Chem. Soc.* **1970**, *92*, 1399-1400. (b) Peet, N. P.; Cargill, R. L.; Bushey, D. F. Synthesis and Acid-Catalyzed Rearrangements of Tricyclo[4.3.2.0]undecanones. *J. Org. Chem.* **1973**, *38*, 1218-1221. (c) Becker, D.; Nagler, M.; Hirsh, S.; Ramun, J. Intramolecular [2 + 2] Photocycloadditions of *E* and *Z* Olefins to Cyclohex-2-enone. *J. Chem. Soc., Chem. Commun.* **1983**, 371-373. (d) Becker, D.; Nagler, M.; Sahali, Y.; Haddad, N. Regiochemistry and Stereochemistry of Intramolecular

- [2 + 2] Photocycloaddition of Carbon-Carbon Double Bonds to Cyclohexenones. *J. Org. Chem.* **1991**, *56*, 4537-4543. (e) Kelly, J. F. D.; Kelly, J. M.; McMurry, T. B. H. Photochemistry of substituted cyclic enones. Part 12.¹ Photocycloaddition of 3-phenylcyclopentenone and 3-phenylcyclohexenone to (*E*)- and (*Z*)-1-phenylpropene. *J. Chem. Soc., Perkin Trans. 2* **1999**, 1933-1941. (f) Brimioulle, R.; Bauer, A.; Bach, T. Enantioselective Lewis Acid Catalysis in Intramolecular [2 + 2] Photocycloaddition Reactions: A Mechanistic Comparison between Representative Coumarin and Enone Substrates. *J. Am. Chem. Soc.* **2015**, *137*, 5170-5176. (g) Poplata, S.; Bauer, A.; Storch, G.; Bach, T. Intramolecular [2+2] Photocycloaddition of Cyclic Enones: Selectivity Control by Lewis Acids and Mechanistic Implications. *Chem. Eur. J.* **2019**, *25*, 8135-8148.
- ²¹ Poplata, S.; Bach, T. Enantioselective Intermolecular [2+2] Photocycloaddition Reaction of Cyclic Enones and Its Application in a Synthesis of (–)-Grandisol. *J. Am. Chem. Soc.* **2018**, *140*, 3228-3231.
- ²² For related structures, see: Piva-Le Blanc, S.; Pete, J.-P.; Piva, O. Conformational effects and cyclodimer formation in intramolecular [2 + 2] photocycloadditions, *Chem. Commun.* **1998**, 235-236.
- ²³ Examples: (a) Eaton, P. E. The Tricyclo[5.3.0.0^{2,6}]decane System. The Photodimers of Cyclopentenone. *J. Am. Chem. Soc.* **1962**, *84*, 2344-2348. (b) Wagner, P. J.; Bucheck, D. J. Comparison of the Photodimerizations of 2-Cyclopentenone and of 2-Cyclohexenone in Acetonitrile. *J. Am. Chem. Soc.* **1969**, *91*, 5090-5097. (c) Reinfried, R.; Belluš, D.; Schaffner, K. Photochemical Reactions. Part 64 [1]. The Photochemistry of 3-Alkyl-cyclopent-2-en-ones. *Helv. Chim. Acta* **1971**, *54*, 1517-1531. (d) Berenjian, N.; De Mayo, P.; Sturgeon, M.-E.; Sydnes, L. K.; Weedon, A. C. Biphasic photochemistry: micelle solutions as media for photochemical cycloadditions of enones. *Can. J. Chem.* **1982**, *60*, 425-436. (e) Yang, J.; Dewal, M. B.;

Profeta, S.; Smith, M. D.; Li, Y.; Shimizu, L. S. Origins of Selectivity for the [2+2] Cycloaddition of α,β -unsaturated Ketones within a Porous Self-assembled Organic Framework. *J. Am. Chem. Soc.* **2008**, *130*, 612-621.

²⁴ A head-to-tail preference is normally expected although the selectivity is not pronounced: (a) Hill, E. A.; Theissen, R. J.; Cannon, C. E.; Miller, R.; Guthrie, R. B.; Chen, A. T. Ring Cleavage Rearrangements of 2-Bicyclo[3.2.0]heptyl and Related Grignard Reagents. *J. Org. Chem.* **1976**, *41*, 1191-1199. (b) Sydnes, L. K. M.; Helge L. Photochemical [2+2] Cycloadditions. II. Photochemical Addition of 3-Butyl-2-cyclopentenone to Some ω -Bromo-1-alkenes. *Acta Chem. Scand. B* **1987**, *41*, 660-667. (c) Rudolph, A.; Weedon, A. C. Radical clocks as probes of 1,4-biradical intermediates in the photochemical cycloaddition reactions of 2-cyclopentenone with alkenes. *Can. J. Chem.* **1990**, *68*, 1590-1597. (d) Hastings, D. J.; Weedon, A. C. Origin of the Regioselectivity in the Photochemical Cycloaddition Reactions of Cyclic Enones with Alkenes: Chemical Trapping Evidence for the Structures, Mechanism or Formation and Fates of 1,4-Biradical Intermediates. *J. Am. Chem. Soc.* **1991**, *113*, 8525-8527. (e) Griesbeck, A. G.; Stadtmüller, S.; Busse, H.; Bringmann, G.; Buddrus, J. Stereoselektivität von Photocycloadditionen, ^{5[1]} Photoreaktionen zwischen 2-Cyclopenten-1-on und Enolethern. *Chem. Ber.* **1992**, *125*, 933-940. (f) Andrew, D.; Hastings, D. J.; Weedon, A. C. The Mechanism of the Photochemical Cycloaddition Reaction between 2-Cyclopentenone and Polar Alkenes: Trapping of Triplet 1,4-Biradical Intermediates with Hydrogen Selenide. *J. Am. Chem. Soc.* **1994**, *116*, 10870-10882. (g) Sydnes, L. K.; Van Ha, D. Cyclobutanes and Cyclobutenes from Photochemical Cleavage of Some Bicyclo[3.2.0]heptan-2-ones. *Aust. J. Chem.* **2009**, *62*, 101-107.

²⁵ Ponaras, A. A.; Meah, M. Y. Synthesis of diosphenol ethers by means of alkoxytrimethylsilanes. *Tetrahedron Lett.* **1986**, *27*, 4953-4956.

- ²⁶ Ponaras, A. A. A New Variant of the Claisen Rearrangement Capable of Creating the Bond between Two Quaternary Centers. *J. Org. Chem.* **1983**, *48*, 3866-3868.
- ²⁷ Bergman, R.; Magnusson, G. Mechanism of Cyclopentene-1-carboxaldehyde Formation by Ring Contraction of 2,3-Epoxycyclohexanols. Improved Synthetic Procedures. *J. Org. Chem.* **1986**, *51*, 212-217.
- ²⁸ Kraft, P.; Jordi, S.; Denizot, N.; Felker, I. On the Dienone Motif of Musks: Synthesis and Olfactory Properties of Partially and Fully Hydrogenated Dienone Musks. *Eur. J. Org. Chem.* **2014**, 554-563.
- ²⁹ Luengo, J. I.; Koreeda, M. Complete Control of the Rearrangement Modes of Enolates of α -Allyloxy Ketones: Reversal from the [3,3]-Claisen to the [2,3]-Wittig Pathway by the Use of the Metalated N,N-Dimethylhydrazones. *J. Org. Chem.* **1989**, *54*, 5415-5417.
- ³⁰ Madelaine, C.; Valerio, V.; Maulide, N. Unexpected Electrophilic Rearrangements of Amides: A Stereoselective Entry to Challenging Substituted Lactones. *Angew. Chem., Int. Ed.* **2010**, *49*, 1583-1586.