




Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/lncn20>

A NOVEL PROTECTING STRATEGY FOR INTERNUCLEOSIDIC PHOSPHATE AND PHOSPHOROTHIOATE GROUPS

Andrei P. Guzaev^a & Muthiah Manoharan^a

^a Department of Medicinal Chemistry, Isis Pharmaceuticals, 2292 Faraday Ave., Carlsbad, California, 92008, U.S.A.

Published online: 07 Feb 2007.

To cite this article: Andrei P. Guzaev & Muthiah Manoharan (2001) A NOVEL PROTECTING STRATEGY FOR INTERNUCLEOSIDIC PHOSPHATE AND PHOSPHOROTHIOATE GROUPS, Nucleosides, Nucleotides and Nucleic Acids, 20:4-7, 1011-1014, DOI: [10.1081/NCN-100002480](https://doi.org/10.1081/NCN-100002480)

To link to this article: <http://dx.doi.org/10.1081/NCN-100002480>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

A NOVEL PROTECTING STRATEGY FOR INTERNUCLEOSIDIC PHOSPHATE AND PHOSPHOROTHIOATE GROUPS

Andrei P. Guzaev* and Muthiah Manoharan

Department of Medicinal Chemistry, Isis Pharmaceuticals,
2292 Faraday Ave., Carlsbad, California 92008

ABSTRACT

The utility of 2-(*N*-isopropyl-*N*-anisoylamino)ethyl group for protection of internucleosidic phosphate linkages in oligonucleotide synthesis was studied. The group demonstrated high coupling yields, favorable deprotection kinetics and a high hydrolytic stability of phosphoramidite building blocks. The mechanism of deprotection was established using a model phosphate triester.

Current oligonucleotide synthesis employs a phosphoramidite method and is carried out automatically by stepwise coupling of nucleoside building blocks to a solid support. This allows an efficient preparation of natural and modified DNA fragments on a routine basis. Phosphoramidite building blocks are most often protected with a 2-cyanoethyl group at the phosphite moiety (1). Treatment of the oligonucleotide with ammonia effects β -elimination in the 2-cyanoethyl group releasing internucleosidic phosphodiester moieties and acrylonitrile as a side product. Recently, acrylonitrile has been demonstrated to alkylate nucleic bases under conditions of deprotection (2). Increasing interest in manufacturing oligonucleotides as drugs motivated our search for alternate protecting groups, whose removal is governed by mechanisms different from β -elimination.

We reported recently the use of 2-[(1-naphthyl)carbamoyloxy]ethyl group for protection of internucleosidic phosphates (3). Here we report the use of a novel 2-(*N*-isopropyl-*N*-anisoylamino)ethyl protection in the oligonucleotide synthesis.

*Corresponding author.




Figure 1. Synthesis of phosphoramidite building blocks 8–12.

For the preparation of novel phosphoramidite building blocks, **1** was converted to a bisamidite **2**, which was isolated as a crystalline compound on silica gel. Nucleosides **3–7** were treated with **2** in the presence of 1*H*-tetrazole to give **8–12** (Fig. 1). These were isolated by column chromatography in 75–89% yield and characterized. The phosphoramidites **8–12** were stable compounds whose shelf life exceeded 9 months. Moreover, as measured by ³¹P NMR, the half-life of disappearance of **8** in 95% aqueous MeCN was 73 h, which indicated its high hydrolytic stability.

Next, the removal of the proposed protecting group (PG) was studied. Model compounds, **13–15**, were synthesized either on a solid support (**13** and **14**) or in solution (**15**, Fig. 2). Half-lives of deprotection of **13** and **14** under various conditions were measured by a gel phase ³¹P NMR. In phosphite triester **13**, the PG was indefinitely stable in MeCN. In contrast, it was rapidly removed from phosphate triester **14** ($\tau_{1/2}$ = 50 min in MeCN). In a similar manner, a model T₁₅ phosphorothioate was assembled on a polystyrene solid support (20 μ mol) using phosphoramidite **8**. As revealed by ³¹P NMR in gel phase, the PG was removed to ca. 90% extent when the synthesis of the oligonucleotide was complete.

Apparently, the extent of deprotection of the oligonucleotide attached to a solid support should depend on the duration of the synthesis and hence on the synthetic




Figure 2a. Structures of model compounds 13–15.






Figure 2b. The fate of **15** under neutral and basic aqueous conditions.

scale employed. On a small scale, the synthesized oligonucleotide remains largely protected. In this case, the remaining protecting groups are removed at the final deprotection step. In order to illustrate the deprotection mechanism under neutral and basic aqueous conditions, the fate of **15** was next studied (Fig. 2). The products, **1** and **19–23**, formed from **15** do not react with nucleic bases. To confirm this, mixtures of **15** (10 equiv) and each of the 5'-DMTr protected nucleosides (dA, dC, dG, and T; 1 equiv) were treated with ammonium hydroxide. As demonstrated by HPLC, each of the nucleosides remain unchanged for at least 48 h at 55°C.

It is known that nucleoside phosphoramidites efficiently and selectively couple at the 5'-position of solid support-bound oligonucleotides with deprotected backbone (4). Phosphoramidites **8–12** were used for preparation of oligodeoxynucleotides and their phosphorothioate and 2'-O-(2-methoxyethyl) analogs, **24–30**, on 1 to 150 μ mol scale (Table). In agreement with previous observations (7), **8–12** were capable of an efficient chain elongation. Analysis of the deprotection mixtures revealed no apparent modifications of nucleic bases or oligonucleotide backbone. The yields of isolated oligonucleotides were similar to those obtained with the standard 2-cyanoethyl phosphoramidites.

Table 1. Oligonucleotides **24–30** Synthesized with the Aid of **8–12**

Compound	Sequence (5' $\rightarrow$ 3')	Backbone	Target
24	GCC CAA GCT GGC ATC CGT CA	P=O	Human ICAM-1
25	TCC CGC CTG TGA CAT GCA TT	P=O	Human C-raf
26	GCC CAA GCT GGC ATC CGT CA	P=S	Human ICAM-1
27	TCC CGC CTG TGA CAT GCA TT	P=S	Human C-raf
28	ATG CAT TCT GCC CCC AAG GA	P=S	Mouse C-raf
29	AGC TTC TTT GCA CAT GTA AA	P=S	Human MDM-2
30	(T*T*T*) ₆ T*T*T*	P=S	—

a. T* stands for 2'-O-(2-methoxyethyl)-5-methyluridine residue.



In conclusion, the 2-(*N*-isopropyl-*N*-anisoylamino)ethyl is a novel, convenient phosphodiester protecting group. It is quantitatively removed either on solid support by treatment with a polar organic solvent or under standard deprotection conditions.

REFERENCES

1. Sinha, N. D.; Biernat, J.; McManus, J.; Köster, H. *Nucleic Acids Res.* **1984**, *12*, 4539–4557.
2. Wilk, A.; Grajkowski, A.; Phillips, L. R.; Beaucage, S. L. *J. Org. Chem.* **1999**, *64*, 7515–7522.
3. Guzaev, A. P.; Manoharan, M. *Tetrahedron Lett.* **2000**, *41*, 5623–5626.
4. Caruthers, M. H.; Kierzek, R.; Tang, J. Y. In *Biophosphates and their Analogues – Synthesis, Metabolism and Activity*; Bruzik, K. S.; Stec, W. J. Eds.; Elsevier: Amsterdam, 1987; pp. 3–21.



Request Permission or Order Reprints Instantly!

Interested in copying and sharing this article? In most cases, U.S. Copyright Law requires that you get permission from the article's rightsholder before using copyrighted content.

All information and materials found in this article, including but not limited to text, trademarks, patents, logos, graphics and images (the "Materials"), are the copyrighted works and other forms of intellectual property of Marcel Dekker, Inc., or its licensors. All rights not expressly granted are reserved.

Get permission to lawfully reproduce and distribute the Materials or order reprints quickly and painlessly. Simply click on the "Request Permission/Reprints Here" link below and follow the instructions. Visit the [U.S. Copyright Office](#) for information on Fair Use limitations of U.S. copyright law. Please refer to The Association of American Publishers' (AAP) website for guidelines on [Fair Use in the Classroom](#).

The Materials are for your personal use only and cannot be reformatted, reposted, resold or distributed by electronic means or otherwise without permission from Marcel Dekker, Inc. Marcel Dekker, Inc. grants you the limited right to display the Materials only on your personal computer or personal wireless device, and to copy and download single copies of such Materials provided that any copyright, trademark or other notice appearing on such Materials is also retained by, displayed, copied or downloaded as part of the Materials and is not removed or obscured, and provided you do not edit, modify, alter or enhance the Materials. Please refer to our [Website User Agreement](#) for more details.

[Order now!](#)

Reprints of this article can also be ordered at

<http://www.dekker.com/servlet/product/DOI/101081NCN100002480>