

Dehydroxylative Fluorination of Tertiary Alcohols

Wei Zhang, Yu-Cheng Gu, Jin-Hong Lin,* and Ji-Chang Xiao*



Cite This: <https://dx.doi.org/10.1021/acs.orglett.0c02438>



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

ABSTRACT: A large number of fluorination methods have been developed, but the construction of a tertiary C–F bond remains challenging. Herein, we describe an efficient dehydroxylative fluorination of tertiary alcohols with Selectfluor via the activation of a hydroxyl group by a $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2/\text{ICH}_2\text{CH}_2\text{I}$ system. Although the reagents appear to be not compatible (Selectfluor with the phosphine and I^- generated in situ), the reactions occur rapidly to give the desired products in moderate to high yields. This work may present a new discovery in fluorination of alcohols since the reported methods are mainly limited to primary and secondary alcohols.



Owing to the unique electronic effects of the fluorine element, the incorporation of a fluorine atom into an organic molecule may profoundly modify its physical, chemical, and biological properties.¹ As tertiary fluorides are important structural motifs in pharmaceuticals such as betamethasone and sofosbuvir, it would be highly desirable to develop efficient fluorination methods for the construction of a tertiary C–F bond. Although many fluorination methods have been developed,² site-selective fluorination for tertiary C–F bond formation remains a challenging task.




Nucleophilic fluorination of alkyl halides is one of the most widely used approaches for fluorine incorporation, but the transformation of tertiary alkyl halides suffers from competing side reactions such as eliminations and rearrangements.^{1c} Nishikata³ and Li⁴ independently found that radical versions could effectively install tertiary C–F bonds (Scheme 1, eq 1). Cesium fluoride and Selectfluor were used as the fluorine sources, respectively. The Nishikata method can only be applied to α -bromo-*N*-arylamides, not to ordinary tertiary alkyl halides. Li's approach works well with ordinary tertiary bromides and iodides, but it is difficult to purify these

substrates because during isolation they would readily undergo an elimination or rearrangement via the formation of tertiary cations. Besides halide groups, other leaving groups derived from hydroxyl functionality can also be substituted by fluorine via a radical reaction or a transition-metal-catalyzed process, but an installation of the leaving groups is required before fluorination (Scheme 1, eq 2).⁵ Decarboxylative fluorination is an effective approach for tertiary C–F bond construction (Scheme 1, eq 3).⁶ Apparently, it is still highly desirable to develop efficient methods for the installation of tertiary C–F bond.


Since alcohols are widely available and hydroxyl group is commonly found in natural products, direct dehydroxylative fluorination of alcohols has received increasing attention. Traditional methods suffer from the use of a hazardous S–F type reagent, such as SF_4 and DAST (diethylaminosulfur trifluoride).⁷ The safety issue has prompted the development of milder dehydroxyfluorination reagents,⁸ including PyFluor by Doyle,⁹ PhenoFluor¹⁰ and AlkylFluor¹¹ by Ritter. Recently, Watson disclosed that fluorination of alcohols with CuF_2 occurred smoothly in the presence of *N,N'*-diisopropylcarbodiimide for the activation of alcohols.¹² All of these dehydroxylative fluorination methods are quite efficient for the conversion of primary and secondary alcohols, but cannot be well applied to tertiary alcohols. We have previously found that the R_3P could react with $\text{ICH}_2\text{CH}_2\text{I}$ to generate iodophosphonium salt $\text{R}_3\text{P}^+\text{I}^-$, which can act as a strong Lewis acid to activate hydroxyl groups,¹³ and thus, the $\text{R}_3\text{P}/\text{ICH}_2\text{CH}_2\text{I}$ system was used to enable nucleophilic fluorination of alcohols.^{13a} This process can also only be extended to

Scheme 1. Tertiary C–F bond formation

Previous work:



This work:



Received: July 21, 2020

primary and secondary alcohols. Herein, we describe the dehydroxylyative fluorination of tertiary alcohols with Selectfluor via the activation of hydroxyl group by a $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2/\text{ICH}_2\text{CH}_2\text{I}$ system (Scheme 1, eq 4). The reactions proceeded rapidly (15 min) at room temperature to give moderate to high yields.

The failure of the $\text{R}_3\text{P}/\text{ICH}_2\text{CH}_2\text{I}$ -promoted nucleophilic fluorination of tertiary alcohols encouraged us to investigate a radical process. We have previously shown $\text{Ph}_3\text{P}/\text{ICH}_2\text{CH}_2\text{I}$ can effectively promote iodination of alcohols,^{13a,b} and it has been reported by Li that tertiary iodides could readily undergo radical fluorination with Selectfluor.⁴ Therefore, it is reasonable to speculate that $\text{Ph}_3\text{P}/\text{ICH}_2\text{CH}_2\text{I}$ -promoted fluorination of tertiary alcohols with Selectfluor might occur if iodination product could be first generated in situ. However, two incompatible pairs of reagents may completely inhibit this fluorination. Ph_3P would readily undergo a redox reaction with Selectfluor.¹⁴ Therefore, Ph_3P may not be able to cooperate with $\text{ICH}_2\text{CH}_2\text{I}$ to activate alcohols, and the consumption of Selectfluor would also suppress fluorination. Furthermore, the iodination requires the presence of iodide anion, which could be produced from the $\text{Ph}_3\text{P}/\text{ICH}_2\text{CH}_2\text{I}$ system.^{13a,b} But the iodide anion may also be easily oxidized by Selectfluor.¹⁵ Apparently, both iodination and fluorination would be suppressed due to the redox reactions of Selectfluor with Ph_3P and I^- anion.

We have found that DMF and CH_3CN are suitable solvents for the $\text{Ph}_3\text{P}/\text{ICH}_2\text{CH}_2\text{I}$ -promoted dehydroxylation of alcohols,¹³ and thus, these solvents were first screened for the fluorination of substrate **1a** (Table 1). No desired product was observed in DMF (entry 1). Although a low yield was obtained in CH_3CN (entry 2), the generation of the desired product

was reflective of the feasibility of the proposed fluorination process. Besides Ph_3P , other trivalent phosphines were also examined (entries 3–9), and $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2$ was found to be a suitable choice (entry 6). As both tertiary iodides and bromides can be fluorinated,⁴ we thought that adding external iodide or bromide anion might be favorable for iodination or bromination and may thus further increase the fluorination yield. A brief survey of the ion additives revealed that using ZnBr_2 significantly increased the yield to 82% (entry 13). A 99% yield was obtained by increasing the loading of ZnBr_2 (entry 15). The reaction occurred rapidly and was completed with 15 min (entry 17).

With the optimal reaction conditions in hand (Table 1, entry 17), we then investigated the substrate scope of the dehydroxylyative fluorination of tertiary alcohols. As shown in Scheme 2, the fluorination process could be extended to a wide

Scheme 2. Dehydroxylyative Fluorination of Tertiary Alcohols*




Table 1. Optimization of Reaction Conditions^a

entry	R_3P	equiv ratio ^a	additive	yield ^b (%)
1 ^c	Ph_3P	1.2:1.2:0		ND
2	Ph_3P	1.2:1.2:0		26
3	$(\text{C}_6\text{F}_5)_3\text{P}$	1.2:1.2:0		9
4	$\text{Ph}_2\text{PCH}=\text{CH}_2$	1.2:1.2:0		16
5	$\text{Ph}_2\text{PCH}_2\text{PPh}_2$	0.6:1.2:0		21
6	$\text{Ph}_2\text{P}(\text{CH}_2)_2\text{PPh}_2$	0.6:1.2:0		65
7	$\text{Ph}_2\text{P}(\text{CH}_2)_3\text{PPh}_2$	0.6:1.2:0		18
8	BINAP	0.6:1.2:0		31
9	Xantphos	0.6:1.2:0		6
10	$\text{Ph}_2\text{P}(\text{CH}_2)_2\text{PPh}_2$	0.6:1.2:0.1	$^t\text{Bu}_4\text{N}^+\text{I}^-$	58
11	$\text{Ph}_2\text{P}(\text{CH}_2)_2\text{PPh}_2$	0.6:1.2:0.1	$^t\text{Bu}_4\text{N}^+\text{Br}^-$	66
12	$\text{Ph}_2\text{P}(\text{CH}_2)_2\text{PPh}_2$	0.6:1.2:0.1	KBr	68
13	$\text{Ph}_2\text{P}(\text{CH}_2)_2\text{PPh}_2$	0.6:1.2:0.1	ZnBr_2	82
14	$\text{Ph}_2\text{P}(\text{CH}_2)_2\text{PPh}_2$	0.6:1.2:0.6	ZnBr_2	89
15	$\text{Ph}_2\text{P}(\text{CH}_2)_2\text{PPh}_2$	0.6:1.2:0.8	ZnBr_2	99
16	$\text{Ph}_2\text{P}(\text{CH}_2)_2\text{PPh}_2$	0.6:1.2:1.0	ZnBr_2	90
17 ^d	$\text{Ph}_2\text{P}(\text{CH}_2)_2\text{PPh}_2$	0.6:1.2:0.8	ZnBr_2	99
18 ^e	$\text{Ph}_2\text{P}(\text{CH}_2)_2\text{PPh}_2$	0.6:1.2:0.8	ZnBr_2	91

^aEquivalent ratio of $\text{R}_3\text{P}/\text{ICH}_2\text{CH}_2\text{I}/\text{additive}$ relative to 1 equiv of substrate **1a**. ^bThe yields were determined by ^{19}F NMR spectroscopy. ^cThe reaction was performed in DMF (2 mL). ^dThe reaction time was 15 min. ^eThe reaction time was 5 min.


*Reaction conditions: alcohol **1** (0.5 mmol), Selectfluor (1.5 mmol), $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2$ (0.3 mmol), $\text{ICH}_2\text{CH}_2\text{I}$ (0.6 mmol), CH_3CN (5 mL), rt, 15 min, N_2 atmosphere. The yields were isolated yields. ^aA gram-scale reaction gave product **2a** in 88% yield. ^bThe yields were determined by ^{19}F NMR spectroscopy. Lower isolated yields were obtained due to high volatility of the products.

range of tertiary alcohols. Various functional groups could be tolerated, such as carbonyl, ester, sulfonate, sulfonamide, and primary halides. The distance between the ester moiety and hydroxyl group affects the reaction efficiency. A short distance led to a low yield (**2e**), probably because the activation of hydroxyl group was suppressed due to the competing coordination of the ester group with the activating Lewis acid generated from the $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2/\text{ICH}_2\text{CH}_2\text{I}$ system. With a long distance, the yield was not affected (e.g., **2f**, **2g**, and **2u**). Bulkier substituents would lead to a decrease

in the yields due to steric effects (**2g**, **2h**, and **2t**). Fluorination of complex molecules also occurred smoothly (**2u–2w**). Surprisingly, *cis*-cyclic alcohol **1x** was exclusively converted into *cis*-product **2x**. To further demonstrate the synthetic potential of this protocol, fluorination of substrate **1a** was carried out on a gram scale. To our delight, product **2a** was isolated in a high yield (1.17g, 88%).

We propose that the reaction may proceed via the first iodination or bromination of alcohols, and the subsequent fluorination of tertiary iodides or bromides. Indeed, the $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2$ /ICH₂CH₂I system can effectively convert alcohol **1a** into the corresponding iodide **3a** (Scheme 3, eq 1).

Scheme 3. Experimental Evidence for the Iodination and Bromination Paths




^aDetermined by ¹H NMR spectroscopy. ^bDetermined by ¹⁹F NMR spectroscopy.

When ZnBr₂ was present, bromide **4a** was also provided (eq 2). Fluorination of both iodide **3a** and bromide **4a** occurred rapidly (eqs 3 and 4), reflecting that iodination and bromination are involved in the fluorination of alcohols. Since the $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2$ /ICH₂CH₂I system alone can efficiently promote iodination, a question arose why bromination with ZnBr₂ should be involved to increase the fluorination yield. The reaction of $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2$ with ICH₂CH₂I would generate iodide anion,¹³ which could readily undergo a redox reaction with Selectfluor.¹⁵ The consumption of iodide anion and Selectfluor would suppress iodination and fluorination, respectively. However, ZnBr₂ cannot be completely dissociated to release bromide anion, and thus the redox reaction of ZnBr₂ with Selectfluor would not easily occur. Although not completely dissociated, ZnBr₂ still can effectively promote bromination. Therefore, ZnBr₂ is quite important for this fluorination process. Although the redox reaction of I[−] with Selectfluor would generate fluoride anion,¹⁵ nucleophilic fluorination could still be excluded on the basis that no desired product was observed by using CsF instead of Selectfluor (eq 5) and that bromide **4a** cannot be converted into fluorinated product within 15 min (eq 6).

I[−]/Selectfluor is not the only incompatible pair in the reaction system. $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2$ is not compatible with Selectfluor either.¹⁴ The issue of this incompatibility can be addressed by sequential addition of reagents because $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2$ reacts with ICH₂CH₂I very fast.

$\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2$ is not soluble in the reaction solvent, CH₃CN, and thus, it would be present as a solid in CH₃CN. However, after ICH₂CH₂I was added, the solid disappeared quickly with stirring the reaction mixture. No solid remained after stirring for about 1 min, and ³¹P NMR analysis revealed that $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2$ was almost completely transformed into a species whose resonance signal appearing at 51.3 ppm, which should correspond to a diiodophosphonium salt, $[\text{Ph}_2\text{P}^+(\text{I})\text{CH}_2\text{CH}_2\text{P}^+(\text{I})\text{Ph}_2] 2\text{I}^-$, according to our previous investigations (Scheme 4, eq 1).^{13a} The diiodophosphonium

Scheme 4. Need for Sequential Addition of Reagents




^aThe yield was determined by ¹H NMR spectroscopy.

salt is quite reactive and may easily undergo side reactions. However, if the substrate alcohol is added before ICH₂CH₂I, the phosphonium salt generated in situ may specifically coordinate to the alcohol and thus side reactions may be suppressed. Indeed, when an alcohol was present, the hydroxyl group was effectively activated and iodination product was obtained in 43% yield right after the solid disappeared (eq 2). For the fluorination procedure, after alcohol/ $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2$ /ICH₂CH₂I was added and the solid disappeared (about 1 min), ZnBr₂ and Selectfluor were then sequentially added without time interval. The issue of incompatibility could be avoided by this procedure.

Based on the above results, we propose the reaction mechanism as shown in Scheme 5. $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2$ reacts

Scheme 5. Plausible Reaction Mechanism



with ICH₂CH₂I rapidly to generate diiodophosphonium salt $[\text{Ph}_2\text{P}^+(\text{I})\text{CH}_2\text{CH}_2\text{P}^+(\text{I})\text{Ph}_2] 2\text{I}^-$, which can act as a Lewis acid to activate hydroxyl group for the iodination or bromination of alcohols.^{13a} A redox reaction of Selectfluor with I[−] anion produces radical cation **A**, which abstracts a halogen atom from tertiary iodides or bromides to deliver alkyl radical. Fluorination of this radical with Selectfluor provides the final product.⁴ The stereoselectivity for the formation of product **2x** should be because of a double-inversion process, iodination or bromination of alcohols with inversion of configuration,^{13a} and

fluorination of the alkyl radical with the second inversion. In this case, probably the alkyl radical and salt **B** are not completely dissociated but are present as a contact ion radical pair, resulting in a back-side combination to the fluorine atom.

In summary, we have described a convenient dehydroxylative fluorination of tertiary alcohols under mild conditions. Compared with previous methods for tertiary C–F bond formation, this protocol features high efficiency (a reaction time of 15 min) and the use of easily available alcohols as substrates. This work may represent a new discovery in fluorination of alcohols since the reported methods are mainly limited to primary and secondary alcohols. The approach may find utility in ^{18}F -labeling due to its high efficiency.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.orglett.0c02438>.

Experimental details; references and notes; copies of ^1H , ^{19}F , and ^{13}C NMR spectra (PDF)

■ AUTHOR INFORMATION

Corresponding Authors

Jin-Hong Lin – Key Laboratory of Organofluorine Chemistry, Shanghai Institute of Organic Chemistry, University of Chinese Academy of Sciences, Chinese Academy of Sciences, Shanghai 200032, China; orcid.org/0000-0002-7000-9540; Email: jlh@sioc.ac.cn

Ji-Chang Xiao – Key Laboratory of Organofluorine Chemistry, Shanghai Institute of Organic Chemistry, University of Chinese Academy of Sciences, Chinese Academy of Sciences, Shanghai 200032, China; orcid.org/0000-0001-8881-1796; Email: jchxiao@sioc.ac.cn

Authors

Wei Zhang – Key Laboratory of Organofluorine Chemistry, Shanghai Institute of Organic Chemistry, University of Chinese Academy of Sciences, Chinese Academy of Sciences, Shanghai 200032, China

Yu-Cheng Gu – Syngenta, Bracknell, Berkshire RG426EY, U.K.

Complete contact information is available at:

<https://pubs.acs.org/doi/10.1021/acs.orglett.0c02438>

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This Paper is Dedicated to Professor Youyou Tu, the 2015 Nobel Prize Laureate of Physiology or Medicine, on the Occasion of Her 90th Birthday. We thank the National Natural Science Foundation (21421002, 21672242, 21971252, 21991122), Key Research Program of Frontier Sciences, Chinese Academy of Sciences (CAS) (QYZDJSSW-SLH049), Youth Innovation Promotion Association CAS (2019256), and the Syngenta PhD Fellowship Awarded to W. Zhang for financial support.

■ REFERENCES

- (1) (a) Purser, S.; Moore, P. R.; Swallow, S.; Gouverneur, V. Fluorine in medicinal chemistry. *Chem. Soc. Rev.* **2008**, *37*, 320–30. (b) Ojima, I. *Fluorine in Medicinal Chemistry and Chemical Biology*; Blackwell Publishing: Chichester, 2009. (c) Kirsch, P. *Modern Fluoroorganic Chemistry: Synthesis, Reactivity, Applications*, 2nd ed.; Wiley-VCH: Weinheim, 2013. (d) Gillis, E. P.; Eastman, K. J.; Hill, M. D.; Donnelly, D. J.; Meanwell, N. A. Applications of Fluorine in Medicinal Chemistry. *J. Med. Chem.* **2015**, *58*, 8315–8359. (e) Meanwell, N. A. Fluorine and Fluorinated Motifs in the Design and Application of Bioisosteres for Drug Design. *J. Med. Chem.* **2018**, *61*, 5822–5880. (f) Wang, J.; Sánchez-Roselló, M.; Aceña, J. L.; del Pozo, C.; Sorochinsky, A. E.; Fustero, S.; Soloshonok, V. A.; Liu, H. Fluorine in Pharmaceutical Industry: Fluorine-Containing Drugs Introduced to the Market in the Last Decade (2001–2011). *Chem. Rev.* **2014**, *114*, 2432–2506. (g) Zhou, Y.; Wang, J.; Gu, Z.; Wang, S.; Zhu, W.; Aceña, J. L.; Soloshonok, V. A.; Izawa, K.; Liu, H. Next Generation of Fluorine-Containing Pharmaceuticals, Compounds Currently in Phase II–III Clinical Trials of Major Pharmaceutical Companies: New Structural Trends and Therapeutic Areas. *Chem. Rev.* **2016**, *116*, 422–518.
- (2) (a) O'Hagan, D.; Deng, H. Enzymatic fluorination and biotechnological developments of the fluorinase. *Chem. Rev.* **2015**, *115*, 634–649. (b) Kohlhepp, S. V.; Gulder, T. Hypervalent iodine(iii) fluorinations of alkenes and diazo compounds: New opportunities in fluorination chemistry. *Chem. Soc. Rev.* **2016**, *45*, 6270–6288. (c) Lee, J.-W.; Oliveira, M. T.; Jang, H. B.; Lee, S.; Chi, D. Y.; Kim, D. W.; Song, C. E. Hydrogen-bond promoted nucleophilic fluorination: Concept, mechanism and applications in positron emission tomography. *Chem. Soc. Rev.* **2016**, *45*, 4638–4650. (d) Yang, L.; Dong, T.; Revankar, H. M.; Zhang, C.-P. Recent progress on fluorination in aqueous media. *Green Chem.* **2017**, *19*, 3951–3992. (e) Szpera, R.; Moseley, D. F. J.; Smith, L. B.; Sterling, A. J.; Gouverneur, V. r. The Fluorination of C–H Bonds: Developments and Perspectives. *Angew. Chem., Int. Ed.* **2019**, *58*, 14824–14848.
- (3) Nishikata, T.; Ishida, S.; Fujimoto, R. Site-Selective Tertiary Alkyl-Fluorine Bond Formation from α -Bromoamides Using a Copper/CsF Catalyst System. *Angew. Chem., Int. Ed.* **2016**, *55*, 10008–10012.
- (4) Chen, H.; Liu, Z.; Lv, Y.; Tan, X.; Shen, H.; Yu, H. Z.; Li, C. Selective Radical Fluorination of Tertiary Alkyl Halides at Room Temperature. *Angew. Chem., Int. Ed.* **2017**, *56*, 15411–15415.
- (5) (a) Cheng, L.-J.; Cordier, C. J. Catalytic Nucleophilic Fluorination of Secondary and Tertiary Propargylic Electrophiles with a Copper-N-Heterocyclic Carbene Complex. *Angew. Chem., Int. Ed.* **2015**, *54*, 13734–13738. (b) Brioché, J. One-pot synthesis of tertiary alkyl fluorides from methyl oxalates by radical deoxyfluorination under photoredox catalysis. *Tetrahedron Lett.* **2018**, *59*, 4387–4391. (c) Aguilar Troyano, F. J.; Ballaschki, F.; Jaschinski, M.; Ozkaya, Y.; Gomez-Suarez, A. Light-Mediated Formal Radical Deoxyfluorination of Tertiary Alcohols through Selective Single-Electron Oxidation with TEDA $^{2+}$. *Chem. - Eur. J.* **2019**, *25*, 14054–14058. (d) Gonzalez-Esguevillas, M.; Miro, J.; Jeffrey, J. L.; MacMillan, D. W. C. Photoredox-catalyzed deoxyfluorination of activated alcohols with Selectfluor. *Tetrahedron* **2019**, *75*, 4222–4227. (e) Ma, J.; Xu, W.; Xie, J. Predictable site-selective radical fluorination of tertiary ethers. *Sci. China: Chem.* **2020**, *63*, 187–191.
- (6) (a) Yin, F.; Wang, Z.; Li, Z.; Li, C. Silver-Catalyzed Decarboxylative Fluorination of Aliphatic Carboxylic Acids in Aqueous Solution. *J. Am. Chem. Soc.* **2012**, *134*, 10401–10404. (b) Ventre, S.; Petronijevic, F. R.; Macmillan, D. W. C. Decarboxylative fluorination of aliphatic carboxylic acids via photoredox catalysis. *J. Am. Chem. Soc.* **2015**, *137*, 5654–5657. (c) Katada, M.; Kitahara, K.; Iwasa, S.; Shibatomi, K. Catalyst-Free Decarboxylative Fluorination of Tertiary β -Keto Carboxylic Acids. *Synlett* **2018**, *29*, 2408–2411. (d) Webb, E. W.; Park, J. B.; Cole, E. L.; Donnelly, D. J.; Bonacorsi, S. J.; Ewing, W. R.; Doyle, A. G. Nucleophilic (Radio)Fluorination of Redox-Active Esters via Radical-Polar Crossover Enabled by Photoredox Catalysis. *J. Am. Chem. Soc.* **2020**, *142*, 9493–9500.
- (7) (a) Markovskij, L. N.; Pashinnik, V. E.; Kirsanov, A. V. Application of Dialkylaminosulfur Trifluorides in the Synthesis of Fluoroorganic Compounds. *Synthesis* **1973**, *1973*, 787–789. (b) Mid-

dleton, W. J. New fluorinating reagents. Dialkylaminosulfur fluorides. *J. Org. Chem.* **1975**, *40*, 574–578.

(8) (a) Lal, G. S.; Pez, G. P.; Pesaresi, R. J.; Prozonic, F. M.; Cheng, H. Bis(2-methoxyethyl)aminosulfur Trifluoride: A New Broad-Spectrum Deoxyfluorinating Agent with Enhanced Thermal Stability. *J. Org. Chem.* **1999**, *64*, 7048–7054. (b) Beaulieu, F.; Beauregard, L.-P.; Courchesne, G.; Couturier, M.; LaFlamme, F.; L'Heureux, A. Aminodifluorosulfonium tetrafluoroborate salts as stable and crystalline deoxyfluorinating reagents. *Org. Lett.* **2009**, *11*, 5050–5053. (c) Umemoto, T.; Singh, R. P.; Xu, Y.; Saito, N. Discovery of 4-*tert*-Butyl-2,6-dimethylphenylsulfur Trifluoride as a Deoxyfluorinating Agent with High Thermal Stability as Well as Unusual Resistance to Aqueous Hydrolysis, and Its Diverse Fluorination Capabilities Including Deoxyfluoro-Arylsulfonylation with High Stereoselectivity. *J. Am. Chem. Soc.* **2010**, *132*, 18199–18205. (d) L'Heureux, A.; Beaulieu, F.; Bennett, C.; Bill, D. R.; Clayton, S.; LaFlamme, F.; Mirmehrab, M.; Tadayon, S.; Tovell, D.; Couturier, M. Aminodifluorosulfonium Salts: Selective Fluorination Reagents with Enhanced Thermal Stability and Ease of Handling. *J. Org. Chem.* **2010**, *75*, 3401–3411. (e) Li, L.; Ni, C.; Wang, F.; Hu, J. Deoxyfluorination of alcohols with 3,3-difluoro-1,2-diarylcyclopropanes. *Nat. Commun.* **2016**, *7*, 13320. (f) Guo, J.; Kuang, C.; Rong, J.; Li, L.; Ni, C.; Hu, J. Rapid Deoxyfluorination of Alcohols with *N*-Tosyl-4-chlorobenzenesulfonimidoyl Fluoride (SulfoxFluor) at Room Temperature. *Chem. - Eur. J.* **2019**, *25*, 7259–7264.

(9) Nielsen, M. K.; Ugaz, C. R.; Li, W.; Doyle, A. G. PyFluor: A Low-Cost, Stable, and Selective Deoxyfluorination Reagent. *J. Am. Chem. Soc.* **2015**, *137*, 9571–9574.

(10) Sladojević, F.; Arlow, S. I.; Tang, P.; Ritter, T. Late-Stage Deoxyfluorination of Alcohols with PhenoFluor. *J. Am. Chem. Soc.* **2013**, *135*, 2470–2473.

(11) Goldberg, N. W.; Shen, X.; Li, J.; Ritter, T. AlkylFluor: Deoxyfluorination of Alcohols. *Org. Lett.* **2016**, *18*, 6102–6104.

(12) Sood, D. E.; Champion, S.; Dawson, D. M.; Chhabra, S.; Bode, B. E.; Sutherland, A.; Watson, A. J. B. Deoxyfluorination with CuF₂: Enabled by Using a Lewis Base Activating Group. *Angew. Chem., Int. Ed.* **2020**, *59*, 8460–8463.

(13) (a) Chen, J.; Lin, J.-H.; Xiao, J.-C. Halogenation through Deoxygenation of Alcohols and Aldehydes. *Org. Lett.* **2018**, *20*, 3061–3064. (b) Chen, J.; Lin, J.-H.; Xiao, J.-C. Dehydroxylation of alcohols for nucleophilic substitution. *Chem. Commun.* **2018**, *54*, 7034–7037. (c) Zhang, W.; Chen, J.; Lin, J.-H.; Xiao, J.-C.; Gu, Y.-C. Rapid Dehydroxytrifluoromethoxylation of Alcohols. *iScience* **2018**, *5*, 110–117.

(14) Yang, Z.; Chen, S.; Yang, F.; Zhang, C.; Dou, Y.; Zhou, Q.; Yan, Y.; Tang, L. PPh₃/Selectfluor-Mediated Transformation of Carboxylic Acids into Acid Anhydrides and Acyl Fluorides and Its Application in Amide and Ester Synthesis. *Eur. J. Org. Chem.* **2019**, *2019*, 5998–6002.

(15) Gilcinski, A. G.; Pez, G. P.; Syvret, R. G.; Lal, G. S. On the relative power of electrophilic fluorinating reagents of the N-F class. *J. Fluorine Chem.* **1992**, *59*, 157–162.