



## Conformationally Constrained Serotonin Analogues: Stereoselective Synthesis of *trans*-3-(2-aminocycloalkyl)Indoles by Aziridine Ring Opening

Jesús Ezquerro, Concepción Pedregal and Carlos Lamas\*

Centro de Investigación Lilly, S. A. Paraje de la Cruz S/N, 28130 Valdeolmos, Madrid, Spain.

Alfredo Pastor, Pilar Alvarez and Juan José Vaquero

Departamento de Química Orgánica, Universidad de Alcalá, 28871 Alcalá de Henares, Madrid, Spain.

**Abstract:** *Trans*-3-(2-aminocyclopentyl)indoles **9a,c** and *trans*-3-(2-aminocyclohexyl)indoles **9b,d** have been stereoselectively prepared by nucleophilic ring opening reaction of *N*-Boc cycloalkylaziridines **6a,b** and the "lower order" magnesium cuprate **II**, generated from the corresponding indolylmagnesium bromides derived from **7a,b**.

The tryptamine nucleus is prevalent in many natural occurring products.<sup>1</sup> Serotonin (**1**, 5-HT) (Figure 1) is one of the most interesting due to its many physiological functions. The molecular manipulation of the serotonin aromatic ring has enabled the development of selective serotonergic drugs such as *Sumatriptan* **2**<sup>2</sup> or benzofuran bioisosteres of tryptamines like **3**<sup>3</sup> among many others.<sup>4</sup> The synthesis of restricted analogues of the tryptamines<sup>5</sup> has provided a useful insight into the ligand recognition requirements for the receptor subtypes.




Figure 1


The synthesis of tryptamines **4** (n=1,2)<sup>6</sup> has been accomplished by the reaction of indolylmagnesium bromide with the corresponding cycloalkene epoxide to yield the 3-(2-hydroxycycloalkyl)indole. Subsequent oxidation of the alcohol to the ketone and then reductive amination, using the Danheiser conditions,<sup>7</sup> gave the *trans* isomers of **4** (n=2) as the major product in the reaction mixture (8:1 ratio). On the other hand, the

conversion of the ketone to the enamine under  $\text{TiCl}_4$  catalysis, followed by reduction with  $\text{NaCNBH}_3$  yielded the *cis* tryptamine.<sup>6b</sup>

Aziridines are very useful synthetic intermediates<sup>8</sup> and react with nucleophiles to form 2-substituted ethylamines. Aziridines require activation by an electronwithdrawing protecting group on the ring nitrogen atom. The most used protecting group has been the *p*-toluenesulphonyl<sup>9</sup> (Ts) and others, like diphenylphosphinyl<sup>10</sup> (Dpp), which avoids the problems associated with the cleavage of the sulphonamide bond.<sup>11</sup> Less attention has been paid to *N*-urethane aziridines<sup>12</sup> due to the problems associated with their preparation<sup>13</sup> and lack of reactivity with hard nucleophiles.<sup>9a</sup>

In this communication we would like to report a new stereoselective synthesis of 3-(2-aminocycloalkyl)indoles by means of nucleophilic *N*-Boc aziridine ring opening.

The activated aziridines **6a,b** (Scheme 1) were prepared by addition of iodine isocyanate to either cyclopentene or cyclohexene. Subsequent treatment with potassium *tert*-butoxide in DMF of the corresponding *trans* 2-iodo-1-isocyanates **5a** (95% yield) and **5b** (90% yield) gave rise to the *N*-Boc aziridines **6a** (75% yield) and **6b** (85% yield).




Scheme 1

This procedure represents a modification of the previously reported one by Hassner<sup>14</sup> where **5b** was treated first with an alcohol, yielding the *trans* 2-iodo-1-carbamate which was further transformed into the *N*-urethane aziridine when treated with base (NaH).

The exemplification of this nucleophilic ring opening reaction (Scheme 2) has been carried out using commercially available 5-methylindole (**7a**) and 7-chloroindole (**7b**), prepared using Bartoli's methodology.<sup>15</sup>

Indolylmagnesium bromides derived from **7a** and **7b** were prepared from the corresponding indoles and  $\text{CH}_3\text{MgBr}$  following literature procedures.<sup>6c</sup> Addition of 0.3 equivalents of  $(\text{CH}_3)_2\text{S}\cdot\text{CuBr}$  resulted in the formation of a "lower order" magnesium cuprate<sup>16</sup> **II** which upon reaction with the *N*-Boc aziridines **6a,b** gave the *trans* cycloalkylamines **8a-d**. Yields are dependent on the substituent present in the reacting indole. Thus, electron donating substituents gave better yields [**8a** (85%); **8b** (80%)] while indoles having electron withdrawing substituents gave lower yields in this nucleophilic ring opening reaction [**8c** (37%); **8d** (43%)]. This reactivity has been observed previously by Kozikowski<sup>12a</sup> in the synthesis of tryptophanes from aziridine-2-carboxylates and various substituted indoles in the presence of zinc triflate as Lewis acid.

Finally, the *N*-Boc protected cycloalkylamines **8a-d** were deprotected to give the amines hydrochlorides **9a-d** with HCl (g) in a 1:1 mixture of  $\text{CH}_2\text{Cl}_2/\text{Et}_2\text{O}$  in almost quantitative yields.



Scheme 2

The *trans* stereochemistry of compounds **9** was ascertain by  $^1\text{H}$  NMR (Figure 2) by measuring the diaxial coupling constant<sup>17</sup> ( $J_{\text{H}_{1\text{ax}},\text{H}_{2\text{ax}}} = 11 \text{ Hz}$ ) for **9b** and by nOe experiments on the *cis* isomer **10**<sup>18</sup> which displayed an 7% enhancement of  $\text{H}_1$  when  $\text{H}_2$  was irradiated. When the same experiment was performed on the parent compound **9b** no substantial nOe was observed.




Figure 2

In conclusion, the presented method represents a novel approach to obtain conformationally constrained tryptamines in a stereoselective fashion by the nucleophilic ring opening of non-polarised *N*-urethane activated aziridines.<sup>19</sup> The easy removal of the aziridine activating group (Boc) makes this procedure superior to the more classical *N*-sulfonyl aziridines, where the final cleavage of the sulphonamide bond is sometimes difficult. Further applications of this methodology to other non-cyclic derived aziridines is in progress in these laboratories and will be reported in due course.

**Acknowledgements:** This research was supported by the Spanish **FARMA III** programme (Ministerio de Industria y Ministerio de Sanidad). A. P. and P. A. are grateful to Lilly, S. A. for a fellowship.

We are also grateful to C. Ramírez for the preparation of some aziridines. We are also grateful to B. P. Clark and S. R. Baker (Lilly Research Center, U.K.) for useful suggestions in the preparation of this manuscript.

### References and Notes

1. Spande, T. F. in *The Chemistry of Heterocyclic Compounds*. Weissberger, A.; Taylor, E. C., Eds. John Wiley & Sons. 1979, Vol. 25 (III), Chapter VIII.
2. Feniuk, W.; Humphrey, P. P. A. *Drug Dev. Res.* **1992**, *26*, 235-240.
3. Tomaszewski, Z.; Johnson, M. P.; Huang, X.; Nichols, D. E. *J. Med. Chem.* **1992**, *35*, 2061-2064.
4. Hibert, M. F.; Mir, A. K. Fozard, J. R. in *Comprehensive Medicinal Chemistry*. Emmett, J. C. Ed. Pergamon Press **1990**. Vol. 3, Chapter 12.9.
5. Macor, J. E. *Tetrahedron Lett.* **1995**, *36*, 7019-7022 and references cited therein.
6. (a) Macor, J. E.; Ryan, K. *Heterocycles* **1990**, *31*, 1497-1504. (b) Ghosh, A.; Wang, W.; Freeman, J. P.; Althaus, J. S.; VonVolgtlander, P. F.; Scahill, T. A.; Mizesak, S. A.; Szmuszkowicz, J. *Tetrahedron* **1991**, *47*, 8653-8662. (c) Audia, J. E.; Colocci, N. *Tetrahedron Lett.* **1991**, *32*, 3779-3782.
7. Danheiser, R. L.; Morin, J. M. Jr.; Salaski, E. J. *J. Am. Chem. Soc.* **1985**, *107*, 8066-8073
8. For a review see: Tanner, D. *Angew. Chem. Int. Ed. Engl.* **1994**, *33*, 599-619.
9. (a) Lygo, B. *Synlett* **1993**, 764-766 and references cited therein. (b) Baldwin, J. E.; Adlington, R. M.; O'Neil, I. A.; Schofield, C.; Spivey, A. C.; Sweeney, J. B. *J. Chem. Soc., Chem. Commun.* **1989**, 1852-1854.
10. (a) Osborn, H. M. I.; Sweeney, J. B.; Howson, B. *Tetrahedron Lett.* **1994**, *35*, 2739-2742. (b) Osborn, H. M. I.; Sweeney, J. B.; Howson, B. *Synlett* **1994**, 145-147.
11. The use of SmI<sub>2</sub> has shown to be quite effective in these deprotection reactions. Vedejs, E.; Lin, S. J. *Org. Chem.* **1994**, *59*, 1602-1603.
12. Urethane activation of aziridines and reaction with indole under Lewis acid catalyst, has been described for the synthesis of different tryptophanes and other derivatives: (a) Sato, K.; Kozikowski A. P. *Tetrahedron Lett.* **1989**, *30*, 4073-4076. (b) Shima, I.; Shimazaki, N.; Imai, K.; Hemmi, K.; Hasimoto, M. *Chem. Pharm. Bull.* **1990**, *38*, 564-566. (c) Legters, J.; Thijs, L.; Zwanenburg, B. *Recl. Trav. Chim. Pays-Bas.* **1992**, *111*, 16-21. (d) Dubois, L.; Mehta, A.; Tourette, E.; Dodd, R. H. *J. Org. Chem.* **1994**, *59*, 434-441.
13. (a) Nakajima, K.; Kawai, H.; Takai, M.; Okawa, K. *Bull. Chem. Soc. Japan*, **1977**, *50*, 917-920. (b) Nakajima, K.; Takai, F.; Tanaka, T.; Okawa, K. *Bull. Chem. Soc. Japan*, **1978**, *51*, 1577- 1578.
14. Hassner, A.; Heathcock, C. *Tetrahedron* **1964**, *20*, 1037-1042.
15. Bartoli, G.; Palmieri, G.; Bosco, M.; Dalpozzo, R. *Tetrahedron Lett.* **1989**, *30*, 2129-2132.
16. Lipshutz, B. H. in "Comprehensive Organic Synthesis", Trost, B. M.; Fleming, I.; Eds. Pergamon, Oxford **1991**; Vol. 1, Chapter 1.4.
17. Karplus, M. *J. Am. Chem. Soc.* **1963**, *85*, 2870-2871
18. This compound was prepared by reductive amination as described in *ref.* 5a
19. Satisfactory spectroscopic data (<sup>1</sup>H NMR, <sup>13</sup>C NMR, IR and HRMS) have been obtained for all compounds reported in this communication.

(Received in UK 13 October 1995; accepted 24 November 1995)