

Tellurium tetrachloride: an improved method of preparation

Nicola Petragnani^{a,*}, Samuel R. Mendes^b, Claudio C. Silveira^{b,*}

^a Instituto de Química, Universidade de São Paulo, Caixa Postal 26077, 05599-970 São Paulo, SP, Brazil

^b Departamento de Química, Universidade Federal de Santa Maria, Caixa Postal 5001, 97105-970 Santa Maria, RS, Brazil

Received 13 December 2007; revised 28 January 2008; accepted 14 February 2008

Available online 19 February 2008

Abstract

An efficient and practical synthesis of tellurium tetrachloride from elemental tellurium and sulfuryl chloride is described.
© 2008 Elsevier Ltd. All rights reserved.

Keywords: Tellurium tetrachloride; Sulfuryl chloride; Tellurium

Tellurium, once regarded as an exotic chemical element, became the key component of a variety of versatile and useful reagents for organic synthesis. The multiple applications of organotellurium chemistry have been well described in a number of review articles and books.^{1–8}


Of the many inorganic tellurium reagents employed to introduce tellurium in organic substrates, such as elemental tellurium, alkali tellurides and ditellurides, hydrogen telluride and tellurium dioxide, tellurium tetrachloride deserves a special role. TeCl₄ reacts with activated aromatic compounds, with organo-chloromercury compounds, with aromatic Grignard reagents and with several active methylene groups; adds to double and triple C–C bond; and participates in the synthesis of tellurium heterocycles.⁸

Tellurium tetrachloride has been usually prepared by the reaction of elemental tellurium with chlorine gas. In spite of the high yields attained, the procedure is rather tedious and laborious, certainly in agreement with the opinion of previous researchers, since a particular glass apparatus^{8,9} is used, in which the tellurium tetrachloride produced is distilled under a continuous chlorine stream into connected tubes. After solidification of the product, the tubes are detached from the apparatus. By this procedure, tellurium tetrachloride is a pale yellow crystalline solid (mp 225 °C).

We describe here a very simple and straightforward procedure for the preparation of tellurium tetrachloride, which employs sulfuryl chloride (SO₂Cl₂) as chlorine source (Scheme 1).

By heating at reflux a suspension of elemental tellurium in excess of sulfuryl chloride, a white to light-gray solid of the tellurium tetrachloride is formed gradually.¹⁰ The yield is almost quantitative. The excess of sulfuryl chloride can be recovered and reused. For most purposes, after the removal of the excess of sulfuryl chloride the reagent is ready to be used in situ. With the aim of proving the merit of this new method, some previously described reactions were performed, attaining good, compatible results:


- Reaction of TeCl₄ with activated aromatic compounds giving aryl tellurium trichlorides **2a–b** and ditellurides **3a–b** by NaBH₄ reduction in situ (Scheme 2).^{11–13}
- Addition of TeCl₄ to phenyl acetylene giving 2-chloro-2-phenylvinyltellurium trichloride **5**, and by the use of phenyl acetylene in excess the corresponding bis-vinyl telluride **6** is formed (Scheme 3).^{14,15}
- Reaction of TeCl₄ with 4 equiv of aryl magnesium bromides, giving symmetrical diaryl tellurides (Scheme 4).^{16,17}




Scheme 1.

* Corresponding authors. Tel./fax: +55 55 32208754 (C.C.S.).


E-mail address: silveira@quimica.ufsm.br (C. C. Silveira).




Scheme 2.



Scheme 3.



Scheme 4.



Scheme 5.

– Reaction of TeCl₄ with alkenes, such as cyclohexene, in ethanol giving 2-ethoxy cyclohexyltellurium trichloride (**9**) (Scheme 5).¹⁸

By our method, the preparation of TeCl₄ can be performed both in small amounts (1–2 mmol) and in large amounts, such as starting with 10 g of tellurium, with comparable high yields.

In conclusion, we described here a very easy and convenient method for the preparation of TeCl₄ that can be used in situ for further transformations or isolated and stored. The procedure is quite simple, for example, allowing the one-pot preparation of diaryl ditellurides from elemental tellurium in very high yields.

Acknowledgement

The authors thank FAPERGS, MCT/CNPq, and CAPES for financial support.

References and notes

1. *Tellurium*; Cooper, W. C., Ed.; Van Nostrand Reinhold: New York, 1971.
2. Engman, L. *Acc. Chem. Res.* **1985**, *18*, 274.
3. Petragnani, N.; Comasseto, J. V. *Synthesis* **1986**, 1.
4. Irgolic, K. Y. In *Houben-Weyl Methods of Organic Chemistry*, 4th ed.; Klamann, D., Ed.; Georg Thieme: Stuttgart, 1990; Vol. E12b.
5. Petragnani, N.; Comasseto, J. V. *Synthesis* **1991**, 793–897.
6. Petragnani, N. In *Tellurium in Comprehensive Organometallic Chemistry II*; McKillop, A., Ed.; Pergamon Elsevier, 1995; Vol. 11, Chapter 14.
7. Petragnani, N.; Stefani, H. A. *Tetrahedron* **2005**, *61*, 1613.
8. Petragnani, N.; Stefani, H. A. *Tellurium in Organic Synthesis—Best Synthetic Methods*, 2nd ed.; Academic Press: London, 2007.
9. Suttle, Y. F.; Smith, C. R. F. *Inorg. Synth.* **1956**, *III*, 140.
10. *Preparation of TeCl₄*: In a two neck flask, equipped with a reflux condenser and a drying tube, tellurium (10.16 g, 80 mmol) was treated with sulfuryl chloride (50 mL; excess) added dropwise. The suspension was refluxed for 12 h, cooled to room temperature and the excess of sulfuryl chloride distilled to dryness, leaving a white to light-gray solid in the flask. Argon atmosphere was then introduced. The solid was washed with dry hexanes (2 × 50 mL), the solvent removed with a syringe and TeCl₄ is ready to use for most purposes. It can be further dried under vacuo and transferred to a glass bottle. Isolated yield: 19.71 g (91.6%); mp 215 °C (lit.⁹ 225 °C).
11. Petragnani, N. *Tetrahedron* **1960**, *11*, 15 and references cited therein.
12. Chieffi, A.; Menezes, P. H.; Comasseto, J. V. *Organometallics* **1997**, *16*, 809.
13. *Preparation of aryltellurium trichlorides and diaryl ditellurides*: A one-necked round-bottomed flask containing tellurium tetrachloride (1.35 g, 5 mmol; prepared as above from 5 mmol of Te) was heated to 120 °C and the corresponding aromatic compound (5 mmol) was added at once, dissolving TeCl₄ promptly. An evolution of HCl was observed and a yellow solid was formed. The system was cooled to room temperature and the trichloride was used crude in the reduction procedure with NaBH₄, according to a procedure described by Comasseto.¹⁶ *Bis(4-methoxyphenyl)ditelluride (3a)*: Yield 1.10 g (94%); mp 57.6–59.0 °C (lit.²⁰ mp 58–59 °C) *Bis(4-methylphenyl)ditelluride (3b)*: Yield 0.89 g (83%); mp 50.1–51.2 °C (lit.¹⁹ mp 51 °C).
14. Moura Campos, M.; Petragnani, N. *Tetrahedron* **1962**, *18*, 527.
15. Cunha, R. L. O. R.; Zuckerman-Schpector, J.; Caracelli, I.; Comasseto, J. V. *J. Organomet. Chem.* **2006**, *691*, 4807; Uemura, S.; Miyoshi, H.; Okano, M. *Chem. Lett.* **1979**, 1357.
16. Rheinboldt, H.; Petragnani, N. *Chem. Ber.* **1956**, *89*, 1270.
17. McWhinnie, W. R.; Patel, M. G. *J. Chem. Soc., Dalton Trans.* **1972**, 199.
18. Sabir Ali, M. E.; Azad Malik, M.; Smith, B. C. C. *Inorg. Chim. Acta* **1989**, *162*, 157.
19. Haller, W. S.; Irgolic, K. J. *J. Organomet. Chem.* **1972**, *38*, 97.
20. Bergman, J. *Tetrahedron* **1972**, *28*, 3323.