




Fig. 1. ORD curves of the acetoxymethane (V).
— in isooctane
--- in methanol

dependent on the polarity of the solvents was not observed.

Experimental

The Hydroxylation of *trans-p*-Menth-2-ene (I) with Peracetic Acid.—*trans-p*-Menth-2-ene (I), b. p. 55–56°C/15 mmHg, $d_4^{25}=0.8160$, $[\alpha]_D^{25}=+120.46$ (neat), was prepared from 1-menthyl tosylate.

I (15 g.) was added drop by drop over a half-hour period to a solution of 30% hydrogen peroxide (25 ml.) in acetic acid (120 ml.) containing a few drops of concentrated sulfuric acid and warmed at 50°C. After the addition of I, the reaction mixture was stirred for another hour. The acetic acid was then removed in vacuo. The residue was dissolved in ether, washed with saturated sodium bisulfite to destroy the excess peracid, and then washed with aqueous sodium carbonate. The evaporation of the dried (sodium sulfate) ether solution yielded a viscous liquid (14.5 g.), b. p. 120–128°C/3 mmHg. On standing at room temperature the liquid gradually deposited crystals of 2-hydroxy-3-acetoxymethane (III) (1.5 g.); after recrystallization from ligroin, it had an m. p. of 114–116°C, $[\alpha]_D^{25}=+16.8^\circ$ (c 2.10, methanol) (Found: C, 66.82; H, 10.52. $C_{12}H_{22}O_3$ requires C, 67.25; H, 10.35%). The acetylation of III with acetic anhydride and sodium acetate gave the diacetate IV, m. p. 48–50°C

(Found: C, 65.82; H, 9.41. $C_{14}H_{24}O_4$ requires C, 65.59; H, 9.44%).

The filtrate (10 g.) was heated in 20% alcoholic potash (100 ml.) under reflux for an hour. After acidification with hydrochloric acid, the product was isolated by ether extraction to yield *p*-menthane-2,3-diol (II) (8 g.) b. p. 98–100°C/0.1 mmHg, $[\alpha]_D^{25}=+66.4^\circ$ (c 2.12, benzene). Diacetate: m. p. and mixed m. p. with acetate IV of III 48–50°C. Di-*p*-nitrobenzoate: m. p. 179–180°C (Found: N, 5.86. $C_{24}H_{26}O_8N_2$ requires N, 5.96%).

The Chromic Acid Oxidation of 2-Hydroxy-3-acetoxymethane (III).—A solution of 2-hydroxy-3-acetoxymethane (III) (3 g.) in pyridine (20 ml.) was added drop by drop to a suspension of chromic anhydride (4.5 g.) in pyridine (45 ml.). After the addition of III, the reaction mixture was stirred for five days at room temperature. After acidification with dilute sulfuric acid, the product was isolated by continuous extraction with ether (five days). The evaporation of the ether gave crude 3-acetoxycarvomenthone (V), which, after purification by vacuum sublimation (120°C (bath), 0.03 mmHg), had an m. p. of 46–47°C; yield 2.1 g. (Found: C, 68.04; H, 9.85. $C_{12}H_{20}O_3$ requires C, 67.89; H, 9.50%). 2,4-Dinitrophenylhydrazone: m. p. 157°C (from alcohol) (Found: N, 14.16. $C_{18}H_{24}O_6N_4$ requires N, 14.28%).

The ORD curves were measured in methanol and in isooctane.

In methanol (c 0.1025):

$[\alpha]_{318}=+975.6^\circ$ (peak)
 $[\alpha]_{280}=-1326.8^\circ$ (trough)

In isooctane (c 0.1035):

$[\alpha]_{322}=+1078.4^\circ$ (peak)
 $[\alpha]_{308}=-850.3^\circ$ (trough)

The Reduction of 3-Acetoxycarvomenthone (V).

—A mixture of 3-acetoxycarvomenthone (V) (2 g.), ethane dithiol (6.5 ml.) and boron trifluoride etherate (6.5 ml.) was allowed to stand for four hours at room temperature. After the addition of water, the product was extracted with ether. The ethereal solution was washed with dilute sodium hydroxide and dried over sodium sulfate. The evaporation of the ether yielded the thioketal of V. Without further purification the thioketal was heated with Raney nickel (from 30 g. of Raney alloy) in dioxane for five hours. After filtration and the removal of the solvent, the residue was treated with *p*-nitrobenzoyl chloride in pyridine at room temperature. The *p*-nitrobenzoate after recrystallization from alcohol showed an m. p. and mixed m. p. with *p*-nitrobenzoate of *d*-neomenthol of 96–99°C. The two *p*-nitrobenzoates showed identical infrared spectra.

Department of Chemistry
Tohoku University
Katahira-cho, Sendai