Enantioselective H-Bond-Directing Approach for Trienaminemediated Reactions in Asymmetric Synthesis**

Łukasz Albrecht, Fabio Cruz Acosta, Alberto Fraile, Anna Albrecht, Jannie Christensen, and Karl Anker Jørgensen*

Control of remote stereocenters constitutes an important challenge in modern organic synthesis.^[1,2] Despite limited generality, asymmetric strategies enabling such functionalizations have found important applications in target-oriented synthesis, thereby indicating their potential.^[1e,2] In recent years, dienamine- and trienamine-mediated reactions have emerged as new tools that enable stereoselective functionalizations of unsaturated aldehydes at the remote γ - or ϵ positions, respectively.^[3,4] In this context, trienamine-based strategies that allow stereocontrolled reactions at the carbon atom seven bonds away from the stereocenter of the catalyst are notable.^[4] Classically, such aminocatalytic transformations are realized using the steric shielding principle for stereochemical induction.^[5] However, important restrictions related to the dienophile scope still exist. One solution to this problem might rely on the application of H-bond-directing aminocatalysts in trienamine chemistry (Scheme 1). Benefits of such strategies relate not only to the dual activation of the reaction partners, which positions them correctly in space, thereby providing a well-defined stereochemical environment; more importantly, such strategies open the possibility of accessing new reactivities and reaction profiles resulting in further development of this field of chemistry.



- Independent activation of both reagents
- Broadened applicabilty of trienamine-mediated reactions

Scheme 1. Remote functionalizations through H-bond-directed trienamine chemistry.

 [*] Dr. Ł. Albrecht, F. Cruz Acosta, Dr. A. Fraile, Dr. A. Albrecht, J. Christensen, Prof. Dr. K. A. Jørgensen
 Center for Catalysis, Department of Chemistry, Aarhus University 8000 Aarhus C (Denmark)
 E-mail: kaj@chem.au.dk
 [**] This work was made possible by grants from Aarhus University,

- OChemSchool, Carlsberg Foundation, and FNU. F.C.A. thanks CSIC for predoctoral JAE grant. We thank Dr. Jacob Overgaard for performing X-ray analysis.
- Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/anie.201204790.

Development of new synthetic strategies that lead to pharmacologically active structural units is of high importance for the chemical community and constitutes an important part of the drug discovery process. The tetrahydroxanthone structural unit (Scheme 2a) is often encountered in nature and among pharmacologically active compounds.^[6]



Scheme 2. a) Importance of the tetrahydroxanthone structural unit, b) natural products containing this unit, and c) novel enantioselective approach to this class of compounds.

Blennolide A, secalonic acid D, and α -diversonolic ester (Scheme 2b) constitute representative examples of a rich family of natural products containing the tetrahydroxanthone unit. These compounds were isolated from fungi, bacteria, and lichens and are well-recognized for their antibiotic activity.^[6,7]

Given the biological properties of optically active tetrahydroxanthone derivatives and their wide occurrence in nature, studies on a synthetic methodology that offers a general and stereoselective access to this class of compounds were initiated. It was anticipated that such an approach for the preparation of tetrahydroxanthone derivatives could rely on the trienamine-mediated reaction between chromones and 2,4-dienals (Scheme 2 c).

Herein, we present a new, enantioselective approach to biologically relevant tetrahydroxanthone derivatives based on a novel trienamine-mediated [4+2] cycloaddition with

Angew. Chem. Int. Ed. 2012, 51, 1-6

© 2012 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim



These are not the final page numbers!



1

2

activated chromones as dienophiles. Furthermore, the possibility to control the stereochemical outcome of organocatalytic trienamine-mediated reactions by an H-bond-directing aminocatalyst is demonstrated.

Initial experiments identified the presence of the cyano group in the 3-position of the starting chromone as a crucial factor ensuring high reactivity, and more importantly, high diastereoselectivity of the tested cycloaddition. Accordingly, optimization studies were initiated using 5-methyl-2,4-hexadienal **1a** and 3-cyanochromone $2a^{[8]}$ as model substrates (Table 1; for full screening results, see the Supporting

Table 1: Enantioselective synthesis of tetrahydroxanthone 3a: optimization studies.^[a]



3	5	0.33	-	20	85	>20:1	45 ^[e]
4	6	0.33	-	20	67	>20:1	76
5	7 a ^[f]	0.33	DEA(4)	20	75	>20:1	87
6	7 b ^[f]	0.33	DEA(4)	20	73	>20:1	82
7	7 a ^[f]	0.1	DEA(4)	48	92	>20:1	89
8	7 a ^[f]	0.05	DEA(4)	48	74	>20:1	n.d.
9	7 a ^[f]	0.05	DEA(8)	48	92 (91)	>20:1	90
10	7 a ^[f]	0.05	DEA(16)	48	87	>20:1	89
11	7 a ^[f]	0.05	DEA(32)	48	> 95	>20:1	86

[a] Reactions performed on a 0.1 mmol scale (see the Supporting Information for details). Tf=trifluoromethanesulfonyl, TFA=trifluoroacetic acid. [b] Conversion of 2a as determined by ¹H NMR spectroscopy. The yield of isolated product **3a** is given in parentheses. [c] Determined by ¹H NMR spectroscopy. [d] Determined after Ramirez olefination by HPLC using a chiral stationary phase. [e] ent-3 a was obtained. [f] 16 mol% of catalyst applied.

Information). Disappointingly, when classical sterically demanding catalysts 4 or 5 were applied (Table 1, entries 1-3), moderate enantioselectivities (45-60% ee) were obtained. To overcome these difficulties we turned our attention to an H-bond-directed catalytic approach.[4f,9] It was reasoned that additional activation of 2a through hydrogen bonding should lead to an increase in its reactivity and provide a well-defined stereochemical environment resulting in enhanced enantioselectivity. To our delight, application of the H-bond-directing catalysts 6 or 7 (Table 1, entries 4-6) led to a significant improvement of the enantioselectivity (76-87% ee) with squaramide-based organocatalyst 7a being the most effective (Table 1, entry 5). It should be noted, that for catalysts 7a,b the use of N,N-diethylacetamide (DEA) was necessary to achieve good catalytic activity. Further screening concerning the effect of concentration (Table 1, entries 5, 7, 8) revealed that higher dilutions were beneficial for the reaction outcome. Finally, the amount of additive was screened (Table 1, entries 8-11), and the results indicate the significant influence of the amount of additive on both reactivity and enantioselectivity of the reaction.

After having established the optimal reaction conditions for the H-bond-directed [4+2] cycloaddition, we turned the attention to the scope of the methodology (Table 2). Delightfully, the developed reaction proved general, as both electronrich (Table 2, entries 2-4) and electron-poor chromones (Table 2, entries 5-7) could be applied to afford the target products 3b-g in good yields (59-98%) and with high enantioselectivities (88-91% ee). Furthermore, in all of the cases, complete diastereoselectivity was observed. Longer reaction times were required to achieve full conversion for chromones 2b-d and 2g, thus indicating the influence of both

Table 2: Enantioselective synthesis of tetrahydroxanthones 3: 3-cyanochromone scope.^[a]



[a] Reactions were performed on a 0.2 mmol scale (see the Supporting Information for details). [b] Determined by ¹H NMR spectroscopy. [c] Determined after Ramirez olefination by HPLC using a chiral stationary phase.

www.angewandte.org

2

© 2012 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

These are not the final page numbers!

electronic and steric factors on the reactivity of the 3cyanochromones.

In further studies, the possibility to utilize differently substituted 2,4-dienals 1 in the developed H-bond-directed [4+2] cycloaddition was evaluated (Scheme 3). To our delight, different substituents could be present in the starting



Scheme 3. Enantioselective synthesis of tetrahydroxanthones **3**: 2,4-dienal scope (see the Supporting Information for details).

2,4-dienals **1**, hereby further increasing the structural diversity of the obtained products **3**. Moreover, the introduction of an additional stereogenic center in the 4-position of the target tetrahydroxanthones proved possible. Notably, the reactions proceeded with a very high and unprecedented^[4] *trans*relationship between the C1 and C4 stereogenic centers in **3j**,**k** that originate from the starting 2,4-dienals **1** (for a discussion of the stereochemical outcome of the reaction, see Scheme 6). Importantly, less reactive and more challenging, linear substrates could also be employed as demonstrated for 2,4-hexadienal **1f** (**1** with $R^1 = R^2 = R^3 = H$ in Scheme 3) providing **3**.

After the successful development of an H-bond-directed approach for trienamine-mediated reactions, the possibilities of synthetic utilization of the obtained cycloadducts **3** were investigated. We were particularly interested in the application of the densely functionalized molecular scaffold of **3** for the construction of various polycyclic structures. Initially we investigated the possibility to introduce a δ -lactone ring system to the target products in a sequence of reactions that involves reduction of the aldehyde moiety and a subsequent hydrolytic cyclization onto the inherent cyano group. When tetrahydroxanthone **3a** was subjected to NaBH₄ reduction, both carbonyl groups in **3a** were reduced (Scheme 4). Importantly, the diastereoselectivity of this process could be



Scheme 4. Enantioselective synthesis of tetrahydroxanthones 9 and 12 that contain δ -lactone rings. DCE = dichloroethane.

fully controlled by decreasing the reaction temperature to -20 °C, thereby affording 8 as a single diastereoisomer. Interestingly, when NaBH(OAc)₃ was employed as the reducing agent, chemoselective reduction of the aldehyde moiety could be achieved, thereby affording 10 in a mixture with its corresponding hemiacetal 11 in a 3:1 ratio. Subsequently, the crude reaction mixtures were subjected to hydrolytic cyclization under acidic conditions. In such a manner, the δ -lactone ring was furnished affording 9 and 12 in good overall yields and as single diastereoisomers.

Further application studies were focused on the utilization of the carbonyl groups of 3 for the introduction of an additional tetrahydropyrane ring in the target products. Two complementary routes enabling such functionalization were elaborated. The first was initiated by a Wittig reaction of tetrahydroxanthone 3a with stabilized ylide 13 followed by chemo- and diastereoselective reduction of the ketone moiety (Scheme 5). A subsequent, TFA-mediated, intramolecular oxa-Michael reaction of 14 afforded the target product 15 in a high 82% overall yield. Notably, the developed reaction sequence allows the introduction of two new stereogenic centers in a fully diastereoselective manner. In the second approach, chemoselective reduction of the aldehyde moiety of 3a was performed as the first step. Hereafter, Lewis acid mediated reductive etherification of the originally formed mixture of alcohol 10 and its hemiacetal 11 using triethylsilane as a reducing agent was performed, and enabled stereoselective construction of the tetrahydropyrane ring in 16 (Scheme 5). Importantly, both developed reaction sequences consisting of either two or three steps could be performed without purification of the intermediates, thereby underlining the high practicality of the developed synthetic strategies.

The absolute configuration of the products was unambiguously assigned by using single-crystal X-ray analysis of the corresponding Ramirez olefination products derived from **3c** and **3j** (Scheme 6a).^[10] The absolute configuration of the remaining products **3a**, **b**, **d**–**i**, **k**, **l** was assigned by analogy. The relative configuration of the newly introduced stereogenic centers in products **9**, **15**, and **16** was assigned by 1D

www.angewandte.org





Scheme 5. Introduction of a tetrahydropyrane ring into tetrahydroxanthone **3 a**.



Scheme 6. Configurational assignments and rationalization for the stereochemical outcome of the H-bond-directed trienamine-mediated cycloadditions. a) Crystal structures of the products of the Ramirez olefination of **3 c** (right) and **3 j** (left). Blue N; red O, white H; gray C, brown Br. b) Transition state models for the H-bond-directed trienamine-mediated cycloadditions and the corresponding products.

NOE experiments. Based on the configurational assignments, transition state models rationalizing the observed absolute and relative stereochemistry of the products **3** were proposed (Scheme 6b). It is postulated that the 3-cyanochromone interacts with the squaramide moiety of the catalyst through the electron-rich cyano group^[11] rather than the carbonyl group. Secondary orbital overlap with participation of the chromone carbonyl group as well as π -stacking interactions between the aromatic ring of the chromone system and the conjugated π system of the corresponding trienamine might further support such an alignment of the dienophile. Importantly, the reaction pathway strongly depends on the substituent in the 4-position of the starting aldehyde **1**, the reaction is proposed to proceed through transition state **A**.

Accordingly, it is assumed that the corresponding trienamine intermediate reacts in the s-cis-(C3=C4-C5=C6) and s-cis-(C1=C2-C3=C4) conformation and the reaction occurs across the remote diene system. However, for steric reasons, the presence of a substituent in the 4-position of 1 forces the reaction to proceed through transition state **B**. In this case, the sterically demanding R¹ substituent positions itself away from the first double bond of the trienamine system as well as from the C2 substituent of the pyrrolidine catalyst. This reactive conformation of the diene can account for the unprecedented trans relationship between the C1 and C4 stereogenic centers of 3. In both cases, proper alignment of the reactive diene system with respect to the chromone dienophile is postulated to be the crucial factor determining the stereochemistry of the product. Nonetheless, the obtained results suggest that the mechanism(s) of trienamine-mediated reactions can be a rather complex issue and might be highly dependent on the structure of the starting dienal. Computational studies in an attempt to understand H-bond-directed trienamine-mediated reactions are currently in progress.

In summary, the first H-bond-directed trienamine-mediated [4+2] cycloaddition was developed, thereby demonstrating the viability of such an activation strategy. The reaction between diversely substituted 2,4-dienals and 3-cyanochromones proceeded smoothly and in a highly stereoselective manner in the presence of a squaramide-containing aminocatalyst. Furthermore, it was demonstrated that the obtained cycloadducts can be chemo- and diastereoselectively transformed into polycyclic products with high molecular and stereochemical complexity that possess up to five stereogenic centers. An unexpected stereochemical outcome of the reaction was observed and a rationalization of the results was provided.

Received: June 19, 2012 Published online: ■■ ■■, ■■■■

Keywords: asymmetric synthesis \cdot organocatalysis \cdot synthetic methods \cdot tetrahydroxanthones \cdot trienamines

- [2] For selected recent examples, see: a) K. Tatsuta, S. Hosokawa, *Chem. Rev.* 2005, 105, 4707; b) S. Sezer, D. Özdemirhan, E. Şahin, C. Tanyeli, *Tetrahedron: Asymmetry* 2006, 17, 2981; c) K. Liu, A. Chougnet, W.-D. Woggon, *Angew. Chem.* 2008, 120, 5911; *Angew. Chem. Int. Ed.* 2008, 47, 5827; d) C. A. Lewis, J. L. Gustafson, A. Chiu, J. Balsells, D. Pollard, J. Murry, R. A. Reamer, K. B. Hansen, S. J. Miller, *J. Am. Chem. Soc.* 2008, 130, 16358.
- [3] For a recent review on dienamine chemistry, see: a) D. B. Ramachary, Y. V. Reddy, *Eur. J. Org. Chem.* 2012, 865; for selected examples, see: b) S. Bertelsen, M. Marigo, S. Brandes, P. Dinér, K. A. Jørgensen, *J. Am. Chem. Soc.* 2006, *128*, 12973; c) G. Bergonzini, S. Vera, P. Melchiorre, *Angew. Chem.* 2010, *122*, 9879; *Angew. Chem. Int. Ed.* 2010, *49*, 9685; d) J. Stiller, E. Marqués-López, R. P. Herrera, R. Fröhlich, C. Strohmann, M.

www.angewandte.org

© 2012 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

These are not the final page numbers!

For selected reviews, see: a) H. Sailes, A. Whiting, J. Chem. Soc. Perkin Trans. 1 2000, 1785; b) K. Mikami, M. Shimizu, H.-C. Zhang, B. E. Maryanoff, Tetrahedron 2001, 57, 2917; c) J. Clayden, N. Vassiliou, Org. Biomol. Chem. 2006, 4, 2667; d) E. J. Thomas, Chem. Rec. 2007, 7, 115; e) J. Clayden, Chem. Soc. Rev. 2009, 38, 817.

Christmann, Org. Lett. 2011, 13, 70; e) J.-L. Li, S.-L. Zhou, P.-X. Chen, L. Dong, T.-Y. Liu, Y.-C. Chen, Chem. Sci. 2012, 3, 1879; f) Ł. Albrecht, G. Dickmeiss, F. Cruz Acosta, C. Rodríguez-Escrich, R. L. Davis, K. A. Jørgensen, J. Am. Chem. Soc. 2012, 134, 2543; g) G. Talavera, E. Reyes, J. L. Vicario, L. Carrillo, Angew. Chem. 2012, 124, 4180; Angew. Chem. Int. Ed. 2012, 51, 4104.

- [4] For a highlight on trienamine chemistry, see: a) E. Arceo, P. Melchiorre, Angew. Chem. 2012, 124, 5384; Angew. Chem. Int. Ed. 2012, 51, 5290; for examples, see b) Z.-J. Jia, H. Jiang, J.-L. Li, B. Gschwend, Q.-Z. Li, X. Yin, J. Grouleff, Y.-C. Chen, K. A. Jørgensen, J. Am. Chem. Soc. 2011, 133, 5053; c) H. Jiang, B. Gschwend, Ł. Albrecht, S. G. Hansen, K. A. Jørgensen, Chem. Eur. J. 2011, 17, 9032; d) Z.-J. Jia, Q. Zhou, Q.-Q. Zhou, P.-Q. Chen, Y.-C. Chen, Angew. Chem. 2011, 123, 8797; Angew. Chem. Int. Ed. 2011, 50, 8638; e) Y.-K. Liu, M. Nappi, E. Arceo, S. Vera, P. Melchiorre, J. Am. Chem. Soc. 2011, 133, 15212; f) Y. Liu, M. Nappi, E. C. Escudero-Adán, P. Melchiorre, Org. Lett. 2012, 14, 1310.
- [5] For general reviews on applications of steric shielding aminocatalysts in aminocatalysis, see: a) G. Lelais, D. W. C. MacMillan, Aldrichimica Acta 2006, 39, 79; b) A. Mielgo, C. Palomo, *Chem. Asian J.* 2008, 3, 922; c) L.-W. Xu, L. Li, Z.-H. Shi, Adv. Synth. Catal. 2010, 352, 243; d) L. Z. Zimmer, C. Sparr, R. Gilmaur, Angew. Chem. 2011, 123, 12062; Angew. Chem. Int. Ed. 2011, 50, 11860; e) K. L. Jensen, G. Dickmeiss, H. Jiang, Ł. Albrecht, K. A. Jørgensen, Acc. Chem. Res. 2012, 45, 248.

- [6] a) K.-S. Masters, S. Bräse, *Chem. Rev.* 2012, *112*, 3717; b) R. N. Kharwar, A. Mishra, S. K. Gond, A. Stierle, D. Stierle, *Nat. Prod. Rep.* 2011, *28*, 1208.
- [7] a) R. Andersen, G. Buechi, B. Kobbe, A. L. Demain, J. Org. Chem. 1977, 42, 352; b) K. C. Nicolaou, A. Li, Angew. Chem. 2008, 120, 6681; Angew. Chem. Int. Ed. 2008, 47, 6579; c) M. E. Maier, Nat. Prod. Rep. 2009, 26, 1105.
- [8] For examples of the application of 3-cyanochromones in Diels-Alder reaction, see: a) R. P. Hsung, *J. Org. Chem.* 1997, 62, 7904;
 b) S. J. Degen, H. C. Shen, G. M. Golding, C. A. Zificsak, R. P. Hsung, *Bioorg. Med. Chem. Lett.* 1999, *9*, 973.
- [9] For recent reviews on H-bond-directed catalysis, see: a) R. I. Storer, C. Aciro, L. H. Jones, *Chem. Soc. Rev.* 2011, 40, 2330;
 b) J. Alemán, A. Parra, H. Jiang, K. A. Jørgensen, *Chem. Eur. J.* 2011, 17, 6890; for the seminal report on squaramide-based catalysts in enantioselective synthesis, see: c) J. P. Malerich, K. Hagihara, V. H. Rawal, *J. Am. Chem. Soc.* 2008, 130, 14416.
- [10] See the Supporting Information for the crystal structures. CCDC 886343 and 886348 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [11] a) S. J. Zuend, M. P. Coughlin, M. P. Lalonde, E. N. Jacobsen, *Nature* 2009, 461, 968; for a review on H-bond strengths, see: b) T. Steiner, *Angew. Chem.* 2002, 114, 50; *Angew. Chem. Int. Ed.* 2002, 41, 48.



Communications



Asymmetric Catalysis

Ł. Albrecht, F. Cruz Acosta, A. Fraile, A. Albrecht, J. Christensen, K. A. Jørgensen* _____ ####-#####

Enantioselective H-Bond-Directing Approach for Trienamine-mediated Reactions in Asymmetric Synthesis





and activated chromones. It is possibile to control the stereochemical outcome of such reactions by employing an H-bonddirecting aminocatalyst.

Biologically relevant tetrahydroxanthones

H-bond donor

H-bond-directed catalysis