This article was downloaded by: [Mount Allison University OLibraries] On: 02 May 2013, At: 01:48 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

http://www.tandfonline.com/loi/gpss20

# Synthesis and Isolation of $\alpha$ -Oxo Sulfines

Theodorus J. G. Damen & Binne Zwanenburg

<sup>a</sup> Department of Organic Chemistry, NSR-Center for Molecular Structure, Design and Synthesis. University of Nijmegen, Toernooiveld, 6525ED Nijmegen, The Netherlands

<sup>b</sup> Department of Organic Chemistry, NSR-Center for Molecular Structure, Design and Synthesis. University of Nijmegen, Toernooiveld, 6525ED Nijmegen, The Netherlands

Published online: 30 Jul 2008.

To cite this article: Theodorus J. G. Damen & Binne Zwanenburg (1997): Synthesis and Isolation of  $\alpha$ -Oxo Sulfines, Phosphorus, Sulfur, and Silicon and the Related Elements, 120:1, 319-320

To link to this article: <u>http://dx.doi.org/10.1080/10426509708545528</u>

### PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <u>http://www.tandfonline.com/page/terms-and-conditions</u>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sublicensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

#### Synthesis and Isolation of α-Oxo Sulfines

THEODORUS J. G. DAMEN and BINNE ZWANENBURG<sup>\*</sup> Department of Organic Chemistry, NSR-Center for Molecular Structure, Design and Synthesis. University of Nijmegen, Toernooiveld, 6525ED Nijmegen, The Netherlands.

The synthesis of several  $\alpha$ -oxo sulfines is discribed. Various  $\alpha$ -oxo sulfines can be isolated as such. The Diels-Alder reaction of these sulfines with 1,3-dienes can be catalyzed by Lewis acids.

KEYWORDS: sulfine, Diels-Alder reaction, Lewis acid catalysis, thiopyran (S-oxide)

#### INTRODUCTION

In recent years our efforts in the area of sulfine chemistry<sup>1,2,3</sup> are particularly focused on the synthetic elaboration of  $\alpha$ -oxo sulfines.<sup>4,5</sup> This type of sulfines can readily be obtained by reaction of either doubly activated methylene compounds, silyl ketene acetals or silyl enol ethers with thionyl chloride in the presence of a suitable tertiary base. Thus far,  $\alpha$ -oxo sulfines have been trapped as their cycloadducts of 1,3-dienes because of their suspected susceptibility toward hydrolysis. However, by a proper modification of the experimental procedures a variety of  $\alpha$ -oxo sulfines can be isolated as such. This finding allows a more detailed study of these sulfines with various reagents.

#### RESULTS

A wide range of differently substituted  $\alpha$ -oxo sulfines has been synthesized and isolated in high yields. Representative examples are collected in Table I. The starting compound can either be an ester (entries 1-4), a ketone (entry 5), an aldehyde (entry 6) or a doubly activated methylene compound (entries 7-8). The last mentioned compound is sufficiently enolized to react directly with thionyl chloride in the presence of base to give the  $\alpha$ -oxo sulfine. In all other cases, the starting carbonyl compound first had to be converted into the corresponding enol silyl ethers. The thermodynamically more stable geometrical isomer, with the sulfine oxygen *trans* to the carbonyl, was formed exclusively.

	R <sup>2</sup> base Me <sub>3</sub> SiCl	R <sup>1</sup> <sup>APA</sup> OSiMe <sub>3</sub> base / SOCl <sub>2</sub>	SOCI <sub>2</sub>	$R^1$ $R^2$
entry	R <sup>1</sup>	R <sup>2</sup>	base	yield (%) <sup>a</sup>
1	Me	OMe	DIPEA	64 <sup>b</sup>
2	i-Pr	OMe	DIPEA	89
3	t-Bu	OMe	DIPEA	85
4	Ph	OMe	DIPEA	94
5	Me	Ph	2,6-Lutidine	92
6	PhCH <sub>2</sub>	Н	2,6-Lutidine	80
7	MeOOC	OMe	Et <sub>3</sub> N	98
8	NC	OEt	Et <sub>3</sub> N	78

TABLE Ι Synthesis of α-oxo sulfines

<sup>a</sup> yield of crude product unless mentioned otherwise; <sup>b</sup> yield after distillation

The Diels-Alder reaction of  $\alpha$ -oxo sulfines with 1,3-dienes conveniently leads to dihydrothiopyran S-oxides which are very interesting synthons for the preparation of a wide range of sulfur-containing heterocycles, including thiosugar derivatives. The Diels-Alder reaction of  $\alpha$ -oxo sulfines with less reactive 1,3-dienes can be catalyzed by a catalytic amount of tin(IV) chloride. This is the first example of Lewis catalysis in the cycloaddition of sulfines. The synthetic prospects of these cycloadducts are currently under active investigation.

#### REFERENCES

- 1. B. Zwanenburg, Recl. Trav. Chim. Pays-Bas 101, 1 (1982)
- B. Zwanenburg and B. G. Lenz, Houben-Weyl, Methoden der Organische Chemie, Band E11, p. 911, George Thieme Verlag (1985)
- 3. B. Zwanenburg, Phosphorus, Sulfur and Silicon 43, 1 (1989)
- 4. J.B. Rewinkel, Thesis University of Nijmegen (1989)
- 5. R.C. de Laet, Thesis University of Nijmegen (1995)