

ChemComm

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: K. Komeyama, S. Kiguchi and K. Takaki, *Chem. Commun.*, 2016, DOI: 10.1039/C6CC03086F.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



COMMUNICATION

A drastic effect of cobalt and chromium catalysts in the borylation of arylzinc reagents†

Kimihiro Komeyama,* Shinnosuke Kiguchi and Ken Takaki

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/chemcomm

A new synthetic approach to arylboronic esters from arylzinc reagents with boryl electrophiles MeOB(OR)_2 has been developed. Furthermore, this protocol could be applied to the cyclization/borylation of alkynylaryl iodides to afford cyclized vinylboronic esters.


Arylboronic esters are recognized as being indispensable building blocks in organic synthesis because of their stability and wide ranging applicability to C–X (X = C, N and O) bond-forming reactions in chemical, medicinal and material science.¹ Over the past 20 years, transition-metal-catalysed borylations of aryl (pseudo)halides with diborons, hydroboranes or metal-boryl reagents have been developed using Pd, Rh, Ni, Cu, Zn, Fe and Co complexes.² These methods facilitate the concise synthesis of arylboronic esters bearing useful functional groups (route a, Scheme 1). More recently, the synthetic strategies used to produce arylboronic esters have shifted to favour direct C–H borylation of arenes, which has been rapidly evolved by precious metal catalysts such as Rh and Ir (route b).^{3,4} Moreover, very recently, the direct borylations have been also accomplished by abundant and inexpensive transition-metal catalysts.⁵ In contrast, transition-metal-free borylation protocols, such as electrophilic borylations of electron-rich arenes (route c),⁶ the Sandmeyer-type borylation (route d),⁷ and boryl substitution reactions with borylsilanes⁸ or borylzincates⁹ (route e), have been frequently reported, although there are still problems with these strategies, such as their narrow substrate scope, loss of the boron component and/or the scarcity of suitable boron sources.

A classical but powerful method for synthesis of arylboronic esters is the substitution of aryllithium or arylmagnesium reagents with trialkylboric esters.¹⁰ This method, however, suffers from poor functional group tolerance. Conversely, compared with the above organometallics, organozinc compounds are highly compatible with a broad range of polar functional groups due to the relatively weak ionic character of the C–Zn bond, which may

undergo chemoselective transformation.¹¹ However, the nucleophilicity of organozinc reagents is quite low; therefore, their reactions with organic electrophiles often require the use of transition-metal catalysts.¹² Despite their high potential for the development of tractable organic transformations, the borylation of organozinc reagents using easy-to-handle boric esters has not been reported,¹³ except for highly electrophilic B-chlorocatecholborane and subporphyrins.¹⁴

Transmetalation is a highly effective way to drastically change the reactivity of an organometallic species. For example, Takagi reported that highly nucleophilic arylchromium intermediates,¹⁵ afforded by the reaction of arylzinc reagents with chromium(III) salts, underwent addition to aldehydes under mild conditions. A similar strategy of changing the nucleophilicity of organometallics via transmetalation has been demonstrated in the Nozaki–Hiyama–Kishi reaction.¹⁶ Here we report the marked additive effect of low-valence cobalt and chromium catalysts in the borylation of arylzinc reagents with MeOB(OR)_2 as the boryl electrophile in the presence of TMSCl (route f). Furthermore, this protocol could be applied to the direct synthesis of arylboronic esters from ubiquitous aryl halides under CoBr_2 , xantphos and $\text{CrCl}_3(\text{thf})_3$ catalyst systems in the presence of Zn.

We began by examining the borylation of the preformed 4-




Scheme 1 Synthetic methods for arylboronic esters.

Department of Applied Chemistry, Graduate School of Engineering, Hiroshima University, Higashi-Hiroshima 739-8527, Japan.

E-mail: kkome@hiroshima-u.ac.jp; Fax: +81-82-424-5494; Tel: +81-82-424-7747

† Electronic Supplementary Information (ESI) available: Additional experimental results, experimental procedures and characterisation for products. See DOI: 10.1039/x0xx00000x


Table 1 Effect of cobalt and chromium salts in the borylation of 4-tolylZnI-LiCl with MeOBpin (**2a**).^a


Entry	Additives (mol %)	Zn (x equiv.)	Yield (%) ^b
1	–	–	Trace
2	CrCl ₃ (thf) ₃ (20)	–	Trace
3	CrCl ₂ (20)	–	Trace
4	CoBr ₂ /xantphos (10), CrCl ₃ (thf) ₃ (20)	2.0	73
5	CoBr ₂ /xantphos (10), CrCl ₃ (thf) ₃ (20)	0.5	67
6	CoBr ₂ /xantphos (10), CrCl ₃ (thf) ₃ (20)	–	Trace
7	CrCl ₃ (thf) ₃ (20)	2.0	Trace
8	CoBr ₂ /xantphos (10)	2.0	32
9	CoBr/xantphos (10) ^c , CrCl ₃ (thf) ₃ (20)	–	13
10	CoBr/xantphos (10) ^c , CrCl ₂ (20)	–	65
11	CoBr (10) ^c	–	trace
12	CoBr/dppe (10) ^c , CrCl ₂ (20)	–	53

^aAll reactions were carried out in the presence of TMSCl (1.2 equiv.). ^bNMR yield. ^cThe borylation was performed after reduction with 20 mol % zinc powder to form the low-valence cobalt complex.

MeC₆H₄ZnI-LiCl¹⁷ with 2-methoxy-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (**2a**, MeOBpin) as model substrates in the presence of TMSCl (1.2 equiv.) at 60 °C for 16 h. These results are summarized in Table 1. As expected, the arylzinc compound was inactive and hardly participates in the borylation, wherein the arylzinc remained unchanged after the reaction (Table 1, entry 1).¹⁸ The addition of CrCl₃(thf)₃ or CrCl₂ catalyst (20 mol %) did not afford the arylboronic ester (entries 2 and 3). Conversely, we observed an unexpected effect arising from the combination of cobalt and chromium catalysts under reduction conditions (entries 4 and 5). Thus, a mixture of CoBr₂/xantphos¹⁹ and CrCl₃(thf)₃ complexes in the presence of zinc powder (0.5–2.0 equiv.) effectively promoted the borylation, giving rise to 4-tolylboronic pinacol ester (**3aa**) in satisfactory yields (67%–73%). Intriguingly, all the components, i.e. the cobalt, chromium and zinc reductant, are crucial for efficient borylation (entries 6–8). Finally, we found that low-valence cobalt and chromium complexes play important roles in the borylation reaction (entries 9 and 10). The role of the xantphos ligand is unclear,²⁰ and it is not absolutely necessary to the borylation. There is no reaction in the absence of the ligand (entry 11); however, the presence of dppe afforded **3aa** in 53% yield (entry 12).

The cobalt/xantphos complex has been reported to be an equally effective catalyst for the transformation of aryl halides into arylzinc reagents under similar reduction conditions.²¹ Based on these reports and our previous work on cobalt-catalysed reactions,²² the present Co/Cr-catalysed borylation with arylzinc reagents was refined into a more simplified protocol starting from 4-tolyl iodide (Table 2, entry 1). Without TMSCl, the desired boronic ester **3aa** was obtained in low yield (27%), even with a longer reaction time (entry 2). This result indicates robust Cr–O bond formation during the reaction, because the bond appears to be difficult to cleave in the catalytic cycle without the aid of TMSCl.²³ Chromium

Table 2 Screening of the reaction conditions for Co/Cr-catalysed borylation starting from 4-tolyl bromide halides **1a** with **2a**.^a


Entry	Change from standard conditions	Yield/% of 3aa ^a
1	–	79 (75)
2 ^b	without TMSCl	27
3	CrCl ₃ (bpy) in place of CrCl ₃ (THF) ₃	12
4	CrCl ₃ (1,10-phen) in place of CrCl ₃ (THF) ₃	4
5 ^c	1,10-phen in place of xantphos	60
















^aNMR yield. The parenthesis value indicates an isolated yield. ^bReaction time: 48 h. ^cMeCN solvent was used instead of THF.

complexes ligated by nitrogen ligands,²⁴ such as 2,2'-bipyridyl (bpy) and 1,10-phenanthroline (1,10-phen), inhibit the borylation in THF solvent (entries 6 and 7). However, the borylation proceeded in acetonitrile with the replacement of xantphos with 1,10-phen, albeit with a slightly lower yield (entry 5).²⁵

Having established the optimum conditions (Table 2, entry 1), we explored substrate scope through the Co/Cr-catalysed borylation of various aryl bromides (Table 3). Electron-neutral (**1b** and **1c**) and -rich aryl bromides (**1d**, **1e** and **1f**) are efficiently converted to the corresponding arylboronic acid pinacol esters **3** in 66%–71% yields (entries 1–5). However, electron-deficient aryl halides with inductively and/or resonance withdrawing substituents were less reactive (entries 6, 8, 10, 12, 14 and 17). Especially, the strongly electron-deficient aryl bromides bearing CN (**1j**) and CF₃ (**1k**, **1m**) substituents were markedly less reactive towards trapping with **2a** (entries 12, 14 and 17). Fortunately, significant improvement was achieved by the replacement of **2a** with 2-methoxy-5,5-dimethyl-1,3,2-dioxaborinane (MeOBnep, **2b**), which is a less sterically hindered electrophile than **2a**.²⁶ This boronate preferentially afforded the desired boronic esters **3jb**, **3kb** and **3mb** in good yields (entries 13, 15 and 18). We also explored the effect of the leaving group, such as the 2-methoxyethoxy substituent, on the boron **2**, but such substituents did not significantly affect the reactivity.²⁷ In the Co/Cr-catalysed borylation, trimethylsilyl aryls (**1l**, entry 16) as well as heteroaryl skeletons such as indole (**1o**, entry 21) and thiophene (**1p**, entry 22) also took part in the borylation. In contrast, the use of aryl chlorides leads to lower product yields.

The borylation reaction presumably involves the generation of aryl cobalt and aryl chromium intermediates. In contrast, the migratory insertion of an alkyne into both aryl-metal complexes has been established.^{25b,28} Based on these findings and our previous results,²² we assumed that if these aryl-metal complexes are active in alkyne-insertion reactions to form a vinyl-metal species, and the products undergo a substitution reaction with boryl electrophiles under identical conditions to those presented above, a useful three-component reaction (carbaboration) could be developed. The carboboration of alkynes is a very important process for regio-

Table 3 Substrate scope for Co/Cr-catalysed borylation of aryl halides.

Ar—X + MeO[B]		10 mol % CoBr ₂ 10 mol % xantphos 20 mol % CrCl ₃ (thf) ₃ Zn (2.0 equiv.) LiCl (1.0 equiv.) TMSCl (1.2 equiv.) THF, 60 °C		Ar—[B]	
1 X = Br, I [B] = Bpin: 2a = Bnep: 2b				3	
Entry	Ar—Br 1	Time /h	Product & Yield /% ^a		
1		16	3ba 71		
2		16	3ca 66		
3		16	3da 66		
4		16	3ea 67		
5 ^b		48	3fa 68		
6		16	3ga 38		
7 ^c		16	3gb 66		
8		16	3ha 62		
9 ^c		16	3hb 75		
10		48	3ia 68		
11 ^c		48	3ib 74		
12 ^b		48	3ja 38		
13 ^{b,c}		48	3jb 83		
14 ^b		48	3ka 23		
15 ^{b,c}		48	3kb 83		
16 ^c		24	3lb 92		
17		48	3ma trace		
18 ^{b,c}		48	3mb 84		
19 ^b		48	3na 65		
20 ^{b,c}		48	3nb 66		
21		48	3oa 72		
22 ^c		48	3pb 80		

^a Isolated yield. ^b 2 equivalent of boryl electrophile was employed. ^c MeOB(nep) was used instead of MeOB(pin).


and stereoselective synthesis of multisubstituted olefins.²⁹

Similar catalytic reactions for alkynes^{13,30} have been reported; however, all examples employed carbon electrophiles. The transformation using boryl electrophiles has been never demonstrated, except for cases employing Grignard reagents.³¹ Our initial attempts at the addition/borylation of 4-octyne with 4-tolyl bromide **1a** and MeOBpin **2a** under identical conditions failed, wherein the major product is arylboronic ester **3aa**. This may be because the reaction of the *in situ* generated aryls-metal species with the boryl electrophile is fast compared to the intermolecular reaction with the alkyne. Based on these experimental results, we next attempted the cyclisation/borylation of alkynyl aryl iodides **4** (Scheme 2). Having refined the conditions,³² the desired cyclisation/borylation was accomplished, affording the cyclised vinyl boronic esters **5a**, **5b** and **5c** in 50, 41 and 43% yields, respectively (Scheme 2). The stereochemistry of the products was determined by NOE measurements, which clearly indicated *syn*-carboboration. In addition, an alkynyl aryl iodide tethered with nitrogen also reacted to afford the corresponding boronic ester **5d** in 26% yield.

Although the actual role of the cobalt catalyst is still unclear, the arylchromium(II) species seems to trigger the borylation step. Thus, the reaction of 4-MeC₆H₄CrCl₂ (prepared via the reaction of the 4-MeC₆H₄Li with CrCl₃(thf)₃)³³ with MeOBpin afforded **3aa** in only 3% GC yield.³⁴ In contrast, the reaction of the 4-MeC₆H₄CrCl provided **3aa** in 67% yield.

In conclusion, we have reported a drastic additive effect for increasing the reactivity of arylzinc reagents, enabling the borylation of unreactive arylzinc compounds to afford various aryl boronic esters. In addition, we found that this protocol could be applied to a more practical borylation starting from ubiquitous aryl halides. The easy-to-operate procedure avoids preparation of air- and moisture-sensitive arylzinc reagents, making it more practical for arylboronic ester synthesis. Furthermore, cyclisation/borylation of arylalkynyl iodides was accomplished by the modified catalytic system to give cyclised vinyl boronic esters, in which carboboration proceeded in an exclusively *syn*-addition manner. Control experiments revealed the important roles of low-valence cobalt and chromium in the borylation. This suggests that a key intermediate is an aryl chromium(II) species, although more data are required in order to understand the mechanistic details of the reaction. Further studies on the mechanism and synthetic applications of the Co/Cr catalyst system are currently underway in our laboratory.

This work was partially supported by a Grant-Aid for Scientific Research (KAKENHI) from the Ministry of Education, Culture, Sports, Science and Technology of Japan (No.

**Scheme 2** Substrate scope for the Co/Cr-catalysed cyclisation/borylation.^a

15K05502).

Notes and references


- (a) N. Miyaura, A. Suzuki, *A. Chem. Rev.* 1995, **95**, 2457. (b) D. G. Hall, *Boronic Acids: Preparation and Applications in Organic Synthesis and Medicine*, Wiley-VCH, Weinheim, 2005.
- Recent reviews for transition metal-catalyzed borylation of aryl (pseudo)halides: (a) W. K. Chow, O. Y. Yuen, P. Y. Choy, C. M. So, C. P. Lau, W. T. Wong, F. Y. Kwong, *RSC Adv.* **2013**, **3**, 12518. (b) M. Murata, *Heterocycles*, 2012, **85**, 1795. With Pd: (c) J. Campos and S. Aldridge, *Angew. Chem. Int. Ed.*, 2015, **54**, 14159. With Rh: (d) M. Tobisu, H. Kinuta, Y. Kita, E. Rémond and N. Chatani, *J. Am. Chem. Soc.*, 2011, **134**, 115. (e) H. Kinuta, M. Tobisu and N. Chatani, *J. Am. Chem. Soc.*, 2015, **137**, 1593. With Ni: (f) M. Tobisu, K. Nakamura and N. Chatani, *J. Am. Chem. Soc.*, 2014, **136**, 5587. (g) X.-W. Liu, J. Echavarren, C. Zarate and R. Martin, *J. Am. Chem. Soc.*, 2015, **137**, 12470. (h) T. Niwa, H. Ochiai, Y. Watanabe and T. Hosoya, *J. Am. Chem. Soc.*, 2015, **137**, 14313. (i) J. Zhou, M. W. Kuntze-Fechner, R. Bertermann, U. S. D. Paul, J. H. J. Berthel, A. Friedrich, Z. Du, T. B. Marder and U. Radius, *J. Am. Chem. Soc.*, 2016, DOI: 10.1021/jacs.6b02337. With Cu: C. Kleeberg, L. Dang, Z. Lin and T. B. Marder, *Angew. Chem. Int. Ed.*, 2009, **48**, 5350. With Zn: (j) S. Bose and T. B. Marder, *Org. Lett.*, 2014, **16**, 4562. (k) S. K. Bose, A. Deisenberger, A. Eichhorn, P. G. Steel, Z. Lin and T. B. Marder, *Angew. Chem. Int. Ed.*, 2015, **54**, 11843. With Fe: (l) R. B. Bedford, P. B. Brenner, E. Carter, T. Gallagher, D. M. Murphy and D. R. Pye, *Organometallics*, 2014, **33**, 5940. (m) F. Labre, Y. Gimbert, P. Bannwarth, S. Olivero, E. Duñach and P. Y. Chavant, *Org. Lett.*, 2014, **16**, 2366. With Co: (n) W. Yao, H. Fang, S. Peng, H. Wen, L. Zhang, A. Hu and Z. Huang, *Organometallics*, 2016, DOI: 10.1021/acs.organomet.6b00161.
- Review for Rh or Ir catalysed aromatic C-H borylation: (a) T. Ishiyama and N. Miyaura, *J. Orgmet. Chem.*, 2003, **680**, 3. (b) I. A. I. Mkhaliid, J. H. Barnard, T. B. Marder, J. M. Murphy and J. F. Hartwig, *Chem. Rev.*, 2010, **110**, 890. (c) J. F. Hartwig, *Chem. Soc. Rev.*, 2011, **40**, 1992. (d) A. Ros, R. Fernández and J. M. Lassaletta, *Chem. Soc. Rev.*, 2014, **43**, 3229.
- (a) S. M. Preshlock, D. L. Plattner, P. E. Maligres, S. W. Krska, R. E. Maleczka and M. R. Smith, *Angew. Chem. Int. Ed.*, 2013, **52**, 12915. (b) S. Kawamori, T. Miyazaki, H. Ohmiya, T. Iwai and M. Sawamura, *J. Am. Chem. Soc.*, 2011, **133**, 19310. (c) Y. Saito, Y. Segawa and K. Itami, *J. Am. Chem. Soc.*, 2015, **137**, 5193. (d) Y. Kuninobu, H. Ida, M. Nishi and M. Kanai, *Nat. Chem.*, 2015, **7**, 712. With Pt: (e) T. Furukawa, M. Tobisu and N. Chatani, *J. Am. Chem. Soc.*, 2015, **137**, 12211.
- With Ni: (a) T. Furukawa, M. Tobisu and N. Chatani, *Chem. Commun.*, 2015, **51**, 6508. (b) H. Zhang, S. Hagihara and K. Itami, *Chem. Lett.*, 2015, **44**, 779. With Fe: (c) T. Hatanaka, Y. Ohki and K. Tatsumi, *Chem. Asian J.*, 2010, **5**, 1657. (d) G. Yan, Y. Jiang, C. Kuang, S. Wang, H. Liu, Y. Zhangb and J. Wang, *Chem. Commun.*, 2010, **46**, 3170. (e) T. Dombray, C. G. Werncke, S. Jiang, M. Grellier, L. Vendier, S. Bontemps, J.-B. Sortais, S. Sabo-Etienne and C. Darcel, *J. Am. Chem. Soc.*, 2015, **137**, 4062. With Fe/Cu: (f) T. J. Mazzacano and N. P. Mankad, *J. Am. Chem. Soc.*, 2013, **135**, 17258. With Co: (g) J. V. Obligation, S. P. Semproni and P. J. Chirik, *J. Am. Chem. Soc.*, 2014, **136**, 4133.
- A. Del Grosso, M. D. Helm, S. A. Solomon, D. Caras-Quintero and M. J. Ingleson, *Chem. Commun.*, 2011, **47**, 12459.
- F. Mo, Y. Jiang and D. Qiu, Y. Zhang and J. Wang, *Angew. Chem. Int. Ed.*, 2010, **49**, 1846.
- E. Yamamoto, K. Izumi, Y. Horita and H. Ito, *J. Am. Chem. Soc.*, 2012, **134**, 19997.
- Y. Nagashima, R. Takita, K. Yoshida, K. Hirano and M. Uchiyama, *J. Am. Chem. Soc.*, 2013, **135**, 18730.
- (a) H. C. Brown and T. E. Cole, *Organometallics*, 1983, **2**, 1316. (b) J. W. Clary, T. J. Rettenmaier, R. Snelling, W. Bryks, J. Banwell, W. T. Wipke and B. Singaram, *J. Org. Chem.*, 2011, **76**, 9602.
- P. Knochel and R. D. Singer, *Chem. Rev.*, 1993, **93**, 2117.
- P. Knochel, M. A. Schade, S. Bernhardt, G. Manolikakes, A. Metzger, F. M. Piller, C. J. Rohbognier and M. Mosrin, *Beilstein J. Org. Chem.*, 2011, **7**, 1261.
- N. Nakagawa, T. Hatakeyama and M. Nakamura, *Chem. Eur. J.*, 2015, **21**, 4257.
- (a) S. Claudel, C. Gosmini, J. M. Paris and J. Périchon, *Chem. Commun.*, 2007, 3667. (b) R. Kotani, K. Yoshida, E. Tsurumaki and A. Osuka, *Chem. Eur. J.*, 2016, **22**, 3320.
- (a) Y. Ogawa, M. Mori, A. Saiga and K. Takagi, *Chem. Lett.*, 1996, 1069. (b) Y. Ogawa, A. Saiga, M. Mori, T. Shibata and K. Takagi, *J. Org. Chem.*, 2000, **65**, 1031.
- (a) K. Takai, K. Nitta and O. Fujimura, *J. Org. Chem.*, 1989, **54**, 4732. (b) K. Takai, N. Matsukawa, A. Takahashi and T. Fujii, *Angew. Chem. Int. Ed.*, 1998, **37**, 152.
- A. Krasovskiy, V. Malakhov, A. Gavryushin and P. Knochel, *Angew. Chem. Int. Ed.*, 2006, **45**, 6040.
- After the reaction, treatment of the reaction mixture with I₂ afforded 4-iodotoluene in nearly quantitative yield.
- M. M. P. Grutters, C. Müller and D. Vogt, *J. Am. Chem. Soc.*, 2006, **128**, 7414.
- M.-N. Birkholz née Gensow, Z. Freixa and P. W. N. M. van Leeuwen, *Chem. Soc. Rev.*, 2009, **38**, 1099.
- M.-Y. Jin and N. Yoshikai, *J. Org. Chem.*, 2011, **76**, 1972.
- (a) K. Komeama, T. Kashihara and K. Takaki, *Tetrahedron Lett.*, 2013, **54**, 5659. (b) K. Komeyama, Y. Okamoto and K. Takaki, *Angew. Chem. Int. Ed.*, 2014, **53**, 11325. (c) R. Asakura, H. Fukuoka and K. Takaki, *Tetrahedron Lett.*, 2015, **56**, 1735.
- F. Alois and N. Shi, *J. Am. Chem. Soc.*, 1996, **118**, 12349.
- K. Namba and Y. Kishi, *J. Am. Chem. Soc.*, 2005, **127**, 15382.
- (a) H. Fillon, C. Gosmini and J. Périchon, *J. Am. Chem. Soc.*, 2003, **125**, 3867. (b) M. Corpet and C. Gosmini, *Chem. Commun.*, 2012, **48**, 11561.
- See Ref. 13 and references therein.
- See Table S1 in electronic supplementary information (ESI).
- With Co catalyst: (a) K. Murakami, H. Yorimitsu and K. Oshima, *Org. Lett.*, 2009, **11**, 2373. (b) B.-H. Tan, J. Dong and N. Yoshikai, *Angew. Chem. Int. Ed.*, 2012, **51**, 9610. With Cr catalyst: (c) K.-I. Kanno, Y. Liu, A. Iesato, K. Nakajima and T. Takahashi, *Org. Lett.*, 2005, **7**, 5453.
- A. B. Flynn and W. W. Ogilvie, *Chem. Rev.*, 2007, **107**, 4698.
- With Pd catalyst: (a) M. Sugino, A. Yamamoto, M. Murakami, *J. Am. Chem. Soc.*, 2003, **125**, 6358. (b) M. Daini and M. Sugino, *Chem. Commun.*, 2008, 5224. (c) M. Daini, A. Yamamoto and M. Sugino, *J. Am. Chem. Soc.*, 2008, **130**, 2918. With Ni catalyst: (d) A. Yamamoto and M. Sugino, *J. Am. Chem. Soc.*, 2005, **127**, 15706. (e) M. Sugino, M. Shirakura and A. Yamamoto, *J. Am. Chem. Soc.*, 2006, **128**, 14438. With Cu catalyst: (f) R. Alfaro, A. Parra, J. Aleman, J. L. G. Ruano, M. Tortosa, *J. Am. Chem. Soc.*, 2012, **134**, 15165. (g) L. Zhang, J. Cheng, B. Carry, Z. Hou, *J. Am. Chem. Soc.*, 2012, **134**, 14314. (h) H. Yoshida, I. Kageyuki and K. Takaki, *Org. Lett.*, 2013, **15**, 952. (i) K. Kubota, H. Iwamoto, E. Yamamoto, H. Ito, *Org. Lett.*, 2015, **17**, 620.
- G.-H. Fang, Z.-J. Yan, M.-Z. Deng, *Synthesis*, 2006, 1148.
- See Table S2 in ESI.
- J. J. Daly, R. P. A. Sneeden, H. H. Zeiss, *J. Am. Chem. Soc.*, 1966, **88**, 4287.
- See Scheme S1 in ESI.

ChemCommun

COMMUNICATION

A drastic effect of cobalt and chromium catalysts in the borylation of arylzinc reagents

Kimihiro Komeyama,* Shinnosuke Kiguchi and Ken Takaki



A new synthetic approach to arylboronic esters from arylzinc reagents with boryl electrophiles MeOB(OR)₂ has been developed. Furthermore, this protocol could be applied to the cyclization/borylation of alkynylaryl iodides to afford cyclized vinylboronic esters.