


Chemical Science

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: A. Saha, M. Chen, M. Lederer, A. Kahnt, X. Lu and D. M. Guldi, *Chem. Sci.*, 2016, DOI: 10.1039/C6SC03207A.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Journal Name

ARTICLE

Sulfur Rich Electron Donors – Formation of Singlet versus Triplet Radical Ion Pair States Featuring Different Lifetimes in the Same Conjugate

Avishek Saha,^{†a} Muqing Chen,^{†b} Marcus Lederer,^a Axel Kahnt,^a Xing Lu,^{*b} and Dirk M. Guldi^{*a}

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

An unprecedented family of novel electron-donor acceptor conjugates based on fullerenes as electron acceptors, on one hand, and triphenyl amines as electron donors on the other hand, have been synthesized and characterized in a variety of solvents using steady state absorption/emission as well as transient absorption spectroscopy. Unprecedented in terms of outcome of radical ion pair formation, that is, singlet versus triplet excited state. This was corroborated by femto- / nanosecond pump probe experiments and by molecular orbital calculations. Not only has the donor strength of the triphenylamines been systematically altered by appending one or two sulfur rich dithiafulvenes, but the presence of the latter changed the nature of radical ion pair state. Importantly, depending on the excitation wavelength, that is, either where the fullerenes or where the triphenylamines absorb, short-lived or long-lived radical ion pair states, respectively, are formed. The short-lived component with a lifetime as short as 6 ps has singlet character and stems from a fullerene singlet excited state precursor. In contrast, the long-lived component has a lifetime of up to 130 ns in THF, has triplet character, and evolves from a triplet excited state precursor. Key in forming more than three orders of magnitude longer lived radical ion pair states is the presence of sulfur atoms, which enhance spin-orbit coupling and, in turn, intersystem crossing. Independent confirmation for the singlet versus triplet character came from temperature dependent measurements with focus on the radical ion pair state lifetimes. Here, activation barriers of 2.4 and 10.0 kJ / mol for the singlet and triplet radical ion pair state, respectively, were established.

Introduction

Natural photosynthesis is one of the essential processes that runs organic life on earth.¹ It operates based on the mutual interplay between light harvesting, energy transfer, electron transfer, and catalysis. Inspired by this, a plethora of covalent and non-covalent electron donor-acceptor systems, in which electron donating and electron accepting building blocks are integrated, have been synthesized and investigated. For instance, the most commonly employed electron donors range from chlorophylls and carotenoids to porphyrins / phthalocyanines and triphenylamines. As far as electron acceptors are concerned, perylenebisdiimides, naphthalenediimides, endohedral fullerenes, and empty fullerenes should be mentioned. In the resulting systems, particular focus has been placed on fundamental aspects of electron transfer.

In the context of electron acceptors, fullerenes were extensively studied in electron donor–acceptor systems. This is mainly due to their unique π -electronic nature, their marked excited-state electronic properties, and their low reorganization energy in electron transfer reactions.^{2–8} A myriad of possible electron donors have been investigated as counterpart to the electron accepting fullerene ranging from organic entities like, for example, tetrathiafulvalenes, to complexes based on porphyrins etc.^{6, 9–16} What renders the resulting electron donor-acceptor systems most appealing is the possibility to adjust their structure and, in turn, to fine-tune their physicochemical features such as quantum yields and lifetimes of charge separation.^{17, 18}

Among the electron donors, triphenylamines and their derivatives stand out owing to their good charge transport ability. As a matter of fact, they were widely used as hole injectors or as active materials in organic electronic devices.^{19–23}

Photophysical assays with a system, where C₆₀ is covalently linked to a TPA, revealed that the photo-induced charge separation process predominantly produces TPA^{•+}-C₆₀^{•-} radical ion pair state in polar solvents via ¹C₆₀^{*} excitation.⁶ Later, in the analogues TPA-Sc₃N@C₈₀, in which TPA is connected via the nitrogen, a significant improved thermal stability resulted in longer lived radical ion pair states, when compared to the corresponding 2-substituted conjugate.²⁴ This led us to consider a different strategy to alter the lifetime of the radical ion pair state, namely TPA modification. For

^a Department of Chemistry and Pharmacy & Interdisciplinary Center for Molecular Materials (ICMM), Friedrich-Alexander University of Erlangen-Nürnberg, Egerlandstr. 3, 91058 Erlangen, Germany. Email: dirk.guldi@fau.de

^b State Key Laboratory of Materials Processing & School of Material Science and Engineering, Huazhong University of Science and Technology, 1037 Luoyu Road, 430074 Wuhan, China. Email: lux@hust.edu.cn

[†]Electronic Supplementary Information (ESI) available: Emission spectra, transient absorption spectra, pulse radiolysis spectra, HPLC spectra, description of the synthesis.

See DOI: 10.1039/x0xx00000x

[†]These authors contributed equally.



ARTICLE

example, dithiafulvenes, derivatives of tetrathiafulvalene, are known for their high electron density and hole mobility.^{25, 26} As such, they bear great potential for modifying TPA and, in turn, for extending the lifetime of the radical ion pair state


Like in photosynthesis, the lifetime of the radical ion pair state is crucial to link energy conversion to an efficient energy storage. To this end, spatially separating electron donors and acceptors is one promising approach, while electronically decoupling them is yet another one.²⁷⁻³⁰ Products of both approaches are long-lived radical ion pair states of singlet or triplet nature. A less frequently explored approach implies the conversion of the singlet radical ion pair state character to the corresponding triplet manifold.^{31, 32} Due to the needed spin-conversion, which proceeds via hyperfine induced singlet-triplet mixing, starting with a triplet excited state precursor in electron donor-acceptor systems enables the efficient formation of long-lived, triplet radical ion pairs. Next to transition metals, whose internal heavy-atom effects facilitate the rapid formation of triplet excited state precursors, second-order vibronic spin-orbit coupling in sulfur-containing building blocks provides similar means.^{33, 34} To the best of our knowledge, formation of singlet versus triplet radical ion pairs in, however, the same electron donor-acceptor conjugate is unprecedented. Especially, the simple variation of the excitation wavelength has not been shown to be linked to such a different electron transfer outcome.

In this regard, we have synthesized two novel TPA-C₆₀ electron donor-acceptor conjugates **4** and **5** bearing one or two dithiafulvenes by means of 1,3-dipolar cycloaddition of azomethine ylides – Scheme 1. We have investigated their photophysical properties with a particular focus on photoinduced electron transfer processes, which resulted in radical ion pair states followed by deactivation to the ground state. Importantly, our investigations have enabled for the first time the selective and direct formation of either singlet or triplet radical ion pair states in the same electron donor-acceptor conjugates. To this end, photoexcitation of either the electron accepting fullerenes or the electron donating triphenylamines / dithiafulvenes provide the means for activating different charge separation and recombination pathways. As such, we believe that our work will trend-set the field of electron donor-acceptor design by means of adapting “triplet” precursors.

Results and discussion

Ground state characterization

In the absorption spectrum of TPA-dithiafulvene **2**, strong absorptions in the near-UV region of the optical spectrum in the form of a maximum at 315 nm and a shoulder around 370 nm are noted. N-methylfulleropyrrolidine **6** gives rise to absorptions in the near-UV and blue region of the optical spectrum with a dominating maximum at 325 nm. Regarding TPA-dithiafulvene-C₆₀ **4**, the ground state absorption is best described as the superimposition of the absorptions of **2** and **7**. In addition, a weak absorption tail that extends into the violet / blue region of the solar spectrum emerges. This is explained by the enhanced conjugated π -system in **4** – see Figure 1.



Scheme 1 Synthesis of **4** and **5** as well as reference **6**.

Turning to **3**, absorptions are seen in the UV region of the solar spectrum with a maximum at 317 nm accompanied by a shoulder-like maximum at around 360 nm. For **5**, optical absorptions are located within the same region of the optical spectrum as observed for **3**, with maxima at 338, 369, and 395 nm. Similar to **4**, the absorption of **5** appears as the superimposition of the absorption spectra of **3** and **6** including an additional red shift by about 40 nm of the absorption characteristics located at 400 nm – Figures 2 and S2.






Figure 1. Absorption spectra of **2** (black), **6** (green), and **4** (red) in toluene. The blue line depicts the sum of the absorption spectra of **2** and **6**.

Pulse radiolysis

In order to characterize the absorption spectra of **2**⁺⁺ and **3**⁺⁺, pulse radiolysis experiments under oxidizing condition were performed. **2** and **3** were dissolved in 1-Chlorobutane (BuCl), saturated with N₂ and irradiated with short high energy electron pulses. These conditions lead to the formation of BuCl⁺⁺, which is known as a strong oxidizing agent.^{35, 36} The ability of BuCl⁺⁺ oxidizing aromatic amines via free electron transfer is well established in the literature.^{37, 38} So it is safe to assume that **2** and **3** are oxidized like TPA - described in the literature forming **2**⁺⁺ and **3**⁺⁺ respectively.³⁷ The pulse radiolysis spectrum of solutions of **2** in BuCl shows after the electron pulse the typical transient absorption of BuCl⁺⁺ with a characteristic transient absorption maximum around 520 nm (not shown).^{35, 36, 39} This transient absorption decays rapidly giving rise to a new set of transient absorptions, one covering the visible part of the optical spectrum and one in the NIR - see Figure S10. These transient absorption bands are attributed to the transient absorption of **2**⁺⁺.

Regarding the reaction of **3** with BuCl⁺⁺, again directly after the electron pulse the characteristic transient absorption features of the BuCl⁺⁺ were observed, which rapidly decay, giving rise to new transient absorption bands maximizing at 580 nm and 1100 nm - see Figure S6 - belonging to the transient absorption of **3**⁺⁺.




Figure 2. Absorption spectrum of **5** in toluene.




Figure 3. Optimized structures and electron densities of LUMOs and HOMOs of (a) **4** and (b) **5** calculated at B3LYP/3-21G level of theory.

Molecular Orbital Calculations

Next, we performed computational studies to shed light onto the geometrical and electronic features of **4**, **5**, and C₆₀-TPA lacking dithiafulvenes. All of the structures were optimized at the B3LYP/3-21G level using Gaussian 09 program package. Figure 3 shows the spatial electron densities of the highest occupied molecular orbital (HOMO) and lowest unoccupied molecular orbital (LUMO) for **4**, **5**, and C₆₀-TPA. First of all, the electron density of the HOMO in C₆₀-TPA was found to be located on the aromatic amine moiety, while the LUMO is localized on the C₆₀ spheroid. These preliminary calculations suggest that the anion radical will be localized on the C₆₀ spheroid, while the radical cation will be localized on the aromatic amine moiety in case of a charge separated state, which is in line with the work of O. Ito et al.⁶ Secondly, for **4** and **5**, the corresponding LUMOs are mainly localized on the fullerene, while the spatial electron densities of the HOMO are extended to the side chains, which bear the dithiafulvene ring. In summary, functionalization of the TPA moiety tunes energies of HOMOs (better electron donating ability) while energies of LUMOs remain essentially unchanged – see Figure 3.

Excited State Characterization

First insight into possible electron donor acceptor interactions came from fluorescence assays. **2**, for example, exhibits a strong emission between 400 and 650 nm with a maximum at 435 nm – Figure S3 in the supporting information – and an emission quantum yield of 0.007 in toluene. When turning to **4**, the TPA-dithiafulvene centered emission is completely quenched in toluene and in THF. For TPA-bis-dithiafulvene **3**, the strong emission between 400 and 500 nm maximizes at 425 nm – Figure S3. A value of 0.015 was obtained for the emission quantum yield in toluene. For **5**, the TPA-



ARTICLE

Journal Name

bis-dithiafulvene based emission is fully quenched in toluene and THF.




Figure 4. (a) Transient absorption spectra of **5** in argon saturated THF upon fs-laser photolysis (387 nm, 400 nJ / pulse) with time delays of 2 ps (black), 10 ps (red), 100 ps (green), 1000 ps (blue), and 5000 ps (cyan) after the laser pulse. (b) Corresponding time absorption profile (black) and exponential decay fit (red) at 590 nm. (c) Nanosecond transient absorption spectra of **5** in argon saturated THF upon laser photolysis (387 nm, 400 nJ / pulse) with time delays of 5 ns (black), 25 ns (red), 50 ns (green), 100 ns (blue) and 500 ns (cyan) after the laser pulse. (d) Corresponding time absorption profile (black) and exponential decay fit (red) at 590 nm.

As a complement, we investigated the C_{60} based fluorescence. Here, in order to excite C_{60} exclusively, 470 nm was selected as excitation wavelength. N-methylfulleropyrrolidine **6** was used as reference component with a reported fluorescence quantum yield of 6.0×10^{-4} in toluene.⁴⁰ **4**, **5**, and **6** all show typical N-methylfulleropyrrolidine centered emissions⁴⁰ with maxima located at 710 ± 1 nm – Figure S4. In toluene, only **5** showed a significant quenching of the C_{60} fluorescence with a quantum yield of $3.2 \pm 0.5 \times 10^{-5}$. Turning to THF, the fluorescence of **4** is quenched by an order of magnitude ($5.3 \pm 0.5 \times 10^{-5}$). In **5**, the fluorescence was quantitatively quenched in THF.

In summary, the complete quenching of the TPA-dithiafulvene and TPA-bis-dithiafulvene fluorescence in **4** and **5** relative to references **2** and **3**, respectively, points to an additional decay mechanism – energy transfer and / or electron transfer. Likewise,

quenching of the C_{60} fluorescence in **4** / THF, in **5** / toluene, and in **5** / THF indicates an additional decay process starting from $^1C_{60}^*$.

Transient absorption spectroscopy based on femtosecond and nanosecond laser photolysis was employed to shed light onto the additional decay processes. In the case of **2** and **3**, the singlet excited states are formed immediately after laser excitation. Both show strong transient absorptions throughout the visible and NIR region of the optical spectrum with maxima at 615 nm for **2** or 612 nm for **3** – Figure S5. The latter decay with a lifetime of ~ 2.5 ps and transform into transient absorptions maximizing at either 565 nm (**2**) or 590 nm (**3**). A likely rationale infers formation of the corresponding triplet manifold by means rapid intersystem crossing due to the presence of the sulfur atoms. In both cases, that is, for **2** and **3**, these transient absorptions decay with lifetimes of 370 and 480 ps, respectively, to repopulate the ground state.



Turning to the electron donor-acceptor conjugate **5** upon 387 nm photoexcitation with femtosecond laser pulses in THF, two transient absorption bands appear 3 ps after the laser pulse.⁴¹ For **5**, the maxima occur around 590 and 1010 nm – see Figures 4a and 4b. Notably, the latter absorption is diagnostic for the one electron reduced C₆₀ radical anion, whereas the former matches well the transient absorption fingerprint of the one electron oxidized TPA-bis-dithiafulvene radical cation. The latter was determined via pulse radiolytic oxidation – Figure S6. From this assignment we conclude the successful formation of the TPA-bis-dithiafulvene^{•+}-C₆₀^{•-} radical ion pair state in conjugate **5**. Taking a closer look at the decay of the TPA-bis-dithiafulvene^{•+}-C₆₀^{•-} radical ion pair state in THF the transient absorption features decay biphasic. In particular, a short-lived component was found along with a long-lived component. For the short lived component a lifetime of 6 ps has been derived while the long lived component decays on a timescale beyond the detection limit of our femtosecond laser photolysis setup of 5500 ps – see Figures 4.

In order to reveal the identity and the lifetime of the long lived transient, the same experiments were repeated with a nanosecond laser photolysis set up. Upon photoexcitation at 387 nm, the same transient absorption appeared as observed in the femtosecond laser photolysis experiments – vide supra. In other words, maxima at 590 and 1010 nm confirm the formation of the TPA-bis-dithiafulvene^{•+}-C₆₀^{•-} radical ion pair state. Next to seeing the short lived component with its 6 ps lifetime, the long lived component was seen to decay with a lifetime of 85 ns – Figures 4c and 4d. In Figure S7, normalized transient absorption spectra from femtosecond and nanosecond laser pulse measurements are shown to demonstrate the resemblance of transients across the different timescales. Any notable differences, as they are noted in the near-infrared parts of the transient absorption spectra are based on the different white light sources used for the femto- and nanosecond laser photolysis set ups.

Our findings that the radical ion pair state shows two lifetimes and that a 387 nm excitation excites TPA-bis-dithiafulvene as well as C₆₀ – Figures 1 and 2 – leads us to hypothesize that two radical ion pair states are formed. One of them is born upon exciting C₆₀, while the other stems from exciting TPA-bis-dithiafulvene. The different radical ion pair state lifetimes are only reasonable when different spin multiplicities, that is, singlet versus triplet, are considered. In other words, we postulate the competition of a singlet versus triplet radical ion pair state.

It is well documented that photoexcitation of C₆₀ leads to its singlet first excited state, for which a lifetime of 1.4 ns is found unless it is quenched.⁴² Please note that the initially populated C₆₀ singlet excited state with its characteristic absorption maximum at around 900 nm is invisible due to the rapid occurring charge separation. Considering that the spin polarization remains constant during the electron transfer it is reasonable to postulate the formation of a short lived, singlet radical ion pair state, originating from C₆₀ singlet excited state. This leaves the long lived radical ion pair state with a triplet electron spin multiplicity. The only feasible pathway in electron donor-acceptor conjugate **5** towards a triplet radical ion pair state is a triplet excited state precursor. Here, the strong spin orbit coupling induced by sulfur atoms comes into play as it facilitates in **2** and **3** the intersystem crossing from the TPA-bis-dithiafulvene and TPA-dithiafulvene singlet first excited states to the corresponding triplet excited states.⁴³

The formation of radical ion pair states with different spin multiplicities has been reported by Fukuzumi *et al.*⁴⁴ A common

verification of the triplet spin multiplicity is the quenching of the parent triplet excited state or the triplet radical ion pair state with molecular oxygen. In the current case, this is, however, unfeasible. The triplet quenching by oxygen is based on a bimolecular energy transfer and the lifetimes observed for our compounds are too short to observe reliable and interpretable oxygen effects.

A different approach to verify our hypothesis is based on taking advantage of the different absorption cross sections. Considering, for example, the absorption spectra of TPA-bis-dithiafulvene **3** – Figure 2 – and C₆₀ reference **7** – Figure 1, we performed femtosecond transient absorption experiments by exciting solutions of **5** at 568 nm. Only contributions from the C₆₀ singlet excited state should be discernable, due to the lack of TPA-bis-dithiafulvene absorption. This is, indeed, the case. Importantly, exciting at 568 nm shows transient absorptions exclusively attributed to short lived radical ion pair state – Figure 5 – which is formed with 1.8 ps.

As a complement, we performed femtosecond transient absorption measurements following 420 nm fs-laser excitation. The latter generates both C₆₀ singlet excited state and TPA-bis-dithiafulvene singlet excited state similar to those experiments, in which 387 nm excitation was employed. Notably, the TPA-bis-dithiafulvene singlet excited state transforms rapidly via intersystem crossing into the corresponding triplet manifold. Like in the 387 nm excitation experiments, biphasic decay kinetics are observed. In particular, a short-lived 6 ps component accompanies a long-lived 85 ns component – Figure S8. At the 420 nm excitation, the relative C₆₀ to TPA-bis-dithiafulvene absorption cross section increases and, in turn, contributions from the C₆₀ singlet excited state dominate over those of the TPA-bis-dithiafulvene triplet excited state. In line with the latter, the relative ratio between the short- and long-lived radical ion pair states intensified.






Figure 5. (a) Transient absorption spectra of **5** in argon saturated THF upon fs-laser photolysis (568 nm, 400 nJ / pulse) with time delays 2 ps (black), 5 ps (red), 10 ps (green), 25 ps (blue), and 100 ps (cyan) after the laser pulse. (b) Corresponding time absorption profile (black) and exponential decay fit (red) at 590 nm.

Additionally, the finding of two different radical ion pair states – one with singlet and one with triplet spin multiplicity – was confirmed in temperature dependent laser photolysis transient absorption measurements with, for example, **5**. When reducing the temperature from 300 to 180 K in 2-methyl-THF the lifetime of the radical ion pair state with singlet spin multiplicity increases from 10 ps (300 K) to 19 ps (180 K) – Figure 6. From the fit to the Arrhenius equation a rather small activation barrier for the charge recombination of the singlet radical ion pair state of 2.4 kJ / mol was obtained. When turning to the temperature dependency of the triplet radical ion pair state of **5** in 2-methyl THF the lifetime increases from 92 ns (300 K) to 1090 ns (180 K) – Figure 7. From an Arrhenius analysis, a fairly substantial activation barrier of 10.0 kJ / mol was derived for the charge recombination of the triplet radical ion pair state.

Both activation barriers, that is, 2.4 kJ / mol and 10.0 kJ / mol for the charge recombination of the singlet and triplet radical ion pair states, respectively, are well in line with our expectations. In **5**, the activation barrier for the recombination of the singlet radical ion pair state is comparable to RT at room temperature and, in turn, negligible.




Figure 6. Plot of rate constant for the charge recombination of the singlet radical ion pair state vs T^{-1} for **5** in 2-methyl THF upon photoexcitation at 387 nm. Linear fit is shown in red. Inset: Corresponding time-absorption profiles at 600 nm obtained at 300 K (black), 270 K (red), 240 K (green), 210 K (blue), and 180 K (cyan).




Figure 7. Plot of the rate constant for the charge recombination of the triplet radical ion pair state vs T^{-1} for **5** in 2-methyl THF upon photoexcitation at 387 nm. Linear fit is shown as in red. Inset: Corresponding time-absorption profiles at 600 nm obtained at 300 K (black), 270 K (red), 240 K (green), 210 K (blue) and 180 K (cyan).




Figure 8. Simplified energy diagrams for **5** in THF illustrating on the left and on the right the different deactivation pathways upon photoexcitation of TPA-bis-dithiafulvene (TPASS) at 387, 420, and 568 nm and C_{60} at 387 and 420 nm, respectively. The energy of the radical ion pair state energy was determined as 1.25 eV for **5** from the one electron reduction of C_{60} as -1.19 V versus Fc/Fc^+ and the one electron oxidation of TPASS as +0.05 V versus Fc/Fc^+ and as 1.25 eV for **4** from the one electron reduction of C_{60} as -1.14 V versus Fc/Fc^+ and the one electron oxidation of TPASS as +0.11 V versus Fc/Fc^+ .



In stark contrast, the activation energy for the triplet radical ion pair state is substantially higher due to the presence of parallel electron spins. The latter requires spin flipping as the rate limiting step in the repopulation of the ground state.

Taking the aforementioned into concert, it is safe to conclude that formation of a singlet radical ion pair state results from C₆₀ photoexcitation in **5**, while a triplet radical ion pair state evolves from TPA-bis-dithiafulvene photoexcitation – Figure 8.

Turning to the electron donor-acceptor system **4** featuring only one TPA-dithiafulvene upon laser excitation with 387 nm femtosecond laser pulses evidence for the short lived and the long lived radical ion pair state is noted. In particular, the transient absorption bands maximize at 585 and 1010 nm – Figures S7 and S11. On one hand, the feature at 1010 nm is a perfect match of the one electron reduced C₆₀ radical anion.^{17,45}

On the other hand, the transient absorption in the visible range of the solar spectrum is in sound agreement with the transient absorption of the one electron oxidized TPA-dithiafulvene radical cation obtained in our pulse radiolysis study – Figures S9 and S10. Our finding that the transient absorption matches the transient absorption of the C₆₀ radical anion and the TPA-dithiafulvene radical cation strongly suggests the formation of the TPA-dithiafulvene^{•+}-C₆₀^{•-} radical ion pair state. Like in the case of **5**, the radical ion pair state of **4** decays in 387 nm excitation experiments with two lifetimes, namely 6 ps and 130 ns. One is due to singlet spin multiplicity and one is due to triplet spin multiplicity. In the presence of air the lifetime of the 130 ns is reduced to 31 ns from which we conclude an activation controlled deactivation of the radical ion pair state.

Considering similar absorption spectra for **4** and **5**, 387 nm excitation leads to the formation of the C₆₀ singlet excited state as well as to the generation of the TPA-dithiafulvene singlet excited state. The latter, however, rapidly transforms via intersystem crossing to the TPA-dithiafulvene triplet excited state. Again, we postulate that from the C₆₀ singlet excited state the short lived, singlet radical ion pair state evolves, while the TPA-dithiafulvene triplet excited state is the precursor to the long lived, triplet radical ion pair state. Support for this notion came from femtosecond transient absorption measurements exciting at 568 nm. Similar to what was described for **5**, the 568 nm excitation results exclusively in the C₆₀ singlet excited state and, in turn, the singlet radical ion pair state formation. Importantly, the TPA-dithiafulvene^{•+}-C₆₀^{•-} transient absorptions decay with a 6 ps component. Here, no long lived transient absorption remains on the time scale beyond 10 ps – Figure S12.

Conclusions

In summary, we have observed fast, photoinduced charge separation in a set of two novel electron donor-acceptor conjugates based on fullerenes and sulfur-containing triphenylamines. Importantly, depending on the excitation wavelength, that is, either where the fullerenes or where the triphenylamines absorb, *for the first time* short-lived or long-lived TPA-dithiafulvene^{•+}-C₆₀^{•-} / TPA-bis-dithiafulvene^{•+}-C₆₀^{•-} radical ion pair states, respectively, are formed in the same conjugate. On one hand, the short-lived component with a lifetime as short as 6 ps has singlet spin multiplicity and stems from a fullerene singlet excited state precursor. On the other hand, the long-lived component has a

lifetime of up to 130 ns in THF, has triplet spin multiplicity, and evolves from a triplet excited state precursor. Both are energetically nearly degenerate, but the exact determination requires application of a varying external field. Key in forming more than three orders of magnitude longer lived triplet radical ion pair states is the presence of sulfur atoms, which enhance spin-orbit coupling and, in turn, render intersystem crossing of TPA-dithiafulvene / TPA-bis-dithiafulvene in the excited state fast with a lifetime of about 2.5 ps. The change in spin multiplicity of the radical ion pair state is likely to be slower. At room temperature, upper and lower limits are given by the radical ion pair states of 6 ps and 130 ns, respectively. Notably, the interconversion between the singlet and triplet radical ion pair states is characterized by magnetic field strength dependence. The physical process, which allows for such detailed analysis, involves Zeeman splitting of the T₀ state by the magnetic field. It can be modulated by the strength of the magnetic field, which is a path that we are currently exploring in our laboratory. Regardless of the respective spin multiplicity both radical ion pair states decay directly to the ground state without forming a triplet excited state. In other words, one state decays in a spin allowed and fast manner, while the other one decays spin forbidden and slow. Independent confirmation for the singlet versus triplet character came from temperature dependent measurements with focus on the radical ion pair state lifetimes. Here, activation barriers of 2.4 and 10.0 kJ / mol for the singlet and triplet radical ion pair state, respectively, were established. Our results document the incentives of a “triplet” precursor and, as such, will pave the way towards new electron donor-acceptor materials for photovoltaics and photocatalysis.

Experimental Section

General Methods:

All reactions were performed under an argon atmosphere. The ¹H and ¹³C NMR chemical shifts are given relative to tetramethylsilane (TMS). All chemicals were purchased and used as received unless otherwise noted. Triphenylamine (TPA) was purchased from Aldrich Chemical Co. 4,5-bis(hexylthio)-1,3-dithiole-2-thione was prepared according to the reported procedures.⁴⁶

Synthesis:

The details of synthesis of **1a**, **1b**, **2**, **3**, **4** and **5** as well as their characterizations are provided in the supporting information.

Photophysical studies:

Steady-state UV/Vis absorption spectra were measured on PerkinElmer Lambda 35 and PerkinElmer Lambda 2 two-beam spectrophotometers. Steady-state emission spectra were recorded with a FluoroMax P spectrometer (Horiba Jobin Yvon). The experiments were performed at room temperature. Emission quantum yields for the TPA-derivative based emission were determined by the comparative method using 9, 10-diphenylanthracene as references with a fluorescence quantum yield of 0.9 in cyclohexane.⁴⁷

Femtosecond transient absorption laser photolysis measurements were performed with an output from a Ti:sapphire laser system (CPA2110, Clark-MXR Inc.): 775 nm, 1 kHz, and 150 fs FWHM pulses. The excitation wavelength was either generated by second harmonic generation (387 nm) or using a NOPA (NOPA Plus – Clark MXR) (420 and 568 nm). For all excitation wavelengths, pulse widths of <150 fs and energies of 400 nJ / pulse were selected. The



ARTICLE

Journal Name

transient absorption measurements were performed with an HELIOS (Ultrafast Systems LLC) transient absorption spectrometer.

Nanosecond transient absorption measurements were performed with an EOS transient absorption spectrometer (Ultrafast Systems LLC). The pump pulses were generated from the amplified Ti:Sapphire laser system described above. The probe pulse (2 kHz, 0.5 ns pulse width), which was generated in a 1064 nm pumped photonic fiber, synchronized with the femtosecond amplifiers.

Conventional nanosecond laser flash photolysis transient absorption measurements were carried out with the output from the third harmonic (355 nm, 10 mJ) of a Nd:YAG laser (Brilliant B, Quantel). The optical detection is based on a pulsed (pulser MSP 05 – OptikElektronik Müller) Xenon lamp (XBO 450, Osram), a monochromator (Spectra Pro 2300i, Acton Research), a fast InGaAs photodiode (Nano 5, Coherent) with 300-MHz amplification, and a 1-GHz digital oscilloscope (WavePro7100, LeCroy).

Pulse radiolysis:

The samples were dissolved in 1-chlorobutane, saturated with N₂ and irradiated with high energy electron pulses (1 MeV, 15 ns duration) by a pulse transformer type electron accelerator (Elit - Institute of Nuclear Physics, Novosibirsk, Russia). The dose delivered per pulse was measured by electron dosimetry.⁴⁸ Doses of 100 Gy were employed. The optical detection of the transients was carried out with a detection system consisting of a pulsed (pulser MSP 05 - Müller Elektronik Optik) Xenon lamp (XBO 450, Osram), a SpectraPro 500 monochromator (Acton Research Corporation), a R 9220 photomultiplier (Hamamatsu Photonics), and a 500 MHz digitizing oscilloscope (TDS 640, Tektronix).

Quantum Chemical Calculations:

Quantum chemical calculations were performed based on the DFT method using the B3LYP/3-21G functional as implemented in the Gaussian 09 program package.⁴⁹

Electrochemical measurements:

Cyclic voltammetry was performed by using a FRA 2 μ Autolab type III (METROHM) potentiostat, in a conventional three-electrode electrochemical cell. A glass carbon disk (1 mm diameter) was used as working electrode, a Pt wire electrode served as the counter electrode, an Ag wire was used as the quasi reference electrode. Typically, 100 μ M solution of **4** and **5** were prepared in dichloromethane while 0.1 M (n-Bu)₄NPF₆ was used as the supporting electrolyte. All potentials were referenced to ferrocenium/ferrocene (Fc⁺/Fc). Measurements were performed using 50 mV/s scan rate after purging with argon at ambient temperature.

Acknowledgements

Financial support from Bavarian “Solar Technologies go Hybrid” – initiative (SolTech) and from the German Science Foundation through SFB 953, is gratefully acknowledged. AK gratefully acknowledges funding from the Deutsche Forschungsgemeinschaft (DFG) via grant KA 3491/2-1. Financial support from The National Thousand Talents Program of China, NSFC (21171061, 51472095), Program for Changjiang Scholars and Innovative Research Team in University (IRT1014) and HUST is gratefully acknowledged.

Notes and references

1. J. Barber, *Chem. Soc. Rev.*, 2009, **38**, 185-196.
2. N. Martín, L. Sánchez, B. Illescas and I. Pérez, *Chem. Rev.*, 1998, **98**, 2527-2548.
3. L. Echegoyen and L. E. Echegoyen, *Acc. Chem. Res.*, 1998, **31**, 593-601.
4. D. M. Guldi and M. Prato, *Acc. Chem. Res.*, 2000, **33**, 695-703.
5. D. Gust, T. A. Moore and A. L. Moore, *Acc. Chem. Res.*, 2001, **34**, 40-48.
6. H.-P. Zeng, T. Wang, A. S. D. Sandanayaka, Y. Araki and O. Ito, *J. Phys. Chem. A*, 2005, **109**, 4713-4720.
7. L. Feng, S. Gayathri Radhakrishnan, N. Mizorogi, Z. Slanina, H. Nikawa, T. Tsuchiya, T. Akasaka, S. Nagase, N. Martín and D. M. Guldi, *J. Am. Chem. Soc.*, 2011, **133**, 7608-7618.
8. S. Kirner, M. Sekita and D. M. Guldi, *Adv. Mater.*, 2014, **26**, 1482-1493.
9. S. Castellanos, A. A. Vieira, B. M. Illescas, V. Sacchetti, C. Schubert, J. Moreno, D. M. Guldi, S. Hecht and N. Martín, *Angew. Chem. Int. Ed.*, 2013, **52**, 13985-13990.
10. G. Kodis, P. A. Liddell, L. de la Garza, A. L. Moore, T. A. Moore and D. Gust, *J. Mater. Chem.*, 2002, **12**, 2100-2108.
11. C. Villegas, E. Krokos, P.-A. Bouit, J. L. Delgado, D. M. Guldi and N. Martín, *Energy Environ. Sci.*, 2011, **4**, 679-684.
12. J. Lehl, M. Vartanian, M. Holler, J.-F. Nierengarten, B. Delavaux-Nicot, J.-M. Strub, A. Van Dorsselaer, Y. Wu, J. Mohanraj, K. Yoosaf and N. Armario, *J. Mater. Chem.*, 2011, **21**, 1562-1573.
13. Y. Takano, S. Obuchi, N. Mizorogi, R. García, M. Á. Herranz, M. Rudolf, D. M. Guldi, N. Martín, S. Nagase and T. Akasaka, *J. Am. Chem. Soc.*, 2012, **134**, 19401-19408.
14. A. Molina-Ontoria, M. Wielopolski, J. Gebhardt, A. Gouloumis, T. Clark, D. M. Guldi and N. Martín, *J. Am. Chem. Soc.*, 2011, **133**, 2370-2373.
15. S. K. Das, B. Song, A. Mahler, V. N. Nesterov, A. K. Wilson, O. Ito and F. D'Souza, *J. Phys. Chem. C*, 2014, **118**, 3994-4006.
16. B. D. Allen, A. C. Benniston, A. Harriman, L. J. Mallon and C. Pariani, *Phys. Chem. Chem. Phys.*, 2006, **8**, 4112-4118.
17. W. Seitz, A. Kahnt, D. M. Guldi and T. Torres, *J. Porphyrins Phthalocyanines*, 2009, **13**, 1034-1039.
18. N. Yoshida, T. Ishizuka, A. Osuka, D. H. Jeong, H. S. Cho, D. Kim, Y. Matsuzaki, A. Nogami and K. Tanaka, *Chemistry – A European Journal*, 2003, **9**, 58-75.
19. Y. Shirota, *J. Mater. Chem.*, 2005, **15**, 75-93.
20. L. Liu, H. Li, J. Bian, J. Qian, Y. Wei, J. Li and W. Tian, *New J. Chem.*, 2014, **38**, 5009-5017.
21. Z. Jiang, T. Ye, C. Yang, D. Yang, M. Zhu, C. Zhong, J. Qin and D. Ma, *Chem. Mater.*, 2011, **23**, 771-777.
22. J.-L. Wang, Z. He, H. Wu, H. Cui, Y. Li, Q. Gong, Y. Cao and J. Pei, *Chem. Asian J.*, 2010, **5**, 1455-1465.
23. H. Shang, H. Fan, Y. Liu, W. Hu, Y. Li and X. Zhan, *Adv. Mater.*, 2011, **23**, 1554-1557.
24. J. R. Pinzón, D. C. Gasca, S. G. Sankaranarayanan, G. Bottari, T. Torres, D. M. Guldi and L. Echegoyen, *J. Am. Chem. Soc.*, 2009, **131**, 7727-7734.
25. C. A. Christensen, A. S. Batsanov and M. R. Bryce, *J. Am. Chem. Soc.*, 2006, **128**, 10484-10490.



26. N. Martín, L. Sánchez, M. Á. Herranz, B. Illescas and D. M. Guldi, *Acc. Chem. Res.*, 2007, **40**, 1015-1024.
27. H. Imahori, D. M. Guldi, K. Tamaki, Y. Yoshida, C. Luo, Y. Sakata and S. Fukuzumi, *J. Am. Chem. Soc.*, 2001, **123**, 6617-6628.
28. S. Gélinas, A. Rao, A. Kumar, S. L. Smith, A. W. Chin, J. Clark, T. S. van der Poll, G. C. Bazan and R. H. Friend, *Science*, 2014, **343**, 512-516.
29. H. Nakanotani, T. Furukawa, K. Morimoto and C. Adachi, *Science Advances*, 2016, **2**.
30. L. R. Sutton, M. Scheloske, K. S. Pirner, A. Hirsch, D. M. Guldi and J.-P. Gisselbrecht, *J. Am. Chem. Soc.*, 2004, **126**, 10370-10381.
31. M. Rudolf, L. Feng, Z. Slanina, T. Akasaka, S. Nagase and D. M. Guldi, *J. Am. Chem. Soc.*, 2013, **135**, 11165-11174.
32. A. Rao, P. C. Y. Chow, S. Gelinas, C. W. Schlenker, C.-Z. Li, H.-L. Yip, A. K. Y. Jen, D. S. Ginger and R. H. Friend, *Nature*, 2013, **500**, 435-439.
33. N. Martín, L. Sánchez, B. Illescas, S. González, M. Angeles Herranz and D. M. Guldi, *Carbon*, 2000, **38**, 1577-1585.
34. A. Rodriguez-Serrano, V. Rai-Constapel, M. C. Daza, M. Doerr and C. M. Marian, *Phys. Chem. Chem. Phys.*, 2015, **17**, 11350-11358.
35. H. Mohan, O. Brede and J. P. Mittal, *J. Photochem. Photobiol. A*, 2001, **140**, 191-197.
36. O. Brede, H. Orthner, V. Zubarev and R. Hermann, *J. Phys. Chem.*, 1996, **100**, 7097-7105.
37. A. Maroz, R. Hermann, S. Naumov and O. Brede, *J. Phys. Chem. A*, 2005, **109**, 4690-4696.
38. O. Brede, A. Maroz, R. Hermann and S. Naumov, *J. Phys. Chem. A*, 2005, **109**, 8081-8087.
39. O. Brede, R. Mehnert and W. Naumann, *Chem. Phys.*, 1987, **115**, 279-296.
40. R. M. Williams, J. M. Zwier and J. W. Verhoeven, *J. Am. Chem. Soc.*, 1995, **117**, 4093-4099.
41. The formation of the radical ion pair state is based on contributions from a C₆₀ precursor state and a TPA-bis-dithiafulvene precursor state including intersystem crossing of the latter.
42. B. Ma, C. E. Bunker, R. Guduru, X.-F. Zhang and Y.-P. Sun, *J. Phys. Chem. A*, 1997, **101**, 5626-5632.
43. Interestingly, the triplet excited state of TPA-dithiafulvene is also overlaid with features of radical ion pair state.
44. S.-H. Lee, A. G. Larsen, K. Ohkubo, Z.-L. Cai, J. R. Reimers, S. Fukuzumi and M. J. Crossley, *Chem. Sci.*, 2012, **3**, 257-269.
45. D. M. Guldi, H. Hungerbühler, E. Janata and K.-D. Asmus, *J. Chem. Soc., Chem. Commun.*, 1993, 84-86.
46. Z. Wan, C. Jia, Y. Duan, X. Chen, Y. Lin and Y. Shi, *Organic Electronics*, 2013, **14**, 2132-2138.
47. S. Hamai and F. Hirayama, *J. Phys. Chem.*, 1983, **87**, 83-89.
48. E. M. Fielden, in *The Study of Fast Processes and Transient Species by Electron Pulse Radiolysis*, ed. F. B. John H. Baxendale, NATO Advances Study Institutes Series C. Mathematical and Physical Sciences 1982, vol. 86, pp. 49-62.
49. M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima,

Y. Honda, O. Kitao, H. Nakai, T. Vreven, J. A. Montgomery Jr., J. E. Peralta, F. Ogliaro, M. J. Bearpark, J. Heyd, E. N. Brothers, K. N. Kudin, V. N. Staroverov, R. Kobayashi, J. Normand, K. Raghavachari, A. P. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, N. J. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, Ö. Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski and D. J. Fox, Gaussian, Inc., Wallingford, CT, USA 2009.

