

This article was downloaded by: [Northeastern University]

On: 04 January 2015, At: 05:00

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

An Improved Synthesis of Symmetrical Diselenides from Aromatic Aldehydes

Yahong Chen^a, Fengshou Tian^{a b}, Maoping Song^b & Shiwei Lu^c

^a Department of Chemistry, Zhoukou Normal University, Zhoukou, Henan, P. R. China

^b Department of Chemistry, Zhengzhou University, Zhengzhou, Henan, P. R. China

^c Dalian Institute of Chemical Physics, Dalian, Liaoning, P. R. China

Published online: 23 Mar 2010.

To cite this article: Yahong Chen, Fengshou Tian, Maoping Song & Shiwei Lu (2010) An Improved Synthesis of Symmetrical Diselenides from Aromatic Aldehydes, *Phosphorus, Sulfur, and Silicon and the Related Elements*, 185:4, 799-802, DOI: [10.1080/10426500902967912](https://doi.org/10.1080/10426500902967912)

To link to this article: <http://dx.doi.org/10.1080/10426500902967912>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

AN IMPROVED SYNTHESIS OF SYMMETRICAL DISELENIDES FROM AROMATIC ALDEHYDES

Yahong Chen,¹ Fengshou Tian,^{1,2} Maoping Song,²
and Shiwei Lu³

¹Department of Chemistry, Zhoukou Normal University, Zhoukou, Henan,
P. R. China

²Department of Chemistry, Zhengzhou University, Zhengzhou, Henan, P. R. China

³Dalian Institute of Chemical Physics, Dalian, Liaoning, P. R. China

An improved method for the synthesis of symmetrical diselenides is described. Reductive selenation of aromatic aldehydes with Se/CO/H₂O/Et₃N system in THF gave the corresponding aromatic diselenides in good yields.

Keywords Aromatic aldehydes; carbon monoxide; diselenides; selenium

INTRODUCTION

Organoselenium compounds are very important organic synthetic reagents and intermediates.¹ Among them, symmetrical diselenides are versatile reagents and potential medicinal agents.² Several methods have been reported for the synthesis of symmetrical diselenides.³ For example, Nishiyama et al. found that aliphatic aldehydes and ketones can reduce selenation to aliphatic diselenides using a Se/CO/H₂O/base.⁴

RESULTS AND DISCUSSION


In the course of our studies on selenium-catalyzed reductive carbonylation of nitroaromatic compounds with carbon monoxide, we developed a new application of the Se/CO/H₂O system to convert aromatic aldehydes,⁵ organic halides,⁶ or imines⁷ to symmetrical diselenides. However, the disadvantages of these methods are long reaction times (7–9 h) and undesirable byproducts (symmetrical urea). We report in this article an efficient method for synthesizing symmetrical diselenides (yields up to 89%) by reactions of selenium and aromatic and heterocyclic aldehydes in the presence of carbon monoxide, water, and triethylamine (Scheme 1).

Treatment of benzaldehyde (**1a**) with one equivalent selenium in THF at 100°C in the presence of carbon monoxide (3 Mpa), water, and Et₃N for 3 h led to efficient reductive

Received 26 September 2008; accepted 14 April 2009.

We acknowledge the financial support from China Postdoctoral Science Foundation (No. 20070410876).

Address correspondence to Fengshou Tian, Department of Chemistry, Zhoukou Normal University, Zhoukou, Henan 466000, P. R. China. E-mail: fengshoutian@163.com



a) Ar = C₆H₅; b) Ar = *p*-MeC₆H₄; c) Ar = *m*-MeC₆H₄; d) Ar = *o*-MeC₆H₄; e) Ar = *p*-MeOC₆H₄; f) Ar = *o*-MeOC₆H₄; g) Ar = *m*-C₆H₄OH; h) Ar = 2-thienyl; i) Ar = 3-thienyl; j) Ar = 3-pyridyl

Scheme 1

selenation to give dibenzyl diselenide (**1b**) in 85% yield (Table I, entry 1). No byproduct such as benzyl alcohol or toluene was detected from the reaction mixture. Although the photochemical reaction of ketones and aldehydes with hydrogen selenide afforded alcohols,⁸ acetophenone did not undergo selenation, and only a trace amount of ethylbenzene was detected in the reaction mixture (GC-MS analysis).⁹ The reaction did not proceed without water or without use of base. The results with other aromatic and heterocyclic aromatic aldehydes are summarized in Table I. Increasing steric hindrance on the aromatic ring led to decreased yields, but steric effects of the substituents only slightly affect yields of the diselenides (Table I, entries 1–6). Salicylaldehyde was unchanged under the reaction conditions (Table I, entry 7). Heterocyclic-aromatic aldehydes (2- and 3-thiophenecarboxaldehydes,

Table I Synthesis of symmetrical diselenides from aromatic aldehydes^a

Entry	Diselenide	Yield (%)	Mp (°C) (lit.)	¹ H NMR (δ)	¹³ C NMR (δ)
1	2a	85 ^b	92–93 (90–91) ¹⁰	7.37–7.26 (m, 10H), 3.88 (s, 4H)	139.05, 129.04, 128.45, 127.21, 32.66
2	2b	89	59–60 (61–62) ¹¹	7.26–7.10 (s, 8H), 3.84 (s, 4H), 2.32 (s, 6H)	149.70, 129.84, 126.56, 112.44, 40.60, 32.90
3	2c	82	Yellow liquid ⁵	7.36–7.22 (m, 8H), 3.98 (s, 4 H), 2.52 (s, 6 H)	138.62, 137.54, 129.5, 128.05, 127.56, 125.82, 32.38, 21.19
4	2d	78	80–81 (86–86.5) ^{3c}	7.42–6.22 (m, 8H), 3.96 (s, 4H), 3.81 (s, 6H)	152.32, 142.85, 111.03, 108.88, 107.70, 24.86, 18.91
5	2e	86	72–73 (72) ¹²	7.26–6.83 (m, 8H), 3.83 (s, 4H), 3.80 (s, 6H)	138.99, 128.21, 125.82, 122.52, 26.36
6	2f	81	66–67 (–) ¹³	7.26–6.84 (m, 4H), 3.98 (s, 4H), 3.87 (s, 6H)	149.84, 148.36, 136.22, 134.52, 123.32, 28.72, 24.53
7	2g	—	—	—	—
8	2h	82	Yellow liquid ⁵	7.43–6.23 (m, 6H), 3.96 (s, 4 H)	152.23, 142.64, 111.35, 108.54, 24.52
9	2i	85	94–96 ⁵ (94–96) ⁵	7.27–7.02 (m, 6H), 3.86 (s, 4 H)	138.99, 128.28, 125.80, 122.52, 26.36
10	2j	88	77–79 ⁵ (77–79) ⁵	8.46–7.18 (m, 8H), 3.72 (s, 4H)	149.84, 148.39, 136.22, 123.32, 28.72

^aAll experimental reaction.

^bYield is only 32% in toluene.

and 3-pyridinecarboxaldehyde) were also efficiently transformed to their corresponding diselenides in good yields (82%–88%) (Table I, entries 8–10).

EXPERIMENTAL

All melting points were recorded on a WRS-1A melting-point apparatus and are uncorrected. All ^1H NMR spectra were recorded on a 400-MHz Bruker AZ 400 spectrometer. Chemical shifts are given as δ value with reference to tetramethylsilane (TMS) as internal standard. Carbon monoxide (99.9%) was dried by zeolite 5A. Elemental selenium (99.5%) and the reagents were received from commercial supply without purification prior to use.

PROCEDURE FOR SYNTHESIS OF AROMATIC DISELENIDES

The aromatic aldehyde (2.5 mmol), selenium (2.5 mmol), water (25 mmol), Et_3N (2.5 mmol), and THF (10 mL) and a magnetic stirring bar were successively loaded into a 70-mL stainless-steel autoclave. The reactor was sealed, flushed three times with 1.0 MPa of carbon monoxide, pressurized with 3.0 MPa of carbon monoxide, and then placed in an oil bath preheated to 100°C . After the reaction was complete, the apparatus was cooled to ambient temperature, and the remaining carbon monoxide was evacuated. Then water (20 mL) was added, and the reaction mixture was extracted with diethyl ether (3×40 mL). The combined organic extracts were dried over anhydrous MgSO_4 , filtered, and the solvent was evaporated under reduced pressure to afford a yellow oil. Further purification by column chromatography on silica gel or TLC gave the pure product. All the products were identified by NMR and/or comparison with authentic samples.

REFERENCES

- (a) A. Krief, In *Comprehensive Organometallic Chemistry*, E. W. Abel, F. G. A. Stone, G. Wilkinson, and A. Mckillop, eds. (Pergamon, Oxford, UK, 1995), vol. II, p. 516; (b) T. G. Back, Ed., *Organoselenium Chemistry. A Practical Approach* (Oxford University Press, Oxford, UK, 1999); (c) G. Mugesh, W. Du Mont, and H. Sies, *Chem. Rev.*, **101**, 2125 (2001).
- (a) B. C. Ranu, T. Mandal, and S. Samanta, *Org. Lett.*, **5**, 1439 (2003); (b) N. Taniguchi and T. Onami, *J. Org. Chem.*, **69**, 915 (2004); (c) T. Nishino, M. Okada, T. Kuroki, T. Watanabe, Y. Nishiyama, and N. Sonoda, *J. Org. Chem.*, **67**, 8696 (2002); (d) G. Ten Brink, J. Vis, I. W. C. E. Arends, and R. A. Sheldon, *J. Org. Chem.*, **66**, 2429 (2001); (e) I. Andreadou, W. M. P. B. Menge, J. N. M. Commandeur, E. A. Worthington, and N. P. E. Vermeulen, *J. Med. Chem.*, **39**, 2040 (1996).
- (a) J. Gladysz, J. Hornby, and J. E. Garbe, *J. Org. Chem.*, **43**, 1204 (1978); (b) L. Syper and J. Mlochowshi, *Tetrahedron*, **44**, 6119 (1988); (c) D. P. Thompson and P. Boudjouk, *J. Org. Chem.*, **53**, 2109 (1988); (d) J. Q. Li, W. L. Bao, P. Lue, and X. J. Zhou, *Synth. Commun.*, **21**, 799 (1991); (e) J. X. Wang, W. Cui, and Y. Hu, *J. Chem. Soc., Perkin Trans. 1*, 2341 (1994); (f) A. Krief and M. Derock, *Tetrahedron Lett.*, **43**, 3083 (2002); (g) A. Krief, C. Delmotte, and W. Dumont, *Tetrahedron*, **53**, 12147 (1997); (h) A. Krief, W. Dumont, and C. Delmotte, *Angew. Chem., Int. Ed.*, **39**, 1669 (2000); (i) P. Salama and C. Bernard, *Tetrahedron Lett.*, **36**, 5711 (1995); (j) P. Salama and C. Bernard, *Tetrahedron Lett.*, **39**, 745 (1998); (k) K. Prabhu and S. Chandrasekaran, *Chem. Commun.*, 1021 (1997); (l) A. Krief, A. F. De Mahieu, W. Dumont, and M. Trabelsi, *Synthesis*, 131 (1988); (m) A. Krief, T. Van Wemmel, M. Redon, W. Dumont, and C. Delmotte, *Angew. Chem., Int. Ed.*, **38**, 2245 (1999).

4. Y. Nishiyama, S. Hamanaka, A. Ogawa, S. Murai, and N. Sonoda, *Synth. Commun.*, **16**, 1059 (1986).
5. F. S. Tian, Z. K. Yu, and S. W. Lu, *J. Org. Chem.*, **69**, 4520 (2004).
6. Y. H. Chen, F. S. Tian, M. P. Song, and S. W. Lu, *Synth. Commun.*, **37**, 2699 (2007).
7. X. D. Zhao, Z. K. Yu, and F. L. Zeng, *Adv. Synth. Catal.*, **347**, 877 (2005).
8. K. Kambe, K. Kondo, S. Murai, and N. Sonoda, *Angew. Chem., Int. Ed. Engl.*, **19**, 1008 (1980).
9. Y. Nishiyama, S. Hamanaka, A. Ogawa, N. Kambe, and N. Sonoda, *J. Org. Chem.*, **53**, 1326 (1988).
10. M. D. Ruan, H. R. Zhao, W. Q. Fan, and X. J. Zhou, *J. Organomet. Chem.*, **19**, 485 (1995).
11. J. W. Lewicki, W. H. H. Günther, and J. Y. C. Chu, *J. Org. Chem.*, **43**, 2672 (1978).
12. V. I. Cohen, *J. Org. Chem.*, **42**, 510 (1977).
13. V. Saravanan, E. Porhiel, and S. Chandrasekaran, *Tetrahedron Lett.*, **44**, 2257 (2003).