

277. Transition Metal Complexes with Bidentate Ligands Spanning *trans*-Positions. VI¹⁾. The Preparation of Some Diaryl- and Dialkyl Derivatives of 2,11-Bis (phosphinomethyl)benzo [*c*]phenanthrene

by Pramesh N. Kapoor and Luigi M. Venanzi



Laboratorium für anorganische Chemie, ETH-Zentrum, CH-8092 Zürich, Switzerland

(4.X.77)

Summary

The preparation of the ditertiary phosphines 2,11-bis(di-*m*-tolylphosphinomethyl)benzo [*c*]phenanthrene (**1b**), 2,11-bis(di-*p*-anisylphosphinomethyl)benzo [*c*]phenanthrene (**1c**), 2,11-bis(di-*m*-trifluoromethylphenylphosphinomethyl)benzo [*c*]phenanthrene (**1d**), 2,11-bis(dicyclohexylphosphinomethyl)benzo [*c*]phenanthrene (**1e**) and 2,11-bis(di-*t*-butylphosphinomethyl)benzo [*c*]phenanthrene (**1f**), by a combination of synthetic routes is described.

Introduction. - In earlier publications [1] the preparation of the ditertiary phosphine 2,11-bis(diphenylphosphinomethyl)benzo [*c*]phenanthrene (**1a**) and some of its coordination chemistry were described. The development of this study required the preparation of complexes of analogous phosphines having different electronic and steric properties and/or different solubility characteristics. This paper describes the preparation of ligands **1b** to **1f**.




- 1a** X = Ph₂P-
- 1b** X = (*m*-CH₃ · C₆H₄)₂P-
- 1c** X = (*p*-CH₃O · C₆H₄)₂P-
- 1d** X = (*m*-CF₃ · C₆H₄)₂P-
- 1e** X = (C₆H₁₁)₂P-
- 1f** X = *t*-Bu₂P-
- 1g** X = Br

Results and discussion. - The new ligands were prepared using a combination of the reactions shown in the *Scheme*. Relevant comments to these reactions are:

Reaction 1. The cleavage of a P-C bond in (*m*-CH₃ · C₆H₄)₃P, using sodium in liquid ammonia [2], followed by hydrolysis, gave (*m*-CH₃ · C₆H₄)₂PH in good yield. The same reaction using (*m*-CF₃ · C₆H₄)₃P, however, led only to the partial recovery of starting material.

¹⁾ Part V: See [1].



(i) Na in liq. NH_3 followed by hydrolysis. (ii) LiAlH_4 in Et_2O . (iii) $n\text{-BuLi}$ in THF followed by **1g**. (iv) **1g** in *iso*- BuC(O)Me followed by NaOH . (v) RMgBr followed by HCl . (vi) $n\text{-BuLi}$ in THF followed by **1g** in THF. (vii) SiHCl_3 and $n\text{-Bu}_3\text{N}$ in MeCN . (viii) $\text{R}'\text{O}^-$ in $\text{R}'\text{OH}$. (ix) **1g**.

Reaction 2. Compounds of the type $\text{R}_2\text{P(O)X}$ have been reduced to corresponding secondary phosphines using a variety of reducing agents, the most common being LiAlH_4 [3]. This reagent was successfully used for the reduction of (*m*- $\text{CF}_3 \cdot \text{C}_6\text{H}_4$) $_2\text{P(O)H}$ to **3d**. However, from the LiAlH_4 reduction of (*p*- $\text{CH}_3\text{O} \cdot \text{C}_6\text{H}_4$) $_2\text{P(O)H}$ no secondary phosphine was formed and only small amounts of (*p*- $\text{CH}_3\text{O} \cdot \text{C}_6\text{H}_4$) $_2\text{PCH}_3$ could be isolated. This type of reaction has been observed previously [3], e.g., the reduction of $\text{Ph}(\textit{p}\text{-CH}_3\text{O} \cdot \text{C}_6\text{H}_4)\text{P(O)H}$ gave a mixture of $\text{Ph}(\textit{p}\text{-CH}_3\text{O} \cdot \text{C}_6\text{H}_4)\text{PH}$ and $\text{Ph}(\textit{p}\text{-CH}_3\text{O} \cdot \text{C}_6\text{H}_4)\text{PCH}_3$. However, (*p*- $\text{CH}_3\text{O} \cdot \text{C}_6\text{H}_4$) $_2\text{P(O)H}$ was not reduced by $\text{Li}[\text{AlH}(\text{O}-t\text{Bu})_3]$ in refluxing THF and its reaction with Si_2Cl_6 in benzene gave several products which were not investigated further.

Reaction 3. This is the standard method of preparation of ligand **1a** [4] and was successfully used for the preparation of ligands **1b** and **1d**. When R_2PH ($\text{R} = \text{C}_6\text{H}_{11}$ and *t*-Bu) were used in this reaction it was shown by ^{31}P -NMR. spectroscopy that mixtures of products were formed. The desired compounds **1e** and **1f** were present in significant amounts but could not be conveniently isolated.

Reaction 4. This reaction has been extensively used for the preparation of tertiary phosphines containing *t*-butyl groups [5]. Ligand **1f** was obtained by this method in ca. 30% yield. Preliminary results indicate that ligand **1e** can also be obtained by this reaction.

Reaction 5. This appears to be one of the most convenient methods for the selective introduction of two R-groups onto a phosphorus atom [6]. It was used to obtain intermediates **2c**, **2d** and **2e**.

Reaction 6. Hays [6] reports the successful preparation of tertiary phosphine oxides by the reaction of alkyl halides on $\text{R}_2\text{P(O)MgX}$ obtained *in situ* by reaction 5. Better results were obtained by starting from pure $\text{R}_2\text{P(O)H}$, adding $n\text{-BuLi}$ followed by **1g**. While this reaction gave the desired product when R was *p*- $\text{CH}_3\text{O} \cdot \text{C}_6\text{H}_4$, when R was C_6H_{11} a complex mixture of products, containing some **4e**, was obtained.

Reaction 7. Although the reduction of tertiary phosphine oxides to the corresponding phosphines can be carried out using a variety of reagents [7], difficulties were encountered in reducing ditertiary bis-phosphine oxides. Thus, $\text{Ph}_2\text{P}(\text{O})\text{CH}_2 \sim \text{CH}_2\text{P}(\text{O})\text{Ph}_2$ was not reduced by SiHCl_3 , in benzene solution in the presence of $p\text{-CH}_3 \cdot \text{C}_6\text{H}_4\text{NEt}_2$, while Si_2Cl_6 in CHCl_3 gave only incomplete reduction. The best reducing system for compounds of type **4** proved to be SiHCl_3 and $n\text{-Bu}_3\text{N}$ in acetonitrile [8].

Reaction 8. Compounds of the type $\text{R}_2\text{P}(\text{OR}')$ are generally prepared by reaction of $\text{R}_2\text{P}\text{Cl}$ and $\text{R}'\text{OH}$ in the presence of a nitrogen base and it is stated that the use of sodium alkoxides is less satisfactory [9]. We find that while $(\text{C}_6\text{H}_{11})_2\text{P}(\text{O}-\text{iso Pr})$ can be obtained from $(\text{C}_6\text{H}_{11})_2\text{P}\text{Cl}$, iso-PrOH and Et_3N , the preparation of $t\text{-Bu}_2\text{P}(\text{O}-\text{iso Pr})$ requires the action of iso-PrONa on $t\text{-Bu}_2\text{P}\text{Cl}$.

Reaction 9. This reaction [10] provides a very convenient method of preparation for compounds of type **4** from $\text{R}_2\text{P}(\text{O}-\text{iso Pr})$ and **1g** even when $\text{R} = t\text{-Bu}$.

The analytical and NMR. data for the new compounds characterized are given in *Tables 1* and *2* respectively.

Table 1. Analytical Data for Ditertiary Phosphines and Their Precursors

Compound		C (%)	H (%)	P (%)	Mol. Wt.
1b ($m\text{-CH}_3 \cdot \text{C}_6\text{H}_4$) ₂ PCH ₂ ~ CH ₂ P($m\text{-CH}_3 \cdot \text{C}_6\text{H}_4$) ₂	Found	84.78	6.15	8.94	668
	Calcd.	84.71	6.22	9.11	681
1c ($p\text{-CH}_3\text{O} \cdot \text{C}_6\text{H}_4$) ₂ PCH ₂ ~ CH ₂ P($p\text{-CH}_3\text{O} \cdot \text{C}_6\text{H}_4$) ₂	Found	77.27	5.80	8.34	794
	Calcd.	77.40	5.68	8.32	745
1d ($m\text{-CF}_3 \cdot \text{C}_6\text{H}_4$) ₂ PCH ₂ ~ CH ₂ P($m\text{-CF}_3 \cdot \text{C}_6\text{H}_4$) ₂	Found	64.40	3.34	7.22	877
	Calcd.	64.30	3.37	6.91	897
1e $(\text{C}_6\text{H}_{11})_2\text{PCH}_2 \sim \text{CH}_2\text{P}(\text{C}_6\text{H}_{11})_2$	Found	81.31	9.03	9.45	629
	Calcd.	81.44	9.00	9.56	649
1f $(t\text{-Bu})_2\text{PCH}_2 \sim \text{CH}_2\text{P}(t\text{-Bu})_2$	Found	79.36	9.29	11.28	537
	Calcd.	79.37	9.25	11.38	545
2c ($p\text{-CH}_3\text{O} \cdot \text{C}_6\text{H}_4$) ₂ P(O)H	Found	64.01	5.78	11.73	
	Calcd.	64.14	5.73	11.83	
2d ($m\text{-CF}_3 \cdot \text{C}_6\text{H}_4$) ₂ P(O)H	Found	49.75	2.69	9.23	
	Calcd.	49.72	2.66	9.17	
2e $(\text{C}_6\text{H}_{11})_2\text{P}(\text{O})\text{H}$	Found	66.36	10.61		
	Calcd.	67.25	10.84		
3d ($m\text{-CF}_3 \cdot \text{C}_6\text{H}_4$) ₂ PH	Found	52.21	2.91	9.53	
	Calcd.	52.19	2.80	9.62	
4b $(m\text{-CH}_3 \cdot \text{C}_6\text{H}_4)_2\text{P}(\text{O})\text{CH}_2$ ~ $\text{CH}_2\text{P}(\text{O})(m\text{-CH}_3 \cdot \text{C}_6\text{H}_4)_2 \cdot 0.25 \text{CHCl}_3$	Found	77.86	5.41		
	Calcd.	78.03	5.73		
4c ($p\text{-CH}_3\text{O} \cdot \text{C}_6\text{H}_4$) ₂ P(O)CH ₂ ~ CH ₂ P(O)($p\text{-CH}_3\text{O} \cdot \text{C}_6\text{H}_4$) ₂	Found	74.26	5.46	8.12	
	Calcd.	72.21	5.44	7.98	
4d ($m\text{-CF}_3 \cdot \text{C}_6\text{H}_4$) ₂ P(O)CH ₂ ~ CH ₂ P(O)($m\text{-CF}_3 \cdot \text{C}_6\text{H}_4$) ₂	Found	62.08	3.29	6.84	
	Calcd.	62.08	3.26	6.68	
4e $(\text{C}_6\text{H}_{11})_2\text{P}(\text{O})\text{CH}_2 \sim \text{CH}_2\text{P}(\text{O})(\text{C}_6\text{H}_{11})_2$	Found	77.05	8.55	8.97	
	Calcd.	77.61	8.59	9.10	
4f $(t\text{-Bu})_2\text{P}(\text{O})\text{CH}_2 \sim \text{CH}_2\text{P}(\text{O})(t\text{-Bu})_2$	Found	75.06	8.74	10.64	
	Calcd.	74.96	8.74	10.75	

Table 2. NMR. Parameters for Diteriary Phosphines and Their Precursors

Compound	$\delta_{1,12^a)}$ [ppm]	δ_{CH_2} [ppm]	δ_{CH_3} [ppm]	$\delta_{=P-H}$ [ppm]	$^1J(^{31}P, ^1H)^b)$ [Hz]	$^2J(^{31}P, ^1H)$ [Hz]	$\delta_{^{31}P^c)}$ [ppm]
1b	8.92(s)	3.70(s)	2.19(s)				- 9.0
1c	8.94(s)	3.63(s)	3.74(s)				- 11.6
1d	9.15(s)	3.75(s)					- 8.6
1e	9.11(s)	3.16(s)					+ 3.8
1f	9.12(s)	3.20(d)	1.20(d) ^{d)}			3.36	+ 35.6
2c			3.78(s)	8.03(d)	475.6		+ 20.5
2d				8.24(d)	492.7		+ 17.8
2e				6.22(d)	433.7		+ 49.5
3d				5.36(d)	219.7		- 40.9
4b	9.41(s)	4.12(d)	2.17(s)			13.80	+ 30.2
4c	9.34(s)	4.05(d)	3.74(s)			13.73	+ 30.3
4d	9.41(s)	4.30(d)				13.20	+ 28.8
4e	9.45(s)	3.52(d)				11.90	+ 49.2
4f	9.60(s)	3.68(d)	1.25(d) ^{e)}			10.68	+ 48.4
5e			1.11(d) ^{f)}				+ 138.5
5f			^{g)}				+ 150.7

a) Shift of protons at positions 1 and 12 of the benzo[c]phenanthrene ring.

b) For H directly bonded to P.

c) Relative to an external H_3PO_4 standard; shifts to higher fields are denoted as negative and those to lower fields as positive.

d) $^3J(^{31}P, ^1H) = 12.29$ Hz.

e) $^3J(^{31}P, ^1H) = 13.12$ Hz.

f) $\delta_{CH} = 3.68$ ppm (doublet of septets); $^3J(^1H, ^1H) = 6.10$ Hz; $^3J(^{31}P, ^1H) = 9.33$ Hz.

g) $\delta_{CH_3(t-Bu)} = 1.09$ (d) ppm; $^3J(^{31}P, ^1H_{(t-Bu)}) = 12.29$ Hz; $\delta_{CH_2(i-Pr)} = 1.22$ (d) ppm; $^3J(^1H_{CH}, ^1H_{CH_3(i-Pr)}) = 6.10$ Hz; $\delta_{CH} = 3.83$ ppm (doublet of septets); $^3J(^{31}P, ^1H) = 1.22$ Hz.

Experimental Part

Melting points were determined using a Büchi melting point apparatus and are uncorrected. 1H - and ^{31}P -NMR. spectra were recorded on a Bruker HX 90 FT spectrometer at frequencies of 90.00 and 36.43 MHz respectively, the 2H -resonance of the deuterated solvent being used as internal lock. 1H -chemical shifts are in ppm with respect to an internal TMS standard whilst the ^{31}P -chemical shifts are in ppm relative to an external H_3PO_4 -standard. Shifts to higher fields are denoted as negative and those to lower fields as positive. ^{31}P -NMR. spectra were obtained under conditions of broad-band 1H -noise decoupling. Elemental analyses and molecular weight determinations were performed by the Micro-analytical Laboratory of the ETH Zürich. All preparations were carried out under a nitrogen atmosphere in dried and de-oxygenated solvents.

1b. - ($m-CH_3 \cdot C_6H_4$)₂PLi was prepared by the addition of 9.0 ml of a 1.67M solution (15.03 mmol) of *n*-BuLi in hexane to a stirred solution of 3.1 g (14.47 mmol) of ($m-CH_3 \cdot C_6H_4$)₂PH [2] in 15 ml THF which had been cooled to 0°. The resulting orange-brown solution was stirred for 1 h at RT., cooled to 0° and treated with a solution of 3.0 g (7.24 mmol) **1g** in 10 ml THF. The resulting light yellow solution was stirred for 2 h at RT. and the solvent evaporated under reduced pressure. The residue was dissolved in 30 ml $CHCl_3$ and the solution washed with 3×20 ml water. The organic layer was dried over $MgSO_4$ and the solution was evaporated under reduced pressure. The resulting oil was washed first with methanol and then with light petroleum (40–60°) leaving a spongy solid which, after drying i.V., gave the product (3.3 g, 67%) as pale yellow powder. This became oily at 140° and melted to a clear liquid from 195 to 200°.

1c. - 6.8 g (5.0 ml, 50.1 mmol) $SiHCl_3$ and 9.1 g (7 ml, 49.0 mmol) *n*-Bu₃N were added to a suspension of 2.0 g (2.57 mmol) **4c** in 100 ml CH_3CN . The mixture was refluxed for 4 h and then left overnight

at RT. The solvent was evaporated under reduced pressure and the waxy residue dissolved in 100 ml CHCl_3 . The solution was washed successively with 80 ml 30% NaOH-solution and 2×100 ml water and dried over MgSO_4 . The yellow oil obtained after evaporation of the solvent was washed with light petroleum (30–60°) and the residual yellow solid was recrystallized from benzene/light petroleum. The product (1.1 g, 57%) melted over the range 145–148°.

Id. - It was prepared and purified as described for compound **1b**. 4.0 g (12.4 mmol) **3d**, 6.1 ml of a 2.04M solution of *n*-BuLi (12.4 mmol) and 2.57 g (6.2 mmol) **1g** gave, after drying i.v., a red-brown spongy solid which was recrystallised from MeOH giving 1.7 g (30%) of the white crystalline product with m.p. 109–110°.

Ie. - It was prepared and purified as described for **1c**. 3.0 g **4e** gave a brown solid which was purified by washing with acetone and recrystallization from benzene/ethanol. The white crystalline product (1.7 g, 60%) had a m.p. 196–199°.

If. - 1.4 g (9.6 mmol) *t*-Bu₂PH [11] was added to a hot solution of 2.0 g (4.83 mmol) **1g** in 40 ml isobutyl methyl ketone. The mixture was refluxed for 6 h and then left overnight. The liquid was decanted, the residue dissolved in 20 ml ethanol and treated with 20 ml 5% NaOH solution. The solvent was decanted from the oily precipitate which was dissolved in 40 ml CHCl_3 . This solution was washed 3 times with 30 ml water, the organic layer dried over MgSO_4 and the solvent evaporated under reduced pressure. The residual spongy solid was recrystallized from ethanol/benzene giving 0.9 g (30%) of the white crystalline product with m.p. 208–210°.

2c. - 19.6 g (76.03 mmol) $(\text{EtO})_2\text{P}(\text{O})\text{H}$ was added slowly to the *Grignard* reagent prepared from 10.5 g (431.89 m-atoms) Mg and 80.0 g (467.72 mmol) *p*- $\text{CH}_3 \cdot \text{C}_6\text{H}_4 \cdot \text{Br}$ in 250 ml ether which had been cooled to 0°. The very viscous solution was refluxed for 2 h and a waxy solid collected at the bottom of the flask. This mixture was hydrolysed with 200 ml 10% HCl-solution followed by 100 ml of water. The ether layer was separated and the residue was extracted with 6 50-ml portions of C_6H_6 . The solid residue was then dissolved in ethanol and this solution extracted with a mixture of C_6H_6 and H_2O . The combined ether and benzene extracts were dried over MgSO_4 , and the solvent evaporated i.v. The pale yellow crude product was purified by recrystallization from acetone/ether and gave 26.6 g (70%) of product with m.p. 115–116° and $\bar{\nu}_{\text{P-H}} = 2340 \text{ cm}^{-1}$.

2d. - It was prepared as described for **2c** from 10.0 g (72.4 mmol) $(\text{EtO})_2\text{P}(\text{O})\text{H}$, 49.0 g (192.1 mmol) *m*- $\text{CF}_3 \cdot \text{C}_6\text{H}_4 \cdot \text{Br}$ and 5.4 g (222.1 m-atoms) Mg and recrystallised from Et_2O by cooling the solution to –60°. The white crystalline product (22.0 g, 90%) had m.p. 62° and $\bar{\nu}_{\text{P-H}} = 2380 \text{ cm}^{-1}$.

2e. - It was prepared as described for **2c** from 20.0 g (144.82 mmol) $(\text{EtO})_2\text{P}(\text{O})\text{H}$, 85.2 g (522.50 mmol) $\text{C}_6\text{H}_{11}\text{Br}$ and 12.6 g (518.26 m-atoms) Mg and recrystallized from hexane. The white solid (18.2 g, 59%) had m.p. 71–72° (Lit. [12]: 73–75°).

3d. - 20.0 g (59.1 mmol) **2d** and 4 g (105.2 mmol) LiAlH_4 in 200 ml Et_2O were refluxed for 15 h. The red-brown solution was cooled to 0° and hydrolysed by the successive addition of 4 ml water, 4 ml 15% NaOH-solution and 12 ml water. The organic layer was separated, dried over Na_2SO_4 and fractionally distilled, the product being collected at 90–92°/0.4 Torr. Yield 10.5 g (54%). $\bar{\nu}_{\text{P-H}} = 2290 \text{ cm}^{-1}$.

4b. - 10 ml of 30% H_2O_2 was gradually added to 1 g of crude **1b** in 30 ml acetone²⁾ at 0°. The reaction mixture was stirred overnight and the solid which had separated was filtered off, dissolved in 20 ml CHCl_3 and chromatographed over alumina using hexane/ CHCl_3 1:1 as eluant. Evaporation of the eluate left a yellow oil which was dissolved in a minimum of ethanol and cooled to –5°. 0.3 g of white crystalline product of m.p. 232–234° were thus obtained.

4c. - 2.0 g *n*-BuLi (31.2 mmol, 20 ml of a 1.56M solution in hexane) was gradually added to a tepid solution of 8.0 g (30.5 mmol) **2c** in 120 ml THF. The dark yellow solution was left at RT. for 2 h and treated with 6.4 g **1g** in 40 ml THF. The solution was refluxed for 2 h, left overnight and the solvent evaporated under reduced pressure. A residue-solution in CHCl_3 was washed with water, dried over MgSO_4 and the solvent evaporated again. The crude product was recrystallized from CHCl_3 /hexane. The crystalline product (8.0 g, 67%) had m.p. 315–322°.

²⁾ The use of H_2O_2 in acetone solution mentioned in this preparation *should be avoided*. Such a mixture is potentially explosive and becomes particularly dangerous after evaporation of the solvent. During the course of subsequent work on a related compound a violent explosion of the oily residue occurred after the acetone had been evaporated. It was later discovered that the explosive nature of H_2O_2 /acetone mixtures had been observed elsewhere, e.g., see A. D. Brewer, Chem. in Britain, 1975, 355.

4d. - It was prepared as described for **4b**. The solvent was evaporated under reduced pressure (*see Footnote to preparation of 4b*) and the oily residue purified by chromatography over alumina using hexane/chloroform mixtures. The waxy solid thus obtained was recrystallized from ethanol by cooling the solution to -5° . 1 g crude **1b** gave 0.7 g of product with m. p. 204-205 $^{\circ}$.

4e. - 2.0 g (4.83 mmol) **1g** and 2.5 g (9.76 mmol) $(C_6H_{11})_2P(O-isoPr)$ were heated together at 140-160 $^{\circ}$ for 2 h, the *iso*-PrBr formed being allowed to distil off. The reaction mixture was then heated to 160-170 $^{\circ}$ for 1 h, cooled and washed with 100 ml Et_2O . The white residual solid was recrystallized from $CHCl_3$ /hexane giving 2.7 g (83%) of product melting from 318 to 322 $^{\circ}$.

4f. - It was prepared and purified as **4e**. 3.6 g (17.56 mmol) *t*-Bu₂P(*O-isoPr*) and 2.5 g (6 mmol) **1g** gave 2.4 g (68.6%) of product melting over the range 270-273 $^{\circ}$.

$(C_6H_{11})_2P(O-isoPr)$ (**5e**). - 18.0 g (77.4 mmol) $(C_6H_{11})_2PCl$, 4.7 g (77.4 mmol) *iso*-PrOH and 7.8 g (77.1 mmol) Et_3N in 50 ml benzene were refluxed for 4 h. After cooling, the solid formed was filtered off and the solution fractionally distilled. The product (15.4 g, 78%) was collected at 110-115 $^{\circ}$ /1.0 Torr and was characterized through its ¹H-NMR. spectrum.

t-Bu₂P(*O-isoPr*) (**5f**). - A solution of *iso*-PrONa, prepared by adding 1.2 g (52.1 m-atoms) Na to 20 ml *iso*-PrOH, was gradually added to 8.9 g (49.3 mmol) *t*-Bu₂PCl in 20 ml benzene. The mixture was refluxed for 14 h, cooled down and the solid filtered off. The solution was fractionally distilled and the product (4.5 g, 43%) was collected at 50 $^{\circ}$ /1.5 Torr. It was characterized through its ¹H-NMR. spectrum.

The authors are indebted to Drs. *L. Maier* and *P. S. Pregosin* for much valuable discussion and to Mr. *G. Balimann* for recording the NMR. spectra.

BIBLIOGRAPHIE

- [1] *F. Bachechi, L. Zambonelli & L. M. Venanzi*, *Helv.* 60, 2815 (1977), and references quoted therein.
- [2] *L. Maier*, 'Primary, Secondary and Tertiary Phosphines', in 'Organic Phosphorus Compounds', Eds. G.M. Kosolapoff and L. Maier, Wiley-Interscience, N.Y., Vol. 1, p. 23, 1972.
- [3] *L. Maier*, loc. cit., pp. 17-20.
- [4] *N.J. DeStefano, D.K. Johnson, R.M. Lane & L. M. Venanzi*, *Helv.* 59, 2674 (1976).
- [5] *L. Maier*, loc. cit., pp. 37-38.
- [6] *H.R. Hays*, *J. org. Chemistry* 33, 3690 (1968).
- [7] *L. Maier*, loc. cit., pp. 45-46.
- [8] *W.S. Knowles, M.J. Sabacky, B.D. Vineyard & D.J. Weinkauff*, *J. Amer. chem. Soc.* 97, 2567 (1975).
- [9] *G.M. Kosolapoff*, 'Organophosphorus Compounds', Wiley, N.Y., 1950, p. 139.
- [10] *R.G. Harvey & E.R. De Sombre*, 'The Michaelis-Arbuzov Reaction', in 'Topics in Phosphorus Chemistry', Eds. M. Grayson and E.J. Griffith, Interscience, N.Y. 1964, Vol. 1, pp. 57-111.
- [11] *L. Maier*, loc. cit., p. 115.
- [12] *M. Epstein & S.A. Buckler*, *Tetrahedron* 18, 1231 (1962).