This article was downloaded by: [Moskow State Univ Bibliote] On: 19 November 2013, At: 13:24 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/gpss20

Heterocyclization of Orthoaminoester and Orthoamino-nitrilethieno[2, 3-c]pyridine: The Facile Synthesis of Fused Pyridothienopyrimidines

E. Kh. Ahmed^a, M. A. Ameen^a & F. F. Abdel-latif^a ^a Chemistry Department, El-Minia University, El-Minia, Egypt Published online: 01 Feb 2007.

To cite this article: E. Kh. Ahmed , M. A. Ameen & F. F. Abdel-latif (2006) Heterocyclization of Orthoaminoester and Orthoamino-nitrile-thieno[2, 3-c]pyridine: The Facile Synthesis of Fused Pyridothienopyrimidines, Phosphorus, Sulfur, and Silicon and the Related Elements, 181:3, 497-510, DOI: <u>10.1080/10426500500263748</u>

To link to this article: http://dx.doi.org/10.1080/10426500500263748

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any

losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <u>http://www.tandfonline.com/page/terms-and-conditions</u>



Heterocyclization of Orthoaminoester and Orthoamino-nitrile-thieno[2,3-*c*]pyridine: The Facile Synthesis of Fused Pyridothienopyrimidines

E. Kh. Ahmed M. A. Ameen F. F. Abdel-latif Chemistry Department, El-Minia University, El-Minia, Egypt

A highly efficient and versatile synthetic approach to the synthesis of pyrido-[4',3':4,5]thieno[2,3-d]pyrimidines (4, 14, 15, 21) and their heterofused (e.g., triazolo-, triazino-, imidazo-, and tetrazolo-,) pyridothienopyrimidines (5–9, 16, 17, 22–24) is described utilizing 2-amino-3-cyano-4,5,6,7-tetrahydro-thieno[2,3-c]pyridine-6-carboxylic acid ethyl ester (2) and diethyl 2-isothio-cyanate-4,5,6,7-tetrahydrothieno[2,3-c]pyridine-3,6-dicarboxylate (10) as starting materials.

 ${\bf Keywords} \ {\rm Cyclization}; heterofused; hydrazine hydrate; or the aminon-itrile$

Orthomino, *o*-isothiocyanate, -ester, and orthoaminocyano thiophene derivatives have been found to be important intermediates in the chemistry of pyrimidine. Therefore, they have attracted much attention during recent years.¹⁻⁹ The resulting pyrimidine derivatives have found a wide application in a variety of synthetic transformations. Various heterocycles have been built to produce the corresponding fused pyrimidines, which show marked biological activities,¹⁰⁻¹³ and these interesting activities have driven our work during the last few years, for which we have been interested in the synthesis of fused pyrimidines.¹⁴⁻¹⁹ The present article follows that line of research by reporting on a new series of reactions leading the formation of fused pyridothieno-pyrimidines starting from the bi-functional *o*-amino-ester and -nitrile compounds.

Received January 12, 2005; accepted May 1, 2005.

Address correspondence to M. A. Ameen, El-Minia University, Chemistry Department, Faculty of Science, El-Minia, 61519 Egypt. E-mail: m_ameen10@yahoo.com

RESULTS AND DISCUSSION

In connection with our interest to study the chemistry of pyrimidines, we have devised the following route for the efficient synthesis of heterofused, 1,2,4-triazoles and 1,2,4-triazine by novel methods. The key compound 2-amino-3-cyano-4,5,6,7-tetrahydro-thieno[2,3-*c*]pyridine-6-carboxylic acid ethyl ester (**2**) was synthesized by the condensation of *N*-carbethoxy-4-piperidone (**1**), sulfer, and malononitrile following the Gewald synthesis.²⁰

The synthesis of model compounds pyridothienopyrimidine **4** and heterofused triazino- and triazolo-derivatives **5–9** required the condensation of compound **2** with triethylorthoformate in the presence of a catalytic amount of acetic anhydride to give 2-ethoxymethyleneamino-3cyano-4,5,6,7-tetrahydrothieno[2,3-c]pyridine-6-carboxylic acid ethyl ester (**3**). The treatment of compound **3** with hydrazine hydrate in ethanol at r.t. afforded the target 3-amino-4-imino-3,4,5,6,7,8hexahydropyrido[4',3':4,5]thieno[2,3-d]pyrimidine-7-carboxylic acid ethyl ester (**4**) in an 85% yield (Scheme 1).



SCHEME 1

The structure of compound **4** was established by ¹H, ¹³C-NMR, and IR spectra as well as elemental analysis. The elemental analysis was consistent with a molecular formula of $C_{12}H_{15}N_5O_2S$. The IR spectroscopy of **4** showed absorption bands at 3300–3200 and 1680 cm⁻¹ related to the NH₂, NH, and carbonyl group, respectively. The ¹H-NMR showed two broad singlets at $\delta_H = 5.60$ and 7.93 ppm, assigned to the NH₂ and NH-imino protons, respectively. The ¹³C-NMR supported the ¹H-NMR spectroscopy by the appearance of three signals at $\delta_C = 151.31$, 154.62, and 156.09 ppm, corresponding to C-2, C=NH, and C=O present in compound **4**.

Interestingly, on condensation of **4** with triethylorthoformate, the corresponding 9,10-dihydropyrido[4'3',4,5]thieno[2,3-d]-1,2,4-triazolo[3,2-f]pyrimidine-8(7H)-carboxylic acid ethyl ester (**5**) was obtained in an 84% yield. Similarly, the triazolo derivatives **6**, **7**, and **8** were prepared by the interaction of **4** with *p*-chloro-benzaldehyde, acetic anhydride, and phenyl isothiocyanate, respectively. On the other hand, the 1,2,4-triazino moiety was built in compound **9** by the treatment **4** with chloroacetyl chloride in DMF catalyzed by triethylamine (Scheme 2).



SCHEME 2 Reagents and conditions: i: $CH_2(CN)_2$, S, morpholine, EtOH. r.t.; ii: $CH(OEt)_3$, Ac_2O , reflux 8 h; iii: $N_2H_4 \cdot H_2O$, EtOH, stirring 2 h; iv: $CH(OEt)_3$, Ac_2O , reflux 3 h; v: P-chlorobenzaldehyde, AcOH, reflux 3 h; vi: Ac_2O , reflux 3 h; vii: PhNCS, pyridine, reflux 4 h; viii: $ClCH_2COCl$, DMF, Et_3N , stirring 5 h.

The ¹H-NMR and IR were used to identify the 1,2,4-triazolo compounds **5–8**; neither IR or ¹H-NMR spectra supported the presence of the NH and NH₂ group protons that existed in **4**. Moreover, the ¹H NMR spectra revealed two singlet signals at $\delta_{\rm H} = 9.42$ and 2.68 ppm, which were assigned to the triazolo proton in compound **5** and CH₃protons in compound **7**, respectively. The ¹H-NMR spectra showed two double-doublets and a multiplet at $\delta_{\rm H} = 7.63$, 7.95, and 7.11–7.52 ppm, corresponding to aromatic protons related to compounds **6** and **8**, respectively.

Promoted by the presence of the bifunctional *ortho*-amino and -ester groups, which is considered as a precursor for the preparation of the thioxopyrimidinone derivative **12** by the reaction of diethyl 2-isothiocyanate-4,5,6,7-tetrahydrothieno[2,3-*c*]pyridine-3,6-dicarboxylate (**10**)¹⁴ with aniline in dichloromethane to give the openform **11**. The latter was subjected to ensuing cyclization in sodium ethoxide solution, yielding the target compound **12**. *S*-methylation of compound **12** with methyl iodide afforded the target product 3-phenyl-2-thiomethyl-4-oxo-3,4,5,6,7,8-hexahydropyrido[4',3':4,5]thieno[2,3-*d*]-pyrimidine-7-carboxylic acid ethyl ester (**13**) (Scheme 3).



SCHEME 3 Reagents and conditions: i: PhNH₂; CH_2Cl_2 , stirring 2 h; ii: EtONa, reflux 1 h; iii: CH_3I , 1N NaOH, stirring 2 h; iv: N_2H_4 · H_2O , EtOH, reflux 15 h; v: HOCH₂CH₂NH₂, 15 h; vi: NaNO₂, HCl, AcOH, stirring; vii: POCl₃, reflux 14 h.

Treating the S-methylated pyridothienopyrimidine **13** with excess hydrazine hydrate afforded the 2-hydrazino compound **14**, which is considered a good starting material for the synthesis of the tetracyclic ring system **16** via a diazotization reaction. The same approach to obtaining 3-phenyl-2-[1-hydroxyethan]-4-oxo-3,4,5,6,7,8-hexahydropyrido[4',3':4,5]thieno[2,3-d]pyrimidine-7-carboxylic acid ethyl ester (**15**) can be achieved by the treatment of compound **13** with ethanolamine. The open structure **15** underwent direct cylization to build the imidazo moiety **17** (Scheme 3).

The structure of compound 13 was proved by the help of the ¹H-NMR, ¹³C-NMR, and IR spectra as well as elemental analysis. Elemental analysis of 13 was in accordance to its molecular formula as C₁₉H₁₉N₃O₃S₂. The ¹H-NMR spectrum of **13** showed a triplet and quartet corresponding to ester protons. The methyl protons were found to resonate as a singlet at $\delta_{\rm H} = 2.49$ ppm, whereas the aromatic protons appeared at $\delta_{\rm H} = 7.27 - 7.55$ ppm. The $^{13}{
m C}$ -NMR supported the ¹H-NMR spectrum by showing three signals at $\delta_{\rm C} = 15.21, 127.11, 128.11, 129.87, \text{ and } 130.98 \text{ ppm corresponding}$ to SCH_3 and (Ar-C) present in compound 13 (see the Experimental section). The IR spectrum of 13 showed the absorption of the CH-arom. and the carbonyl groups at 3000 and 1710-1680 cm⁻¹, respectively. Combining these spectral and analytical data, compound 13 was identified as 3-phenyl-2-thiomethyl-4-oxo-3,4,5,6,7,8-hexahydropyrido[4',3':4,5]thieno[2,3-d]pyrimidine-7-carboxylic acid ethyl ester.

Analogously, compound **2** was proved to be a key intermediate for other subsequent conversions. Thus, when it was allowed to react with carbon disulfide in alcoholic KOH, the reaction produced compound **18**. This was triturated with methyl iodide to yield ethyl-2,4-bithiomethyl-5,8-dihydropyrido[4',3' :4,5]thieno[2,3-*d*]pyrimidine-7(6*H*)-carboxylate (**19**), which was confirmed easily by the ¹H-NMR spectrum that exhibited a singlet for the two CH₃-protons at $\delta_{\rm H} = 2.61$ ppm (Scheme 4).

Surprisingly, on reacting **19** with excess hydrazine hydrate in the presence of pyridine as a solvent to synthesize the 2,4dihydrazino compound, the process removed only one $-SCH_3$ by $-NHNH_2$ in ethyl-4-hydrazino-2-thiomethyl-5,8-dihydropyrido[4',3' :4,5]thieno[2,3-d]pyrimidine-7(6H)-carboxylate (**21**). The structure of **21** was synthetically proved by its cyclization using triethylorthoformate and carbon disulfide to give **22** and **23**. On reducing **22** by Raney Ni in ethanol, we isolated our known compound ethyl-7,10dihydropyrido[4',3':4,5]thieno[2,3-d]-1,2,4-triazolo[2,1-f]pyrimidine-8(9H)-carboxylate (**24**).¹³ The aforementioned synthetic proof indicated the prevalence of SCH_3 group in position 2 without change in compound **21** (Scheme 4).

In conclusion and by the results in hand, we might serve in the synthesis new classes of fused pyridothienopyrimidines using efficient, facile, and elegant methods of preparation.



SCHEME 4 Reagents and conditions: i: CS_2 , alc. KOH, reflux 5 h; ii: CH_3I , 1N NaOH, stirring 1 h; iii: N_2H_4 · H_2O , pyridine, reflux 4 h; iv: $CH(OEt)_3$, reflux 4 h; v, iv: CS_2 , KOH, CH_3OH , reflux 3 h; vi: raney Ni, ethanol, reflux 2 h.

EXPERIMENTAL

Melting points were determined on a Boetius melting point apparatus and are uncorrected. Elemental analyses were performed on a Carol Erba CHN-S Elemental Analyzer 1108. The ¹³C and ¹H NMR spectra were obtained using sa Bruker AC 300 (¹H: 300.13 MHz, ¹³C :75.5 MHz). The solvents were deuterated dimethyl sulfoxide and chloroform. The δ -values are given in ppm and the internal standard was tetra-methylsilane. The IR spectra were recorded on a Nicolet 250 FT-IR spectrophotometer as potassium bromide pellets.

2-Ethoxymethyleneamino-3-cyano-4,5,6,7-tetrahydrothieno[2,3-c]pyridine-6-carboxylic Acid Ethyl Ester (3)

A mixture of **2** (2.51 g, 10 mmol) with a few drops of acetic anhydride in triethylorthoformate (20 mL) was refluxed for 8 h. The separated crystalline product was obtained on cooling at r.t. Yield: 2.4 g (80%), colorless plates (ethanol), m.p. 90–92°C, IR (KBr): $v^{-} = 2210$ (CN), 1715 (ester C=O), 1618 (C=N) cm⁻¹, ¹H NMR (DMSO-d₆): $\delta = 1.21$ (t, 3H, COOCH₂CH₃), 1.35 (t, 3H, OCH₂CH₃), 2.85 (t, 2H, H-4), 3.65 (t, 2H, H-5), 4.11 (q, 2H, COOCH₂CH₃), 4.35 (q, 2H, OCH₂CH₃), 4.55 (s, 2H, H-7), 8.55 (s, 1H, CH), ¹³C NMR (DMSO-d₆): $\delta = 14.38$ (CH₃), 14.38 (CH₃), 23.39 (C-4), 40.93 (C-5), 42.02 (C-7), 60.86 (CH₂), 60.97 (CH₂), 113.38 (CN), 123.99 (C-3), 129.60 (C-7a), 146.09 (C-3a), 154.55 (C=O), 159.07 (CH=N), 163.58 (C-2).

3-Amino-4-imino-3,4,5,6,7,8-hexahydropyrido[4',3':4,5] thieno[2,3-*d*]pyrimidine-7-carboxylic Acid Ethyl Ester (4)

To a stirred suspension of **3** (6.1 g, 20 mmol) in absolute ethanol (40 mL), hydrazine hydrate (1.5 g, 30 mmol) was added. The reaction mixture was stirred at r.t. for 2 h. The precipitate that formed was filtered off, washed with cold ethanol, and recrystalized from dioxane to produce white needles. Yield: 4.9 g (85%), m.p. 152–154°C, (Found: C, 49.02; H, 5.01; N, 23.63; S, 10.91. Calc. for $C_{12}H_{15}N_5O_2S$ (293.35): C, 49.13; H, 5.15; N, 23.87; S, 10.73), IR (KBr): v = 3360-3280 (NH₂, NH), 1708 (ester C=O), 1615 (C=N) cm⁻¹, ¹H NMR (DMSO-d₆): $\delta = 1.21$ (t, J = 7 Hz, 3H, COOCH₂CH₃), 3.00 (t, J = 5.6 Hz, 2H, H-5), 3.65 (t, J = 5.6 Hz, 2H, H-6), 4.11 (q, J = 7 Hz, 2H, COOCH₂CH₃), 4.55 (s, 2H, H-8), 5.60 (b, 2H, NH₂), 7.93 (b, 1H, NH), 8.35 (s, 1H, CH), ¹³C NMR (DMSO-d₆): $\delta = 14.42$ (CH₃), 26.09 (C-5), 40.87 (C-6), 42.79 (C-8), 60.92 (CH₂), 119.68 (C-4a), 127.95 (C-4b), 129.84 (C-8a), 148.22 (C-2), 151.31 (C-9a), 154.62 (C=NH), 156.09 (C=O).

9,10-Dihydropyrido[4',3':4,5]thieno[2,3-*d*]-1,2,4-triazolo [3,2-*f*]pyrimidine-8(7*H*)-carboxylic Acid Ethyl Ester (5)

Compound 4 (0.59 g, 2 mmol) in triethylorthoformate (5 mL) was heated under reflux for 3 h. The solid product that formed was collected and recrystalized from methanol to produce white needles. Yield: 0.5 g (84%), m.p. 160–161°C, IR (KBr): v = 1700 (ester C=O), 1610 (C=N) cm⁻¹, ¹H NMR (DMSO-d₆): $\delta = 1.24$ (t, 3H, COOCH₂CH₃), 3.17 (t, 2H, H-10), 3.69 (t, 2H, H-9), 4.18 (q, 2H, COOCH₂CH₃), 4.68 (s, 2H, H-7), 9.12 (s, 1H, H-4), 9.42 (s, 1H, H-2).

2-(4-Chlorophenyl)-9,10-dihydropyrido[4',3':4,5]thieno[2,3-*d*]-1,2,4-triazolo[3,2-*f*]pyrimidine-8(7*H*)-carboxylic Acid Ethyl Ester (6)

A mixture of 4 (1.47 g, 5 mmol) and *p*-chlorobenzaldehyde (0.7 g, 5 mmol) in 15 mL of acetic acid was heated under reflux for 3 h. The resulting precipitate was collected and crystallized from dioxane to produce pale yellow needles. Yield: 1.8 g (90%), m.p. 175–176°C, IR (KBr): v = 3090 (arom. CH), 1705 (ester C=O), 1605 (C=N) cm⁻¹, ¹H NMR (DMSO-d₆): $\delta = 1.29$ (t, 3H, COOCH₂CH₃), 3.15 (t, 2H, H-10), 3.85 (t, 2H, H-9), 4.18 (q, 2H, COOCH₂CH₃), 4.75 (s, 2H, H-7), 7.63 (d, 2 ArH), 7.95 (d, 2 ArH), 9.12 (s, 1H, H-4).

2-Methyl-9,10-dihydropyrido[4',3':4,5]thieno[2,3-*d*]-1,2,4triazolo[3,2-*f*]pyrimidine-8(7*H*)-carboxylic Acid Ethyl Ester (7)

A solution of 4 (0.59 g, 2 mmol) in acetic anhydride (5 mL) was refluxed for 3 h. The reaction mixture was allowed to cool and poured in ice water. The precipitate that formed was filtered off, washed with cold water, and recrystalized from ethanol to produce white needles. Yield: 0.56 g (88%), m.p. 181–182°C, IR (KBr): v = 2950 (aliph. CH), 1705 (ester C=O), 1610 (C=N) cm⁻¹, ¹H NMR (DMSO-d₆): $\delta = 1.30$ (t, 3H, COOCH₂CH₃), 2.68 (s, 3H, CH₃), 3.27 (t, 2H, H-10), 3.87 (t, 2H, H-9), 4.20 (q, 2H, COOCH₂CH₃), 4.83 (s, 2H, H-7), 9.09 (s, 1H, H-4).

2-(Phenylamino)-9,10-dihydropyrido[4',3':4,5]thieno[2,3-*d*]-1,2,4-triazolo[3,2-*f*]pyrimidine-8(7*H*)-carboxylic Acid Ethyl Ester (8)

A mixture of 4 (0.59 g, 2 mmol) and phenyl isothiocyanate (0.27 g, 2 mmol) in pyridine (10 mL) was refluxed for 4 h. On cooling, the

formed solid product was collected by filtration. Yield: 0.57 g (72%), yellow crystals (ethanol), m.p. 200–202°C, IR: (KBr) $\upsilon = 3075$ (arom. CH), 1708 (ester C=O), 1620 (C=N) cm⁻¹, ¹H NMR (DMSO-d_6): $\delta = 1.18$ (t, 3H, COOCH₂CH₃), 3.05 (t, 2H, H-10), 3.65 (t, 2H, H-9), 4.16 (q, 2H, COOCH₂CH₃), 4.64 (s, 2H, H-7), 7.11–7.52 (m, 5ArH), 9.18 (s, 1H, H-4), 10.45 (s, 1H, NH).

4-Oxo-2,3,4,8,10,11-hexahydropyrido[4',3':4,5]thieno[2,3-*d*]-1,2,4-triazolo[3,2-*f*]pyrimidine-9-carboxylic Acid Ethyl Ester (9)

To a mixture of **4** (0.293 g, 1 mmol) and chloroacetyl chloride (0.17 g, 1.5 mmol) in DMF (5 mL), triethylamine (0.101 g, 1 mmol) was added. The reaction mixture was stirred at r.t. for 5 h and then poured into ice water. The precipitate that formed was filtered off, washed with water, and recrystalized from ethanol to produce pale yellow needles, 0.25 g (75%), m.p. 139–140°C, IR (KBr): v = 3280 (NH), 1710 (ester C=O), 1665 (C=O), 1615 (C=N) cm⁻¹, ¹H NMR (DMSO-d₆): $\delta = 1.18$ (t, 3H, COOCH₂CH₃), 3.14 (t, 2H, H-11), 3.65 (t, 2H, H-10), 4.14 (q, 2H, COOCH₂CH₃), 4.30 (s, 2H, C-2), 4.67 (s, 2H, H-8), 9.15 (s, 1H, H-5), 10.83 (s, 1H, NH).

Diethyl-2-(3-phenyl-thioureido)-4,7-dihydro-5*H*-thieno[2,3-*c*]-pyridine-3,6-dicarboxylate (11)

Isothiocyanate compound **10** (3.4 g, 10 mmol) was dissolved in dichloromethane (30 mL) and then an equivalent amount of aniline was added. The mixture was stirred at r.t. for 2 h. The solid product that precipitated was filtered off and recrystalized from methanol to produce white crystals, 3.9 g (90%), m.p. 179–181°C, IR (KBr): v = 3270 (NH), 3080 (arom. CH), 1705 (ester C=O), 1680 (ester C=O) cm⁻¹, ¹H NMR (DMSO-d₆): $\delta = 1.26$ (t, 6H, 2COOCH₂CH₃), 2.85 (t, 2H, H-4), 3.65 (t, 2H, H-5), 4.11 (q, 4H, 2COOCH₂CH₃), 4.55 (s, 2H, H-7), 7.26–7.56 (m, 5H, ArH), 9.55 (s, 1H, NH), 11.02 (s, 1H, NH).

Ethyl-4-oxo-3-phenyl-2-thioxo-1,2,3,4,5,,8-hexahydropyrido-[4',3':4,5]thieno[2,3-d]pyrimidine-7(6H)-carboxylate (12)

A mixture of **11** (4.3 g, 10 mmol) and a solution of sodium ethoxide (0.3 g sodium in 30 mL absolute ethanol) was refluxed for 1 h. After removing the solvent, the residue was diluted with water and acidified with HCl. The product was collected by filtration, washed with water, dried and recrystallized from DMF/n hexane to produce white crystals. Yield: 3.6 g (92%), m.p. 273–275°C, IR (KBr): v = 3250 (NH), 3040 (arom. CH), 1708

(ester C=O) cm⁻¹, ¹H NMR (DMSO-d₆): δ = 1.21 (t, 3H, COOCH₂CH₃), 2.95 (t, 2H, H-5), 3.72 (t, 2H, H-6), 4.14 (q, 2H, COOCH₂CH₃), 4.63 (s, 2H, H-8), 7.22–7.54 (m, ArH), 10.50 (s, H, NH).

Ethyl-4-oxo-3-phenyl-2-thiomethyl-3,4,5,8-tetrahydropyrido-[4',3':4,5]thieno[2,3-*d*]pyrimidine-7(6*H*)-carboxylate (13)

To a suspension of **12** (1.16 g, 3 mmol) in 1*N* sodium hydroxide (12 mL), methyl iodide (0.72 g, 5 mmol) was added. The reaction mixture was stirred at r.t. for 2 h. The solid was precipitated, collected, washed with water, and dried. Yield: 1.14 g (95%), colorless crystals (methanol), m.p. 160–161°C, (Found: C, 56.64; H, 4.59; N, 10.31; S, 15.94. Calc. for $C_{19}H_{19}N_3O_3S_2$ (401.51): C, 56.84; H, 4.77; N, 10.47; S, 15.81), IR (KBr): v = 3060 (arom. CH), 1710 (ester C=O), 1680 (C=O) cm⁻¹, ¹H NMR (DMSO-d₆): $\delta = 1.30$ (t, 3H, COOCH₂CH₃), 2.49 (s, 3H, SCH₃), 3.05 (t, 2H, H-5), 3.75 (t, 2H, H-6), 4.21 (q, 2H, COOCH₂CH₃), 4.68 (s, 2H, H-8), 7.27–7.55 (m, 5ArH), ¹³C NMR (DMSO-d₆): $\delta = 14.23$ (CH₃), 15.21 (SCH₃), 24.87 (C-5), 40.21 (C-6), 41.20 (C-8), 60.79 (CH₂), 110.94 (C-4a), 114.63 (C-4b), 121.45 (C-8a), 127.11, 128.11, 129.87, 130.98 (Ar-C), 131.69 (C-9a), 155.22 (C-2), 157.12 (C=O), 164.86 (C=O).

Synthesis of 14 and 15

To a stirred suspension of 13 (10 mmol) in absolute ethanol (10 mL), hydrazine hydrate 80% or amino ethanol (5 mL) was added. The reaction mixture was heated under reflux for 15 h. Excess solvent was removed and the residue was poured into water. The precipitates that formed were filtered off, washed with cold ethanol, and recrystalized from methanol.

Ethyl-2-hydrazino-4-oxo-3-phenyl-3,4,5,,8-tetrahydropyrido-[4',3':4,5]thieno[2,3-*d*]pyrimidine-7(6*H*)-carboxylate (14)

Yield: 2.1 g (55%), m.p. 112–113°C, IR (KBr): $\upsilon = 3370-3280$ (NH₂, NH), 3070 (arom. CH), 1710 (ester C=O), 1680 (C=O) cm⁻¹, ¹H NMR (DMSO-d₆): $\delta = 1.28$ (t, 3H, COOCH₂CH₃), 2.65 (b, 2H, NH₂), 3.07 (t, 2H, H-5), 3.52 (t, 2H, H-6), 4.17 (q, 2H, COOCH₂CH₃), 4.64 (s, 2H, H-8), 5.26 (b, H, NH), 7.25–7.54 (m, 5H, ArH).

Ethyl-2-(hydroxyethylamino)-4-oxo-3-phenyl-3,4,5,,8tetrahydro-pyrido[4',3':4,5]thieno[2,3-*d*]pyrimidine-7(6*H*)-carboxylate (15)

Yield: 1.6 g (40%), m.p. 88–90°C, IR (KBr): v = 3430-3300 (OH), 3230 (NH), 3080 (arom. CH), 2920 (aliph. CH), 1709 (ester C=O) cm⁻¹, ¹H

NMR (DMSO-d₆): $\delta = 1.29$ (t, 3H, COOCH₂CH₃), 2.51 (b, H, OH), 2.97 (t, 2H, H-5), 3.47 (t, 2H, H-6), 3.54–3.71 (m, 4H, 2CH₂), 4.16 (q, 2H, COOCH₂CH₃), 4.58 (s, 2H, H-8), 4.63 (s, H, NH), 7.26–7.58 (m, 5H, ArH).

Ethyl-5-oxo-4-phenyl-4,5,,6,9-tetrahydropyrido[4',3':4,5] thieno[2,3-*d*]tetrazolo[5,1-*b*]pyrimidine-8(7*H*)- carboxylate (16)

To a solution of **14** (0.39 g, 1 mmol) in conc. HCl (5 mL) and glacial acetic acid (5 ml) at 0°C, a cold solution of (10%) sodium nitrite (7 mL, 10 mmol) was added with stirring for 10 min. The formed precipitate was filtered, washed with water, and recrystalized from ethanol to produce buff crystals. Yield: 0.25 g (62%), m.p. 205–207°C, IR (KBr): v = 3065 (arom. CH), 1712 (ester C=O), 1675 (C=O) cm⁻¹,¹H NMR (DMSO-d₆): $\delta = 1.21$ (t, 3H, COOCH₂CH₃), 2.89 (t, 2H, H-6), 3.53 (t, 2H, H-7), 4.13 (q, 2H, COOCH₂CH₃), 4.55 (s, 2H, H-9), 7.25–7.51 (m, 5H, ArH).

Ethyl-5-oxo-4-phenyl-1,2,4,5,,6,9-hexahydropyrido[4',3':4,5] thieno[2,3-*d*]imidazo[2,1-*b*]pyrimidine-8(7*H*)carboxylate (17)

A mixture of compound **15** (0.42 g, 1 mmol) and phosphoryl chloride (10 mL) was refluxed for 14 h. Excess solvent was removed and the residue was poured into water. After extraction with chloroform, the organic layer was washed with sodium bicarbonate and water, and then dried over anhydrous sodium sulfate. The solvent was evaporated under vacuum. Yield: 0.14 g (35%), yellow crystals (methanol/n-hexane), m.p. 132–133°C, IR (KBr): v = 3085 (arom. CH), 1707 (ester C=O) cm⁻¹, ¹H NMR (DMSO-d₆): $\delta = 1.24$ (t, 3H, COOCH₂CH₃), 2.86 (t, 2H, H-6), 3.65 (t, 2H, H-7), 3.85–4.05 (m, 4H, 2CH₂), 4.17 (q, 2H, COOCH₂CH₃), 4.55 (s, 2H, H-9), 7.28–7.59 (m, 5H, ArH).

Ethyl-2,4-dithioxo-1,2,3,4,5,8-hexahydropyrido[4',3':4,5]thieno [2,3-*d*]pyrimidine-7(6*H*)-carboxylate (18)

To suspension of **2** (2.5 g, 10 mmol) in 10 % alcoholic potassium hydroxide (15 mL), 5 mL of carbon disulfide was added. The mixture was refluxed for 5 h. Excess solvent was removed. The residue diluted with water and acidified with HCl. The product was filtered off, washed with water, dried and recrystallized from DMF/water to produce yellow crystals, 3 g (93%), m.p. 251–252°C, IR (KBr): v = 3270 (NH), 1710 (ester

C=O), 1612 (C=N) cm⁻¹,¹H NMR (DMSO-d₆): δ = 1.21 (t, J = 7 Hz, 3H, COOCH₂CH₃), 3.07 (t, J = 5.6 Hz, 2H, H-5), 3.62 (t, J = 5.6 Hz, 2H, H-6), 4.06 (q, J = 7 Hz, 2H, COOCH₂CH₃), 4.53 (s, 2H, H-8), 13.25 (bs, 2H, 2NH), ¹³C NMR (DMSO): δ = 14.37 (CH₃), 27.22 (C-5), 40.92 (C-6), 42.58 (C-8), 60.93 (CH₂), 115.33 (C-4a), 126.19 (C-4b), 130.73 (C-8a), 148.70 (C-9a), 154.70 (C=O) 169.21 (C=S), 181.27 (C=S)

Ethyl-2,4-dithiomethyl-5,8-dihydropyrido[4',3':4,5]thieno[2,3d]pyrimidine-7(6H)-carboxylate (19)

To a suspension of **18** (0.98 g, 3 mmol) in 1*N* sodium hydroxide (12 mL), methyl iodide (1.44 g, 10 mmol) was added. The reaction mixture was stirred at r.t. for 1 h. The solid was precipitated, filtered off, washed with water, and dried. Yield: 0.92 g (86%), orange crystals (methanol). m.p. 134–135°C, IR (KBr): v = 1710 (ester C=O), 1615 (C=N) cm⁻¹, ¹H NMR (DMSO-d₆): $\delta = 1.21$ (t, 3H, COOCH₂CH₃), 2.61 (s, 6H, 2CH₃), 2.96 (t, 2H, H-5), 3.68 (t, 2H, H-6), 4.06 (q, 2H, COOCH₂CH₃), 4.64 (s, 2H, H-8).

Ethyl-4-hydrazino-2-thiomethyl-5,8-dihydropyrido[4',3':4,5] thieno[2,3-*d*]pyrimidine-7(6*H*)-carboxylate (21)

To a stirred suspension of **19** (3.5 g, 10 mmol) in pyridine (30 mL), hydrazine hydrate 80% (2 mL) was added. The reaction mixture was heated under reflux for 4 h. The precipitate that formed after cooling was filtered off, washed with cold ethanol, and recrystalized from methanol to produce gray needles. Yield: 2.2 g (65%), m.p. 176–177°C, (Found: C, 44.38; H, 5.17; N, 30.19; S, 9.89. Calc. for $C_{12}H_{17}N_7O_2S$ (323.38): C, 44.57; H, 5.30; N, 30.32; S, 9.68), IR (KBr): v = 3380-3270 (NH₂, NH), 1702 (ester C=O), 1630 (C=N) cm⁻¹,¹H NMR (DMSO-d₆): $\delta_{=}1.21$ (t, 3H, COOCH₂CH₃), 2.56 (s, 3H, SCH₃), 2.61 (b, 2H, NH₂), 2.85 (b, 1H, NH), 3.08 (t, 2H, H-5), 3.78 (t, 2H, H-6), 4.21 (q, 2H, COOCH₂CH₃), 4.85 (s, 2H, H-8).

Ethyl-4-thiomethyl-7,10-dihydropyrido[4',3':4,5]thieno[2,3-*d*]-1,2,4-triazolo[2,1-*f*]pyrimidine-8(9*H*)-carboxylate (22)

A mixture of compound **21** (0.65 g, 2 mmol) and triethylorthoformate (10 mL) was refluxed for 4 h. After cooling at r.t., the formed product was filtered off, dried, and recrystallized from ethanol to produce colorless crystals. Yield: 0.50 g (72%). m.p. 192–194°C, IR (KBr): v = 1706 (ester C=O), 1613 (C=N) cm⁻¹, ¹H NMR (DMSO-d₆): $\delta = 1.30$ (t, 3H,

COOCH₂CH₃), 2.81 (s, 3H, SCH₃), 3.29 (t, 2H, H-10), 3.85 (t, 2H, H-9), 4.22 (q, 2H, COOCH₂CH₃), 4.77 (s, 2H, H-7), 8.79 (s, H, CH).

Ethyl-3-thioxo-4-thiomethyl-2,3,7,10tetrahydropyrido[4',3':4,5]thieno[2,3-*d*]-1,2,4-triazolo[2,1f]pyrimidine-8(9*H*)-carboxylate (23)

To a mixture of **21** (0.65 g, 2 mmol) and KOH (0.112 g, 2 mmol) in methanol (10 mL), carbon disulfide (1 mL) was added. The mixture was refluxed for 3 h. After removing the solvent on vacuum, the residue was diluted with water and acidified with HCl. The solid product was filtered off, washed with water, dried, and recrystalized from DMF/water to produce yellow crystals. Yield: 0.47 g (62%), m.p. 285–287°C, IR (KBr): v = 3220 (NH), 1705 (ester C=O), 1610 (C=N) cm⁻¹, ¹H NMR (DMSO-d₆): $\delta = 1.28$ (t, 3H, COOCH₂CH₃), 2.71 (s, 3H, SCH₃), 3.24 (t, 2H, H-10), 3.82 (t, 2H, H-9), 4.20 (q, 2H, COOCH₂CH₃), 4.73 (s, 2H, H-7), 9.86 (b, 1H, NH).

Ethyl-7,10-dihydropyrido[4',3':4,5]thieno[2,3-d]-1,2,4triazolo[2,1-f]pyrimidine-8(9H)-carboxylate (24)

To a solution of compound **22** (0.7 g, 2 mmol) in ethanol (10 mL), Raney nickel catalyst (0.2 g) was added. The mixture was gently refluxed under a hydrogen atmosphere for 2 h. The catalyst was removed by filtration. The solvent was removed under vacuum to give white needles in a 61% yield. (M.p. 212–214°C Lit.¹³)

REFERENCES

- M. Modica, M. Santagati, A. Santagati, V. Cutuli, N. Mangno, and A. Caruso, *Pharmazie*, 55, 500 (2000).
- [2] Y.-W. Ho, J. Chin Chem Soc., 48, 1163 (2001).
- [3] C. G. Dave, A. B. Shah, and H. C. Shah, J. Heterocycl Chem., 34, 937 (1997).
- [4] H. Vicueg, S. Leistner, N. Bohm, J. Prantz, and G. Wagnez, Pharmazie, 48, 26 (1993).
- [5] O. A. Abd-allah, Phosphorus, Sulfur and Silicon, 178, 1115 (2003).
- [6] R. Bohmen, and R. Pech, Pharmazie, 39, 19 (1984).
- [7] S. Leistner, U. Krasselt, S. Dumke, and G. Wagner, Pharmazie, 47, 682 (1992).
- [8] H. S. El-Kadhef, A. A. Geies, A. M. Kamal, and A. A. Abdel-Hafez, J. Chem. Technol. Biotechnol., 57, 15 (1993).
- [9] H. Marouko, K. Yamagata, and M. Yamazaki, J. Heterocyclic Chem. 38, 269 (2001).
- [10] E. Bousquet, G. Romero, F. Guerrera, A. Caruso, and M. A. Roxas, *Farmaco Ed. Sci.*, 40, 869 (1985).
- [11] E. Bousquet, F. Guerrera; N. A. Siracusa, A. Caruso, and M. A. Roxas, *Farmaco Ed Sci.*, 39, 110 (1984).

- [12] C. G. Dave, P. R. Shah, C. K. Dave, and J. V. Patel, J. Indian Chem. Soc., 66, 48 (1989).
- [13] E. Kh. Ahmed, A. M. N. Gohar, and M. A. Ameen, Pharmazie, 55, 31 (2000).
- [14] E. Kh. Ahmed, U. Sensfuss, and W. D. Habicher, Heterocyclic Chem., 36, 1119 (1999).
- [15] E. Kh. Ahmed, Heteroatom Chemistry, 13, 280 (2002).
- [16] E. Kh. Ahmed, Phosphorus, Sulfur and Silicon, 177, 989 (2002).
- [17] E. Kh. Ahmed, Phosphorus, Sulfur and Silicon, 177, 1323 (2002).
- [18] E. Kh. Ahmed, Phosphorus, Sulfur and Silicon, 178, 1 (2003).
- [19] E. Kh. Ahmed, Heteroatom Chemistry, 14, 201 (2003).
- [20] K. Gewald, E. Schinke, and H. Bottcher, Chem. Ber., 99, 94 (1966).