

This article was downloaded by: [George Mason University]

On: 21 December 2014, At: 16:25

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides and Nucleotides

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/Incn19>

Enantioselective Synthesis and Conformational Analysis of Cyclohexene Carbocyclic Nucleosides

Jing Wang^a & Piet Herdewijn^a

^a Laboratory of Medicinal Chemistry, Rega Institute for Medical Research, K. U. Leuven, Minderbroedersstraat 10, B-3000, Leuven, Belgium

Published online: 04 Oct 2006.

To cite this article: Jing Wang & Piet Herdewijn (1999) Enantioselective Synthesis and Conformational Analysis of Cyclohexene Carbocyclic Nucleosides, *Nucleosides and Nucleotides*, 18:4-5, 593-594, DOI: [10.1080/15257779908041505](https://doi.org/10.1080/15257779908041505)

To link to this article: <http://dx.doi.org/10.1080/15257779908041505>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>


ENANTIOSELECTIVE SYNTHESIS AND CONFORMATIONAL ANALYSIS OF CYCLOHEXENE CARBOCYCLIC NUCLEOSIDES

Jing Wang and Piet Herdewijn*

Laboratory of Medicinal Chemistry, Rega Institute for Medical Research, K.U.Leuven, Minderbroedersstraat 10, B-3000 Leuven, Belgium


ABSTRACT: An enantioselective approach towards the synthesis of optically pure cyclohexene nucleosides **3** has been developed starting from (*R*)-carvone. The key steps are the regio- and stereoselective hydroboration of an exo double bond, the selective reduction of an enone intermediate and introduction of a base moiety by Mitsunobu reaction. Conformational analysis showed that the adenine base adopts predominantly in a pseudo-axial position.

Carbocyclic nucleosides are of broad interest as antiviral agents and as building blocks for oligonucleotides due to their better chemical and enzymatic stabilities, as compared to nucleosides. The difference is due to the replacement of the ring-oxygen atom by a methylene group, hence, to the loss of the anomeric center. In contrast to five-membered ring carbocyclic nucleosides, little is known about the conformational behaviour and biological activity of six-membered ring analogues.




Previous research on six-membered carbohydrate nucleosides such as **1** and their carbocyclic congeners **2** (R = H) has shown that these two series of nucleosides exhibit a striking conformational difference, presumably responsible for their different biological activity. Indeed, the hexitol nucleosides **1**,¹ possessing antiviral activity, exist predominantly in the chair-like C1 conformation with the base moiety in an axial position, mimicking the C3'-endo form of a furanose nucleoside. However, their carbocyclic congeners **2** (R = H),² which adopt the opposite 1C chair conformation with

the base moiety in an equatorial position, are devoid of antiviral activity. Introduction of an additional α -oriented hydroxyl group (**2**, R = OH) failed to force the molecule into the desired C1 conformation. In order to facilitate interconversion between conformations in a biological system, which seems to be crucial for biological activity, we synthesized the cyclohexene nucleosides **3** in which a double bond is introduced in the cyclohexyl ring. The energy difference between the two pseudo-chair conformers of **3** is expected to be lower than in the cyclohexane nucleosides **2**. An enantioselective approach to the synthesis of optically pure **3** starting from (*R*)-carvone is presented.



Reagents: i. 9-BBN, THF; ii. TBSCl (1eq), imidazole, DMF; iii. MsCl, Et₃N, CH₂Cl₂; iv. Pd-C(10%), HCOONH₄, MeOH; v. MnO₂, CH₂Cl₂; vi. NaBH₄, CeCl₃·7H₂O, MeOH; vii. (a) adenine, PPh₃, DEAD, dioxane, r.t., 66% and 17% *N*₇-isomer; (b) TBAF, THF.

According to NMR analysis and molecular modeling, **3** exists predominantly in the C3'-endo conformation, orienting the adenine base in a pseudo-axial position.



REFERENCES

1. (a) Verheggen, I.; Van Aerschot, A.; Toppet, S.; Snoeck, R.; Janssen, G.; Balzarini, J.; De Clercq, E.; Herdewijn, P. *J. Med. Chem.* **1993**, *36*, 2033-2040; (b) Verheggen, I.; Van Aerschot, A.; Van Meervelt, L.; Rozenski, J.; Wiebe, L.; Snoeck, R.; Andrei, G.; Claes, P.; De Clercq, E.; Herdewijn, P. *J. Med. Chem.* **1995**, *38*, 826-835.
2. Maurinsh, Y.; Schraml, J.; De Winter H.; Blaton, N.; Peeters, O.; Lescrinier, E.; Rozenski, J.; Van Aerschot, A.; De Clercq, E.; Busson, R.; Herdewijn P. *J. Org. Chem.* **1997**, *62*, 2861-2871.
3. Wang, J.; Busson, R.; Blaton, N.; Rozenski, J.; Herdewijn, P. *J. Org. Chem.* **1998**, *63*, 3051-3058.