

Synthesis of a Potent Antimalarial Amphilectene

Sergey V. Pronin and Ryan A. Shenvi*

Department of Chemistry, The Scripps Research Institute, 10550 North Torrey Pines Road, La Jolla, California 92037, United States

Supporting Information

ABSTRACT: 7-Isocyano-11(20),14-epiamphilectadiene, the most potent of antimalarial amphilectenes, is synthesized in seven steps from readily available materials. The synthesis is enabled by a new dendrimeric triene (Danishefsky [3]-dendralene) and a new method for stereo- and chemoselective isocyanation. This chemistry provides a useful entry into an underexplored yet promising family of antimalarial terpenoids.

In 2009, clinicians working along the Thai–Cambodia border confirmed reports of malaria infections that showed decreased susceptibility to artemisinin-combination therapies (ACT), which are the standard of care for treating malignant malaria (*P. falciparum*).¹ The incidence of artemisinin resistance is on the rise,² with some communities reporting reduced ACT-sensitivity in 50% of the patient population.^{3b} Consequently, there is an urgent need to identify, produce, and analyze promising new drugs for the treatment of malaria.³ The isocyanoamphilectenes (e.g., 1–3, Figure 1) are marine




Figure 1. Antimalarial amphilectenes. ^a IC_{50} (*P. falciparum* K1); ^bS.I. = IC_{50} (human fibroblast)/^a; ^c IC_{50} (*P. falciparum* W2); ^dS.I. = IC_{50} (human fibroblast)/^c.

diterpenes⁴ which have demonstrated antimalarial activity and selectivity comparable to the broadly prescribed medicines chloroquine and mefloquine.⁵ Notably, 3 is highly active ($IC_{50} = 31 \text{ nM}$) toward chloroquine-resistant (W2) *P. falciparum*. Unfortunately, isolation from tropical sponges over the course of three decades has not proved a viable means of procuring the mono- or diisocyanoamphilectenes in large quantity,⁶ and chemical synthesis routes have been lengthy (1 in 27 and 25 steps; and 2 in 34 steps).⁷ Here we demonstrate a concise entry into this isocyanoterpene (ICT) family, the total synthesis of 3, and an unusual stereoinversion of a tertiary alcohol.


Our lab is developing syntheses of medicinally important small molecules from linear polyunsaturated oligomers using controlled polycyclizations.⁸ The global strategy of converting small monomers to linear polyunsaturated oligomers, followed

by selective polycyclization, is a common strategy in secondary metabolite biosynthesis⁹ and benefits from both synthetic efficiency and divergency. We realized that the amphilectene skeleton might be reasonably disconnected by scission into two polyunsaturated chains (Figure 1), one of which resembled a dendrimeric isoprene dimer.


Dendrimeric polyenes, or dendralenes, are attractive Diels–Alder cycloaddition reactants because their initial reaction with a dienophile generates a new diene, which can undergo further cycloaddition to yield multiple rings in one step.¹⁰ This process of cycloaddition and diene transmission can proceed in $n + 1$ iterations, if n is equal to the number of conjugated trienes in the initial dendralene. Unfortunately, dendralenes have seen only limited use in natural product synthesis¹⁰ since substituted variants can exhibit poor chemo-, regio-, and stereoselectivity upon cycloaddition.^{10,11} In contrast, highly polarized, non-dendrimeric dienes, especially the Danishefsky diene and its derivatives, undergo facile and highly regioselective cycloadditions with electron-deficient dienophiles.¹² We reasoned that providing Danishefsky-diene character to a dendralene might expand the utility of these polyenes and provide rapid access to the amphilectene skeleton.

Scheme 1. Synthesis of 4 and Attempted Diels–Alder Cascade

a. Synthesis of a Danishefsky dendralene




b. Attempted dendralene Diels–Alder cascade to access the amphilectene skeleton.



Received: October 13, 2012

Scheme 2. Synthesis of (+)-7-Isocyano-11(20),14-epiamphilectadiene (3)



Danishesky dendralene 4 (Scheme 1a) can be synthesized from commercially available *trans*-4-methoxy-3-buten-2-one by (1) iodination to yield 5;¹³ (2) Negishi coupling of 5 with vinylzinc bromide to yield 6; and (3) silylation of 6 using *tert*-butyldimethylsilyl triflate (TBSOTf) and triethylamine. This process is scalable and reproducible; however organoiodide 5 colors at ambient temperature, which corresponds to a drop in the conversion of Negishi coupling, so 5 must be used fresh.

Initial attempts to apply dendralene 4 to the synthesis of the amphilectene skeleton were frustrated on several fronts (Scheme 1b). First, although unsaturated ketone 7a (X = H) underwent facile and regioselective cycloaddition with 4, treatment of cycloadduct 8a with Yb(OTf)₃ caused rapid and irreversible isomerization to 9. To circumvent this isomerization, we synthesized α -chloro ketone 7b (X = Cl), which also underwent clean cycloaddition, but 10b was too unstable to undergo a second Diels–Alder reaction and rapidly decomposed, even upon storage at –20 °C. The third tactic involved use of the corresponding *tert*-butyl ester 7c (X = CO₂*t*-Bu), which underwent smooth cycloaddition to 8c. Unfortunately, the energy of activation for subsequent cycloaddition exceeded that of thermal hetero-retro-ene (loss of isobutylene, 10c→11), and only 9 was observed after prolonged heating. Photochemical cycloadditions were attempted using 10c, but only strained cyclobutane 12 was formed, which resisted subsequent vinyl cyclobutane rearrangement to our target tricycle.

Eventually, we found that methyl ester 7d serves as a competent dienophile (Scheme 2), but this highly reactive compound lacks the steric bulk of 7c and is subject to decomposition, presumably through oligomerization. Therefore 7d is generated under dilute conditions and then concentrated in the presence of dendralene 4. The utility of this reaction for the total synthesis of 3 is shown in Scheme 2. Cycloaddition between 4 and 7d occurs at ambient temperature, and addition of 5 mol % Yb(OTf)₃ generates the intermediate cross-conjugated enone¹⁴ (see Scheme 1b, 10d), which undergoes a second Diels–Alder cycloaddition upon heating in *o*-dichlorobenzene in a microwave reactor. Tricycle 15 can be isolated in 56% yield (calculated from vinylketone 6), and its diastereomer (probably the *endo*-cycloadduct) is detected in only small quantities (d.r. > 10: 1). The reaction 7d + 4 → 15 can also be conducted in one pot, but the yield is lower (31%) due to the presence of Yb(OTf)₃, which causes competitive decomposition.


Efficient conversion of 15 to amphilectene 3 requires an exact choreography of steps, initiated by immediate excision of the methyl carboxylate.¹⁵ Due to the sensitivity of 15 toward conjugate nucleophilic additions, we utilize a modified Krapcho decarboxylation¹⁶ to remove the superfluous methyl ester, which provides the diketone with high stereoselectivity. A tandem conjugate addition/1,2 addition succeeds due to the intermediacy of a silylenol ether which prevents addition of

methylmagnesium bromide to the C11 ketone. As predicted, the product of this sequence, 16, possesses the required configurations at C1, C7, C8, and C12 for conversion to 3.

Transformation of 16 into 3 involves C11 ketone methylenation and a challenging stereoselective displacement of the C7 tertiary alcohol via *N*-alkylation of cyanide. Activation of both functions in 16 is achieved by addition of trimethylsilylmethyl cerium chloride to the C11 ketone,¹⁷ followed by selective trifluoroacetylation of the C7 tertiary alcohol to yield 17. However, final installation of the equatorial isonitrile pharmacophore required extensive exploration since existing methods were not suited to substrates bearing additional sites of unsaturation. For example, application of tertiary alcohol isocyanation reported by Tada¹⁸ and Kitano¹⁹ led first to alkene formation and then to the slow and indiscriminate addition of isocyanide to alkenes mediated by a Brønsted acid. The conditions of Corey and Magriotis for isocyanation were reported to exhibit low stereoselectivity,²⁰ and we also observed competitive formation of tertiary chlorides and cyanides. Therefore we developed our own method for isocyanation²¹ using model decalin 18 (Table 1).

Table 1. Comparison of Isocyanation Methods

entry		conditions	% 19a,b ^a	19a : 19b ^a
1	18b, 15 equiv. TMSCN, 20 equiv. TiCl ₄ , CH ₂ Cl ₂ , 22 °C		9 ^b	6 : 94
2	18a, 3 equiv. TMSCN, 3 equiv. ZnBr ₂ , CH ₂ Cl ₂ , 22 °C		39 ^c	18 : 82
3	18b, 15 equiv. TMSCN, 80 °C		0 ^d	n/a
4	18b, 15 equiv. TMSCN, 3 mol% Sc(OTf) ₃ , 22 °C		86 ^e	88 : 12



^aAccording to GC. ^bMajor byproducts were alkyl chlorides. ^cMajor byproducts were alternative isomeric isonitriles ^dProducts were alkenes. ^eIsolated yield of 19a,b from entry 4 is 66%; isolated yield of 19a is 51%.

As shown in entries 1 and 2, the conditions of Corey and Tada, respectively, produce *axial* isonitrile 19b from 18a or 18b, in preference to equatorial isonitrile 19a.²² We interpret these results to indicate the intermediacy of a solvent separated carbocation, which is expected to undergo facile axial attack.²³ Since inversion of configuration at the C7 tertiary trifluoroacetate was desired, we explored displacement conditions²⁴ that would instead induce reaction at a contact ion pair.²⁵

Unfortunately, attempts at thermal solvolysis^{24a} of **18b** provide alkenes exclusively (entry 3). However, ionization of **18b** with a weak, oxophilic Lewis acid,²⁶ Sc(OTf)₃, in the presence of 15 equiv of TMSCN and in the absence of any other solvent provides **19a** with high stereoselectivity and little elimination (entry 4). Notably, the major diastereomer derives from inversion of configuration and suggests either (1) attack of a contact ion pair or (2) departure of the trifluoroacetate coincident with attack by the TMSCN nucleophile (structure **20**). It should be noted that addition of TMSCN to a 4-*tert*-butyl-1-methylcyclohexyl carbocation in dichloromethane is reported to proceed with primarily the opposite selectivity to yield an axial isonitrile.¹⁹ Application of this new method to penultimate intermediate **17** also effects elimination of the C11 alcohol and provides **3** in good yield and high diastereoselectivity.

In conclusion, we have disclosed a short synthesis of **3**, which is a low nanomolar antimalarial agent and the most potent of the amphilectene isocyanoterpenes. The synthesis of **3** relies on a new dendrimeric variant of the Danishefsky diene which allows rapid assembly of polycycles from linear unsaturated polyenes. Furthermore, we have uncovered a potentially useful, chemoselective inversion of tertiary alcohols with TMSCN; this reactivity is currently being explored. We anticipate that the approach reported herein will be applicable to the asymmetric synthesis of the entire amphilectene family,²⁷ related isocyanoterpenes, and easily derived analogs which could constitute a new arsenal of medicines in the fight against malaria.

■ ASSOCIATED CONTENT

■ Supporting Information

Experimental procedures and spectroscopic data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

rshenvi@scripps.edu

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was supported by an ADI grant from TSRI and a very generous donation of equipment and supplies from Exelixis. We thank the Baran lab for use of their microwave reactor and the Yu lab for use of their LCMS.

■ REFERENCES

- (1) Dondorp, A. M.; Nosten, F.; Yi, P.; Das, D.; Phyto, A. P.; Tarning, J.; Lwin, K. M.; Ariey, F.; Hanpithakpong, W.; Lee, S. J.; Ringwald, P.; Silamut, K.; Imwong, M.; Chotivanich, K.; Lim, P.; Herdman, T.; An, S. S.; Yeung, S.; Singhasivanon, P.; Day, N. P.; Lindegardh, N.; Socheat, D.; White, N. J. *N. Engl. J. Med.* **2009**, *361*, 455.
- (2) Carrara, V. I.; Zwang, J.; Ashley, E. A.; Price, R. N.; Stepniewska, K.; Barends, M.; Brockman, A.; Anderson, T.; McGready, R.; Phaiphun, L.; Proux, S.; van Vugt, M.; Hutagalung, R.; Lwin, K. M.; Phyto, A. P.; Preechapornkul, P.; Imwong, M.; Pukrittayakamee, S.; Singhasivanon, P.; White, N. J.; Nosten, F. *PLoS ONE* **2009**, *4*, e4551.
- (3) (a) Enserink, M. *Science* **2010**, *328*, 844. (b) Eisenstein, M. *Nature* **2012**, *484*, S16.
- (4) Wratten, S. F.; Faulkner, D. J.; Hiratsu, K.; Clardy, J. *Tetrahedron Lett.* **1978**, *20*, 4345.

- (5) (a) König, G. M.; Wright, A. D.; Angerhofer, C. K. *J. Org. Chem.* **1996**, *61*, 3259. (b) Wattanapiromsakul, C.; Chanthathamrongsi, N.; Bussarawit, S.; Yuenyongsawad, S.; Plubrukarn, A.; Suwanborirux, K. *Can. J. Chem.* **2009**, *87*, 612.
- (6) Limited material precluded bioassay is several cases: Wright, A. D.; McCluskey, A.; Robertson, M. J.; MacGregor, K. A.; Gordon, C. P.; Guenther, J. *Org. Biomol. Chem.* **2011**, *9*, 400.
- (7) For **1**, see: (a) Piers, E.; Llinas-Brunet, M. *J. Org. Chem.* **1989**, *54*, 1483. (b) Piers, E.; Llinas-Brunet, M.; Oballa, R. M. *Can. J. Chem.* **1993**, *71*, 1484. For **2**, see: (c) Miyaoka, H.; Okubo, Y. *Synlett* **2011**, *4*, 547. For a 26 step ICT synthesis, see also: (d) Piers, E.; Romero, M. A. *Tetrahedron* **1993**, *49*, 5791.
- (8) (a) Pronin, S. V.; Tabor, M. G.; Jansen, D. J.; Shenvi, R. A. *J. Am. Chem. Soc.* **2012**, *134*, 2012. (b) Pronin, S. V.; Shenvi, R. A. *Nat. Chem.* **2012**, *4*, 915.
- (9) Dewick, P. M. *Medicinal Natural Products: A Biosynthetic Approach*; Wiley: West Sussex, U.K., 2009.
- (10) For a review, see: Hopf, H.; Sherburn, M. S. *Angew. Chem., Int. Ed.* **2012**, *51*, 2298.
- (11) For example: (a) Bradford, T. A.; Payne, A. D.; Willis, A. C.; Paddon-Row, M. N.; Sherburn, M. S. *Org. Lett.* **2007**, *9*, 4861. Cyclic dienophiles are uniquely stereoselective, see: (b) Kwon, O.; Park, S. B.; Schreiber, S. L. *J. Am. Chem. Soc.* **2002**, *124*, 13402.
- (12) (a) Danishefsky, S. J. *Acc. Chem. Res.* **1981**, *14*, 400. (b) Danishefsky, S. J. *Aldrichimica Acta* **1986**, *59*.
- (13) Kozmin, S. A.; Iwama, T.; Huang, Y.; Rawal, V. H. *J. Am. Chem. Soc.* **2002**, *124*, 4628.
- (14) Inokuchi, T.; Okano, M.; Miyamoto, T.; Madon, H. B.; Takagi, M. *Synlett* **2000**, 1549.
- (15) There is another low, near-equal-energy configuration available to the amphilectene tricycle **16** by inversion of stereochemistry at both the C8 and C12 bridgehead carbons. An alternative choreography of steps (i.e., initial conjugate addition to **15**) yields this alternative stereoisomer.
- (16) Hayakawa, I.; Asuma, Y.; Ohyoshi, T.; Aoki, K.; Kigoshi, H. *Tetrahedron Lett.* **2007**, *48*, 6221.
- (17) Johnson, C. R.; Tait, B. D. *J. Org. Chem.* **1987**, *52*, 281.
- (18) Kitano, Y.; Chiba, K.; Tada, M. *Tetrahedron Lett.* **1998**, *39*, 1911.
- (19) Okada, I.; Kitano, Y. *Synthesis* **2011**, *24*, 3997.
- (20) Corey, E. J.; Magriotis, P. *J. Am. Chem. Soc.* **1987**, *109*, 287.
- (21) The method of Mukaiyama was not successful in our hands. See: Masutani, K.; Minowa, T.; Hagiwara, Y.; Mukaiyama, T. *Bull. Chem. Soc. Jpn.* **2006**, *79*, 1106.
- (22) Assigned by ¹³C NMR: Crews, P.; Naylor, S.; Hanke, F. J.; Hogue, E. R.; Kho, E.; Braslau, R. *J. Org. Chem.* **1984**, *49*, 1371.
- (23) Elakovitch, S. D.; Traynham, J. G. *J. Org. Chem.* **1973**, *38*, 873.
- (24) (a) Hughes, E. D.; Ingold, C. K.; Martin, R. J. L.; Meigh, D. F. *Nature* **1950**, *166*, 679. (b) Doering, W. v. E.; Zeiss, H. H. *J. Am. Chem. Soc.* **1953**, *75*, 4733.
- (25) Winstein, S.; Clippinger, E.; Fainberg, A. H.; Heck, R.; Robinson, G. C. *J. Am. Chem. Soc.* **1956**, *78*, 328.
- (26) Ferraris, D.; Drury, W. J., III; Cox, C.; Lectka, T. *J. Org. Chem.* **1998**, *63*, 4568.
- (27) For examples of asymmetric Danishefsky Diels–Alder reactions with alkene dienophiles, see: Sudo, Y.; Shirasaki, D.; Harada, S.; Nishida, A. *J. Am. Chem. Soc.* **2008**, *130*, 12588. Hiraoka, S.; Harada, S.; Nishida, A. *J. Org. Chem.* **2010**, *75*, 3871.