

Organic & Biomolecular Chemistry

This article is part of the

OBC 10th anniversary
themed issue

All articles in this issue will be gathered together
online at

www.rsc.org/OBC10



Cite this: *Org. Biomol. Chem.*, 2012, **10**, 5811

www.rsc.org/obc

COMMUNICATION

Synthesis of 1,2,4-triazines and the triazinoisoquinolinedione DEF ring system of noelaquinone†‡§

Liming Cao, John P. Maciejewski, Stephan Elzner, David Amantini and Peter Wipf*


Received 20th February 2012, Accepted 19th March 2012

DOI: 10.1039/c2ob25353d


The intramolecular Staudinger-aza-Wittig reaction is used for a general synthesis of 1,2,5,6-tetrahydro-1,2,4-triazines, a structural motif reported for the natural product noelaquinone. The DEF moiety of noelaquinone was obtained in 13 steps and 2% overall yield, and the structure of the synthetic product was confirmed by X-ray analysis.

Noelaquinone **1** (Fig. 1) was isolated in 1998 by Paul Scheuer and coworkers from the Indonesian *Xestospongia* sp.¹ The structure assignment of this marine metabolite was mainly based on NMR analyses, and its biological activities have not yet been explored. However, the hexacyclic noelaquinone ring system is closely related to halenaquinone, xestoquinone, and the viridin class of furanosteroids, many of which have shown potent kinase inhibitory effects.^{2,3} A distinct feature of noelaquinone is its DEF 3,4-dihydro-2*H*-[1,2,4]triazino[2,3-*b*]isoquinoline-6,11-dione, a heterocycle that has hitherto never been synthesized.

The absence of any information on its biological profile and the presence of a unique heterocyclic ring system make noelaquinone an attractive target for total synthesis. Herein, we report a new method for the formation of fused 1,2,4-triazines and



Scheme 1 Retrosynthesis of 1,2,5,6-tetrahydro-1,2,4-triazines.

Scheme 2 Preparation of triazino-isoindolones from *N*-aminophthalimide.

its application to the first preparation of the DEF ring system in **1**.

As a key step for the expedient formation of 1,2,4-triazines, we planned to develop a variant of the intramolecular Staudinger-aza-Wittig reaction. This method has proven to be a useful tool in natural products synthesis,⁴ and we decided to further explore its application for the ring closure of an azido-hydrazone precursor (Scheme 1).

Due to its ready availability and high chemical reactivity, we first used *N*-aminophthalimide **3** as a model system (Scheme 2). Azide **5** was obtained in three steps and 47% yield from **3** by condensation with α -bromoacetaldehyde **4**⁵ in the presence of 4 Å molecular sieves to give the α -bromohydrazone. Since this intermediate was found to be very labile, it was reduced without isolation with NaCNBH₃ to give the corresponding bromoethyl derivative, which was converted to alkyl azide **5** with sodium azide in DMF.





Fig. 1 Polycyclic furans isolated from the marine sponge *Xestospongia*.

Department of Chemistry and Center for Chemical Methodologies and Library Development, 219 Parkman Avenue, 15260 Pittsburgh, PA, USA. E-mail: pwipf@pitt.edu; Fax: +1 412 624 0787; Tel: +1 412 624 8606


† This article is part of the *Organic & Biomolecular Chemistry* 10th Anniversary issue.

‡ Dedicated to the memory of Prof. Robert E. Ireland.

§ Electronic supplementary information (ESI) available: Synthetic protocols and spectroscopic data. CCDC 868588–868590. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c2ob25353d




Scheme 3 Failed hydrazone formation.

Scheme 4 Preparation of homophthalimide **18** by a key copper-catalyzed C-arylation.

When azide **5** was heated in toluene at 100 °C in the presence of 2 equiv of PBu_3 , a clean conversion to the desired triazinone **6** occurred. Alternatively, the hydrazide was *N*-protected with benzyl bromide in the presence of tetrabutylammonium hydrogen sulfate (TBAHS), and the phosphine treatment afforded the corresponding triazinone **7** in 97% yield. The *N*-benzylated triazinone **7** was much easier to isolate and purify than the unprotected **6**.

The successful conversion of model azido-hydrazide **5** into triazino[3,2-*a*]isoindolones **6** and **7** encouraged us to explore the preparation of triazino[2,3-*b*]isoquinolines. However, we were unable to isolate the requisite intermediate α -bromohydrazone **9** from hydrazide **8** under typical condensation conditions with aldehyde **4** (Scheme 3). While the hydrazone appeared to form, decomposition during the reaction resulted in the formation of a complex mixture at workup.

After several unsuccessful attempts to add the azide moiety to the homophthalimide ring system, we were able to develop a strategy that installed the cyclic imide on a prefunctionalized hydrazide (Scheme 4). Protection of 2-hydroxyethylhydrazine with ethyl pyruvate **10** gave hydrazone **11** in 59% yield. Mesylation of the alcohol and treatment with NaN_3 led to azide **12**, which was selectively benzylated and then hydrolyzed to give the azido-hydrazine **14**. After acylation of **14** with *o*-iodobenzoylchloride **15**, a C-arylation⁶ of diethylmalonate **17** in the



Scheme 5 Formation of isomeric triazinoisoquinolines under acidic acetal hydrolysis conditions.

presence of 5 mol% of Cu(I)-catalyst, 10 mol% of picolinic acid, and 3 equiv of Cs_2CO_3 in dioxane led to the α -aryl malonate intermediate, which was used crude after workup. The homophthalimide ring was readily obtained by treatment with catalytic *p*-toluenesulfonic acid (*p*-TSA) in aqueous toluene, which promoted a cascade cyclization–decarboxylation to yield **18** in 73% over 2 steps.

In preparation for the introduction of the benzylic ketone function, **18** was converted to thioacetal **20** by treatment with trimethylene dithiosylate **19** (Scheme 5).⁷ Acetal exchange to the dimethyl acetal **21** was accomplished with iodobenzenebis-trifluoroacetate (PIFA) in methanol in the presence of trifluoroacetic acid.⁸ The Staudinger-aza-Wittig reaction of **21** proceeded smoothly under microwave irradiation conditions to give the desired triazine **22** in 61% yield, but all attempts to hydrolyze the acetal to unmask the carbonyl group either decomposed the product or provided a 2.5 : 1 mixture of the isomeric triazinoisoquinolines **23a** and **23b** in modest 45% combined yield. The structures of **23a** and **23b** were unambiguously assigned by X-ray crystallographic analysis (Fig. 2).

While the formation of **23b** accomplished our goal to access the DEF ring system of **1**, we were unable to identify acetal hydrolysis conditions avoiding the equilibration of this heterocyclic system to the isomeric 3,4-dihydro-2*H*-[1,2,4]triazino-[4,3-*b*]isoquinoline-6,11-dione **23a**. Furthermore, we were unable to cleave the benzyl protecting group in either **23a** or **23b**.

Accordingly, we modified the synthetic route shown in Schemes 4 and 5 by using the *p*-methoxybenzyl (PMB) protected azido hydrazine **24** (Scheme 6). Acylation of **15**, C-arylation with malonate **17**, and cyclization–decarboxylation proceeded uneventfully to give homophthalate **25**. In two steps, the benzylic methylene group in **25** was converted to the dimethyl acetal and the Staudinger-aza-Wittig reaction provided triazine **26** in high yield. Interestingly, controlled hydrolysis in cold sulfuric acid removed the PMB group in 72% yield without cleavage of the dimethyl acetal. The resulting dimethyl acetal **27**





Fig. 2 X-Ray structures of **23a** (top) and **23b** (bottom).



Scheme 6 Preparation of the DEF ring system of noelaquinone from the PMB-protected azido hydrazine **24**.

was subjected to hydrolysis in conc. H₂SO₄ at room temperature to yield the desired unprotected triazino[2,3-*b*]isoquinoline **28** in the absence of any rearrangement products. The difference in the reactivities of **22** and **27** in the acid catalyzed hydrolysis is quite striking. We speculate that the increased solvation of the less lipophilic **27** increases the stability of the intermediate oxocarbenium ion, thus allowing for a quicker acetal cleavage at slightly lower temperature and preventing the triazine isomerization. The structural assignment of **28** was confirmed by X-ray analysis (Fig. 3).




Fig. 3 X-Ray structure of **28**.

Conclusions

We have demonstrated an efficient method to generate azido hydrazines, thus enabling the application of the Staudinger-aza-Wittig reaction of imides as a general method for the preparation of tetrahydro-1,2,4-triazines. For the synthesis of the DEF ring system of noelaquinone, the Cu(I)-catalyzed C-arylation of diethyl malonate was used to access the key intermediate homophthalimides **18** and **25**. The use of the PMB protective group and controlled acidic hydrolysis conditions led to the first preparation of the heterocycle **28**, previously reported for the DEF ring moiety of the natural product noelaquinone. In the course of this synthesis, we also observed the equilibration of triazinoisoquinolines **23a** and **23b** under acidic conditions, most likely via an ANRORC⁹ mechanism.

Acknowledgements

The authors thank the National Institutes of Health (P50GM067082) for support of this research and Dr. Steven Geib (University of Pittsburgh) for X-ray analyses.

Notes and references

- Y. Zhu, W. Y. Yoshida, M. Kelly-Borges and P. J. Scheuer, *Heterocycles*, 1998, **49**, 355.
- (a) P. Wipf and R. J. Halter, *Org. Biomol. Chem.*, 2005, **3**, 2053; (b) B. Wakefield, R. J. Halter and P. Wipf, *Org. Lett.*, 2007, **9**, 3121; (c) Y. Lang, F. E. S. Souza, X. Xu, N. J. Taylor, A. Assoud and R. Rodrigo, *J. Org. Chem.*, 2009, **74**, 5429; (d) A. D. Findlay, A. Gebert, I. A. Cade and M. G. Banwell, *Aust. J. Chem.*, 2009, **62**, 1173; (e) K. M. Muller and B. A. Keay, *Synlett*, 2008, 1236; (f) E. H. Sessions and P. A. Jacobi, *Org. Lett.*, 2006, **8**, 4125; (g) P. Wipf, D. J. Minion, R. J. Halter, M. I. Berggren, C. B. Ho, G. G. Chiang, L. Kirkpatrick, R. Abraham and G. Powis, *Org. Biomol. Chem.*, 2004, **2**, 1911; (h) N. T. Ihle, R. Williams, S. Chow, W. Chew, G. Paine-Murrieta, D. J. Minion, R. J. Halter, P. Wipf, R. Abraham, L. Kirkpatrick and G. Powis, *Mol. Cancer Ther.*, 2004, **3**, 763.
- (a) J. R. Hanson, *Nat. Prod. Rep.*, 1995, **12**, 381; (b) A. Longeon, B. R. Copp, M. Roue, J. Dubois, A. Valentin, S. Petek, C. Debitus and M.-L. Bourguet-Kondracki, *Bioorg. Med. Chem.*, 2010, **18**, 6006.
- (a) S. Eguchi, *ARKIVOC*, 2005, **ii**, 98; (b) P. H. Lambert, M. Vaultier and R. Carrie, *J. Org. Chem.*, 1985, **50**, 5352; (c) H. Takeuchi and S. Eguchi, *Tetrahedron Lett.*, 1989, **30**, 3313; (d) J. Chen and C. J. Forsyth, *Org. Lett.*, 2003, **5**, 1281–1283; (e) S. Bräse, C. Gil, K. Knepper and V. Zimmermann, *Angew. Chem., Int. Ed.*, 2005, **44**,

- 5188; (f) Z.-B. Chen, D. Hong and Y.-G. Wang, *J. Org. Chem.*, 2009, **74**, 903; (g) A. Tarraga, P. Molina, D. Curiel and M. D. Velasco, *Tetrahedron*, 2001, **57**, 6765; (h) R. Brossmer and H. Mack, *Tetrahedron Lett.*, 1981, **22**, 933.
- 5 M. Jachak, M. Mittelbach and H. Junek, *Org. Prep. Proced. Int.*, 1993, **25**, 469.
- 6 (a) X. Xie, G. Cai and D. Ma, *Org. Lett.*, 2005, **7**, 4693; (b) S. F. Yip, H. Y. Cheung, Z. Zhou and F. Y. Kwong, *Org. Lett.*, 2007, **9**, 3469.
- 7 R. B. Woodward, I. J. Pachter and M. K. Scheinbaum, *Org. Synth.*, 1974, **54**, 39.
- 8 G. Stork and K. Zhao, *Tetrahedron Lett.*, 1989, **30**, 287.
- 9 H. C. van der Plas, *Adv. Heterocycl. Chem.*, 1999, **74**, 1.